

HETEROCYCLES, Vol. 84, No. 2, 2012, pp. 1227 - 1233. © 2012 The Japan Institute of Heterocyclic Chemistry
 Received, 26th April, 2011, Accepted, 23rd May, 2011, Published online, 3rd June, 2011
 DOI: 10.3987/COM-11-S(P)14

FORMAL SYNTHESIS OF (-)-CLAVEPICTINE A AND (+)-CLAVEPICTINE B FROM A SULFINIMINE (*N*-SULFINYLIMINE)-DERIVED CHIRAL BUILDING BLOCK[#]

Franklin A. Davis* and He Xu

Department of Chemistry, Temple University, Philadelphia, Pennsylvania 19122
 U. S. A.

fdavis@temple.edu

Abstract – The sulfinimine-derived chiral building block *trans*-2,6-disubstituted 1,2,5,6-tetrahydropiperidine (+)-**5** was employed in a concise formal asymmetric synthesis of the cytotoxic marine alkaloids clavepictines (-)-A and (+)-B (**1a** and **1b**). This synthesis is highlighted by a highly diastereoselective hydroboration-oxidation reaction to install the C-3 hydroxyl group.

Clavepictines A and B (**1a** and **1b**) are quinolizidine alkaloids isolated from the marine Bermudan tunicate *Clavelina picta*.^{1,2} These alkaloids have an unusual *cis*-ring fused quinolizidine skeleton where the methyl and acetoxy (hydroxyl) groups occupy axial positions.² Antifungal, antimicrobial, and antitumor activity have been attributed to these compounds.¹ Couty and co-workers, in a structure-reactivity relationship (SAR) study concluded that their cytotoxicity is primarily determined by the side chain at C-10.³





Figure 1. Structures of Clavepictine A and B

Three asymmetric total syntheses of clavepictines A and B have been described.^{4,6} All of these syntheses

[#]Dedicated to Professor Albert Padwa on the occasion of his 75th birthday and in recognition of his pioneering contributions to the art of synthesis.

utilized an intramolecular Michael-type cyclization of a functionalized piperidine to prepare the quinolizidine ring. In each case reduction of an intermediate acyl iminium ion species, generated from an *N*-acyl enamine was used to form the core *trans*-2,6-disubstituted piperidine ring. Momose and co-workers, in the first total synthesis of (+)-**1a** and (+)-**1b**, prepared the enamine from a lactone diol,⁴ while the Ma group used an enamine sulfone.⁵ Ha and Cha prepared the *N*-acyl enamine via a Comins' cross-coupling of an enamide triflate.⁶ This group also generated the *trans*-2,6-disubstituted piperidine ring using Beak's α -lithiation-substitution chemistry.


Recently we introduced a new method for the asymmetric synthesis of *trans*-2,6-disubstituted 1,2,5,6-tetrahydropiperidines such as (+)-**5**. These unsaturated heterocycles are valuable building blocks for the asymmetric synthesis of polysubstituted piperidines because of the many methods available for carbon-carbon double bond functionalization (Scheme 1).^{7,8} Our procedure involves a one-pot five-step cascade reaction of a sulfinimine-derived *N*-sulfinyl- δ -amino- β -ketophosphonate (+)-**2** to a 3-phosphoryl dihydropyridone (+)-**3** via an intermediate enamino. ^{7,9,10} Addition of methyl cuprate to (+)-**3** gave the *trans*-piperidine (+)-**4** which was transformed into 1,2,5,6-tetrahydropiperidine (+)-**5**. Compound (+)-**5** was employed in the first total synthesis of (-)-myrtine (**6**) the unnatural isomer of the alkaloid (+)-myrtine. We describe here a formal synthesis of clavepictines A and B (**1a** and **1b**) from building block (+)-**5**.



Scheme 1

It was decided first to form the quinolizidine ring and introduce the C-3 hydroxy group later. The benzyl protecting group in (+)-**5** was removed by treatment with sodium in liquid NH_3 at -78 °C affording


tetrahydropyridine alcohol (2*S*,6*S*)-(+)-**7** in 90% yield (Scheme 2). Alcohol (+)-**7** was oxidized with 1.5 equiv of Dess-Martin periodinane reagent to give an intermediate aldehyde **8**, which was not purified, but was used directly in a Horner-Wadsworth-Emmons reaction with 1.1 equiv of diethyl phenylsulfonylethylphosphonate¹¹ and 1.1 equivalence of NaH, affording the piperidine vinyl sulfone (2*S*,6*S*)-(+)-**9** in 66% yield over the two steps (Scheme 2). Attempts to remove the *N*-Boc group in (+)-**9** with TMS-I were unsuccessful leading to unidentified products. However, with excess TFA in CH₂Cl₂ (+)-**9** gave a salt **10** that was treated with 10 equiv of Cs₂CO₃ in MeOH to give quinolizidine (+)-**11** in 74% yield along with its minor isomer (Scheme 2). The intramolecular Michael cyclization gave the desired stereochemistry at C-10 as determined by X-ray crystallography as shown in Figure 2 and by its conversion to a compound of known absolute configuration (see below).



Scheme 2

To install the C-3 hydroxyl group epoxidation of the double bond in (+)-**11** was next considered. However, the basic nitrogen atom in (+)-**11** and the anticipated problems in the regioselective ring opening of the epoxide led us to abandon this approach. In a study of the hydroboration of unsaturated piperidines Lyle and coworkers have reported that the direction of borane addition is strongly influenced by the positive charge on nitrogen; i.e. transition state **TS-A** should be favored over transition state **TS-B** (Scheme 3).¹² Indeed hydroboration-oxidation of (+)-**11** gave the desired alcohol (-)-**12** in 50% isolated yield as a single isomer. Conversion of the alcohol into the MOM ether with methyl chloromethyl ether afforded (-)-**13** in 82% yield (Scheme 3). Quinolizidine (-)-**13** is identical in all respect to that prepared in Momose's synthesis of the clavicipitines.⁴ The synthesis of quinolizidine (-)-**13** represents a formal

asymmetric synthesis of clavепictines (-)-**1a** and (+)-**1b** and confirms the stereochemical assignments for (+)-**11** and (-)-**12**.



Scheme 3

In summary, a new formal synthesis of the cytotoxic marine alkaloids Clavепictines (-)-A and (+)-B (**1a** and **1b**) is described from sulfinimine-derived building block *trans*-2,6-disubstituted 1,2,5,6-tetrahydropiperidins (+)-**5**. The synthesis is highlighted by a highly diastereoselective hydroboration-oxidation reaction to install the C-3 hydroxyl group.

ACKNOWLEDGEMENT

This work was supported by grants from the National Institutes of General Medical Sciences (GM57870 and GM51982).

EXPERIMENTAL

General procedure. All reagents were used as received unless otherwise noted. Tetrahydrofuran (THF), diethyl ether (Et₂O), dichloromethane (DCM), and toluene were purified by filtration on a solvent purification system. Unless otherwise mentioned, all reagents were carried under argon atmosphere. Column chromatography was performed on silica gel, 230-400 mesh. TLC plates were visualized with UV, in an iodine chamber, or with phosphomolybdic acid, unless otherwise noted. ¹H NMR and ¹³C NMR spectra were recorded on 500 and 400 MHz NMR spectrometers. (2*S*,6*S*)-(+)-*tert*-Butyl 6-(benzyloxy)butyl-2-methyl-5,6-dihydropyridine-1-carboxylate (**5**) was prepared as previously described.⁷

(2*S*,6*S*)-(+)-*tert*-Butyl 6-(4-hydroxybutyl)-2-methyl-5,6-dihydropyridine-1 (2*H*)-carboxylate (**7**). In a

50-mL oven-dried three-neck round bottom flask equipped with a magnetic stirring bar, rubber septum, a cold finger, an ammonia gas inlet, and an argon inlet was placed Na metal (0.06 g, 2.56 mmol). The flask was cooled to $-78\text{ }^{\circ}\text{C}$ as the liquid ammonia (10 mL) was collected using the cold finger and reacted with Na to give a dark blue color solution. To the flask was slowly added *t*-BuOH (0.1 g in 1 mL THF) followed by (+)-**5** (0.115 g, 0.32 mmol) in THF (6 mL). The solution was stirred at $-78\text{ }^{\circ}\text{C}$ for 10 min. The dry ice bath was removed and the solution was warmed to rt to evaporate the liquid ammonia before it was quenched with sat. aqueous NH_4Cl (5 mL). The aqueous layer was extracted with EtOAc (2×10 mL). The combined organic phases were washed with H_2O (2×5 mL), brine (5 mL), dried (MgSO_4), and concentrated. Flash chromatography (EtOAc:hexanes:MeOH, 10:90:10) afforded 0.086 g (90%) of a colorless oil; $[\alpha]_{\text{D}}^{20} +50.8$ (*c* 0.85, CHCl_3); IR (neat) 3410, 2213, 1700, 1232, 1169 cm^{-1} ; ^1H NMR (CDCl_3) δ 5.73 (m, 2 H), 4.14 (m, 1 H), 3.98 (m, 1 H), 3.62 (d, $J = 6.4$ Hz, 2 H), 2.28 (m, 1 H), 2.09 (m, 1 H), 1.61-1.22 (m, 6 H), 1.47 (s, 9 H), 1.27 (d, $J = 6.4$ Hz, 3 H); ^{13}C NMR (CDCl_3) δ 155.9, 132.0, 122.6, 79.7, 63.2, 51.9, 48.6, 33.4, 32.9, 28.9, 27.2, 23.6, 21.9. HRMS calcd for $\text{C}_{15}\text{H}_{28}\text{NO}_3$ ($\text{M} + \text{H}$) 270.2069. Found 270.2068.

(2*S*,6*S*)-(+)-*tert*-Butyl 2-methyl-6-(5-(phenylsulfonyl)pent-4-enyl)-5,6-dihydropyridine-1-carboxylate (9). In a 25-mL, one-necked, round-bottom flask fitted with magnetic stirring bar, rubber septum, and argon inlet was placed (+)-**7** (0.067 g, 0.249 mmol) in CH_2Cl_2 (8 mL). To the flask was added Dess-Martin periodinane (1.06 mL, 15% in CH_2Cl_2) and the solution was stirred for 2 h. The reaction mixture was quenched with aqueous Na_2SO_3 (5 mL) and stirred for 10 min until the solution was clear. The mixture was then extracted with Et_2O (3×10 mL) and the combined organic phases were washed with sat. aqueous NaHCO_3 (2×5 mL), brine (5 mL), dried (MgSO_4), and concentrated to give a yellow residue of the aldehyde **8**. The crude aldehyde was used in the next step without purification. In another 25-mL one-necked, round-bottom flask fitted with magnetic stirring bar, rubber septum, and argon inlet was placed NaH (0.01 g, 0.249 mmol, 60% in mineral oil), which was washed with petroleum ether (3×5 mL). To this flask was added $(\text{EtO})_2\text{POCH}_2\text{SO}_2\text{Ph}^{11}$ (0.073 g, 0.249 mmol) in THF (10 mL) followed by the crude aldehyde. The mixture was stirred for 30 min and quenched with sat. aqueous NH_4Cl solution (5 mL). The phases were separated, the aqueous phase was extracted with EtOAc (2×10 mL) and the combined organic phase were washed with H_2O (5 mL), brine (5 mL), dried (MgSO_4), and concentrated. Flash chromatography (EtOAc:hexanes, 20:80) afforded 0.077 g (76%) of a colorless oil; $[\alpha]_{\text{D}}^{20} +45.7$ (*c* 1.5, CHCl_3); IR (neat) 2360, 1684, 1365, 1147 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.86 (m, 1 H), 7.84 (m, 1 H), 7.59 (m, 1 H), 7.52 (m, 2 H), 6.96 (m, 1 H), 6.30 (m, 1 H), 5.71 (m, 2 H), 4.09 (m, 1 H), 3.98 (m, 1 H), 2.31 (m, 1 H), 2.23 (m, 2 H), 1.50-1.38 (m, 4 H), 1.44 (s, 9 H), 1.26 (d, $J = 6.4$ Hz, 3 H); ^{13}C NMR (CDCl_3) δ 155.8, 147.1, 141.1, 133.5, 132.1, 131.0, 129.6, 127.9, 122.4, 79.8, 33.4, 31.7, 28.9,

27.3, 25.5, 22.0. HRMS calcd for C₂₂H₃₁NO₄NaS (M + Na) 428.1872. Found 428.1859.

(4S,6S,9aS)-(+)-4-Methyl-6-(phenylsulfonylmethyl)-4,6,7,8,9,9a-hexahydro-1H-quinolizine (11). In a 25-mL, one-necked, round-bottom flask fitted with magnetic stirring bar, rubber septum, and argon inlet was placed (+)-**9** (0.31 g, 7.64 mmol) in CH₂Cl₂ (20 mL). To the flask was added TFA (2.84 mL, 38.2 mmol) and the solution was stirred for 50 min. The solvent was removed and the flask was placed under vacuum for 1 h to afford a yellow residue. The residue was dissolved in MeOH (30 mL), Cs₂CO₃ (2.5 g, 76.4 mmol) was added and the reaction mixture was stirred for 30 min. At this time, the mixture was transferred to a pre-mixed solvent of EtOAc (150 mL) and brine (30 mL) in a 500-mL separatory funnel and shaken vigorously for 1 min. The organic phase was washed with brine (2 × 20 mL), dried (MgSO₄), and concentrated. Flash chromatography (EtOAc:hexanes, 50:50) afforded 0.17 g (74%) of a white solid; mp 69-72 °C; [α]_D²⁰ + 51.2 (c 0.45, CHCl₃); IR (neat) 2360, 1303, 1142 cm⁻¹; ¹H NMR (CDCl₃) δ 7.93 (d, *J* = 7.6 Hz, 2 H), 7.61 (d, *J* = 7.2 Hz, 1 H), 7.53 (d, *J* = 7.2 Hz, 2 H), 5.49 (m, 1 H), 5.36 (m, 1 H), 3.58 (m, 2 H), 3.30 (q, *J* = 8.0 Hz, 2 H), 2.81 (m, 1 H), 1.86 (m, 2 H), 1.67 (m, 2 H), 1.54 (m, 2 H), 1.36 (m, 1 H), 1.13 (m, 1 H), 1.00 (d, *J* = 6.4 Hz, 3 H); ¹³C NMR (CDCl₃) δ 140.8, 133.6, 130.4, 129.3, 128.5, 121.8, 59.4, 50.2, 49.0, 48.7, 29.8, 25.8, 23.8, 21.2, 20.8; HRMS calcd for C₁₇H₂₄NO₂S (M + H) 306.1528, found 306.1516.

(3R,4S,6S,9aS)-(-)-4-Methyl-6-(phenylsulfonylmethyl)-octahydro-1H-quinolizine-3-ol (12). In a 25-mL, one-necked, round-bottom flask fitted with magnetic stirring bar, rubber septum, and argon inlet was placed (+)-**11** (0.030 g, 0.0982 mmol) in THF (5 mL) at 0 °C. To the flask was added BH₃·THF (0.393 mL, 0.393 mmol, 1 M in THF) and the solution warmed to rt and stirred for 2 h. At this time, the rubber septum was removed and replaced with a reflux condenser, and 6 N NaOH (0.5 mL) and 30% H₂O₂ (0.1 mL) were added to the solution. The mixture was heated at 50-55 °C for 2 h, cooled to rt, diluted with H₂O (2 mL), and extracted with EtOAc (3 × 10 mL). The combined organic phases were washed with H₂O (5 mL), brine (5 mL), dried (MgSO₄), and concentrated. Flash chromatography (EtOAc:hexanes, 50:50) afforded 0.016 g (50%) of a colorless oil; [α]_D²⁰ -12.5 (c 0.21, CHCl₃); IR (neat) 3520, 1311, 1149 cm⁻¹; ¹H NMR (CDCl₃) δ 7.92 (d, *J* = 7.6 Hz, 2 H), 7.63 (t, *J* = 7.2 Hz, 1 H), 7.54 (t, *J* = 7.2 Hz, 2 H), 3.83 (m, 1 H), 3.70 (m, 1 H), 3.27 (dd, *J* = 14.0, 5.2 Hz, 1 H), 3.02 (m, 1 H), 2.71 (m, 2 H), 2.10-0.76 (m, 13 H); ¹³C NMR (CDCl₃) δ 140.8, 133.7, 129.4, 128.5, 73.1, 58.4, 56.0, 50.1, 49.4, 28.6, 27.9, 24.2, 23.0, 20.5, 15.4. HRMS calcd for C₁₇H₂₆NO₃S (M + H) 324.1633. Found 324.1644.

(3R,4S,6S,9aS)-(-)-3-(Methoxymethoxy)-4-methyl-6-(phenylsulfonyl-methyl)-octahydro-1H-quinolizine (13). In a 25-mL, one-necked, round-bottom flask fitted with magnetic stirring bar, rubber septum, and argon inlet was placed (-)-**12** (0.020 g, 0.0618 mmol) in THF (5 mL). To the flask was added chloromethyl methyl ether (0.019 mL, 0.247 mmol) and DIPEA (0.043 mL, 0.247 mmol). The solution

was stirred for 8 h and the solvent was concentrated to give a residue. Flash chromatography (EtOAc:hexanes, 50:50) afforded 0.017 g (82%) of a colorless oil; $[\alpha]_D^{20}$ -11.0 (*c* 0.30, CHCl₃), [lit.⁴ $[\alpha]_D^{26}$ -10.95 (*c* 0.81, CHCl₃)]; IR (neat) 3058, 1446, 1303, 1148 cm⁻¹; ¹H NMR (CDCl₃) δ 7.92 (d, *J* = 7.6 Hz, 2 H), 7.61 (t, *J* = 7.6 Hz, 1 H), 7.54 (t, *J* = 7.6 Hz, 2 H), 4.70 (d, *J* = 6.8 Hz, 1 H), 4.58 (d, *J* = 6.8 Hz, 1 H), 3.83 (m, 1 H), 3.64 (dd, *J* = 14.0, 6.4 Hz, 1 H), 3.36 (s, 3 H), 3.28 (dd, *J* = 14.4, 5.2 Hz, 1 H), 2.79 (m, 2 H), 2.62 (m, 1 H), 1.84-1.66 (m, 4 H), 1.55 (m, 1 H), 1.37-1.19 (m, 3 H), 1.05 (d, *J* = 5.6 Hz, 3 H), 1.02-0.93 (m, 2 H); ¹³C NMR (CDCl₃) δ 140.6, 131.1, 128.8, 128.1, 95.6, 79.4, 58.2, 55.5, 53.1, 49.2, 48.9, 27.7, 25.8, 22.3, 21.0, 20.4, 15.4. HRMS calcd for C₁₉H₃₀NO₄S (M + H) 368.1896. Found 368.1881. The spectra were consistent with literature values.⁴

REFERENCES

1. M. F. Raub, J. H. Cardellina, M. I. Choudhary, C.-Z. Ni, J. Clardy, and M. C. Alley, *J. Am. Chem. Soc.*, 1991, **113**, 3178.
2. F. Kong and D. J. Faulkner, *Tetrahedron Lett.*, 1991, **32**, 3667.
3. C. Agami, F. Couty, G. Evano, F. Darro, and R. Kiss, *Eur. J. Org. Chem.*, 2003, 2062.
4. N. Toyooka, Y. Yotsui, Y. Yoshida, and T. Momose, *J. Org. Chem.*, 1996, **61**, 4882.
5. S. Yu, X. Pu, T. Cheng, R. Wang, and D. Ma, *Org. Lett.*, 2006, **8**, 3179.
6. (a) J. D. Ha, D. Lee, and J. K. Cha, *J. Org. Chem.*, 1997, **62**, 4550; (b) J. D. Ha and J. K. Cha, *J. Am. Chem. Soc.*, 1999, **121**, 10012.
7. F. A. Davis, H. Xu, and J. Zhang, *J. Org. Chem.*, 2007, **72**, 2046.
8. For reviews see: (a) P. M. Weintraub, J. S. Sabol, J. M. Kane, and D. R. Borcharding, *Tetrahedron*, 2003, **59**, 2953; (b) S. Laschat and T. Dickner, *Synthesis*, 2000, 1781; (c) M. S. M. Pearson, M. Mathe-Allainmat, V. Fargeas, and J. Lebreton, *Eur. J. Org. Chem.*, 2005, 2159; (d) F.-X. Felpin and J. Lebreton, *Current Organic Synthesis*, 2004, **1**, 83; (e) M. G. P. Buffat, *Tetrahedron*, 2004, **60**, 1701.
9. For leading references to the chemistry of *N*-sulfinylimines see: F. A. Davis, *J. Org. Chem.*, 2006, **71**, 8993.
10. For reviews on *N*-sulfinylimines see: (a) D. Morton and R. A. Stockman, *Tetrahedron*, 2006, **62**, 8869; (b) C. H. Senanayake, D. Krishnamurthy, Z.-H. Lu, Z. Han, and I. Gallou, *Aldrichimica Acta*, 2005, **38**, 93; (c) P. Zhou, B.-C. Chen, and F. A. Davis, *Tetrahedron*, 2004, **60**, 8003.
11. T. A. Blumenkopf, *Synth. Commun.*, 1986, **16**, 139.
12. R. E. Lyle, R. R. Carle, C. R. Ellefson, and C. K. Spicer, *J. Org. Chem.*, 1970, **35**, 802.