

HETEROCYCLES, Vol. 86, No. 2, 2012, pp. 1391 - 1400. © 2012 The Japan Institute of Heterocyclic Chemistry  
 Received, 10th July, 2012, Accepted, 17th August, 2012, Published online, 24th August, 2012  
 DOI: 10.3987/COM-12-S(N)95

## Mn(OAc)<sub>3</sub>-BASED $\alpha'$ -OXIDATIVE ACETOXYLATION OF *N*-TRIFLUOROACETYL VINYLOGOUS AMIDES


Hong-Yu Lin and Barry B. Snider\*

Department of Chemistry MS 015, Brandeis University, 415 South Street,  
 Waltham, Massachusetts 02454-9110, USA; E-mail: snider@brandeis.edu

**Abstract** – Vinylogous amides **1** and **11** are oxidized by dried Mn(OAc)<sub>3</sub> in benzene at 90 °C to pyridines **5** and **12**, respectively, whereas the analogous *N*-trifluoroacetyl vinylogous amides **6** and **13** are oxidized by dried Mn(OAc)<sub>3</sub> in benzene at 120 °C to give acetates **7** and **14**. The  $\alpha'$ -oxo radicals that are generated in this oxidation cyclize to proximal double bonds to give tricyclic alkenes such as **20**.


### INTRODUCTION

The lycopodium alkaloid sauroine (7,8-dihydroxylycopodine, **4**) was recently shown to improve memory retention in male Wistar rats making it a significant synthetic target (see Scheme 1).<sup>1</sup> We chose to first address the synthesis of the simpler, related alkaloid 7-hydroxylycopodine (**3**) and recently reported a six step synthesis of **3** from vinylogous amide **1**.<sup>2</sup> For the synthesis of sauroine (**4**) we needed to hydroxylate the vinylogous amide **1** in the  $\alpha'$ -position to obtain **2**, which could then be elaborated to sauroine (**4**) by the route that was used to prepare **3** from **1**.



Scheme 1

Unfortunately, a variety of standard hydroxylation procedures failed to convert **1** to **2** indicating that the chemistry of vinylogous amides such as **1** is quite different from that of enones.<sup>2b</sup> For instance, Watt and Demir developed Mn(OAc)<sub>3</sub>-based  $\alpha'$ -acetoxylation of enones that proceed in high yield with dried Mn(OAc)<sub>3</sub> in benzene at reflux.<sup>3</sup> They found that the use of both Mn(OAc)<sub>3</sub>•2H<sub>2</sub>O that had been dried in vacuum over P<sub>2</sub>O<sub>5</sub> for several days and benzene at reflux as the solvent was crucial for the success of the  $\alpha'$ -acetoxylation. Although only 2 equiv of dried Mn(OAc)<sub>3</sub> are required by the stoichiometry, the reaction is heterogeneous and in practice requires 4-6 equiv of dried Mn(OAc)<sub>3</sub>. This procedure also works well on  $\beta$ -alkoxyenones (vinylogous esters),<sup>3a</sup> but we found that it converted vinylogous amide **1** to pyridine **5** in 65% yield, rather than the desired acetate ester of **2** (see Scheme 2).<sup>2b</sup> Clearly oxidation proceeds at the nitrogen rather than by enolization and formation of the  $\alpha'$ -oxo radical. We thought that acylation of the nitrogen should both prevent this undesired oxidation and favor the formation of the required  $\alpha'$ -oxo radical by increasing the acidity of the  $\alpha'$ -proton.




Scheme 2

## RESULTS AND DISCUSSION

Electron deficient acyl groups should be the most effective at increasing the acidity of the  $\alpha'$ -proton. We therefore treated **1** with trifluoroacetic anhydride and triethylamine in CH<sub>2</sub>Cl<sub>2</sub> for 1 h at 0 °C to generate *N*-trifluoroacetyl vinylogous amide **6** in 95% yield.<sup>4</sup> To our delight, we found that oxidation of **6** with dried Mn(OAc)<sub>3</sub> in benzene in a sealed tube at 120 °C for 30 h cleanly afforded the desired acetate **7** as an inseparable ~1 : 1 mixture of isomers. This reaction is considerably slower than the  $\alpha'$ -acetoxylation of  $\beta$ -alkoxyenones (vinylogous esters), which require 20-48 h in benzene at reflux. The rate determining step is formation of the Mn(III) enolate and the  $\alpha'$ -protons of vinylogous imide **6** are probably less acidic

than those of vinylogous esters. Hydrolysis of both the acetate and trifluoroacetamide of **7** with potassium carbonate in MeOH at 25 °C for 6 h gave 31% of the desired trans alcohol **2** and 28% of the undesired cis alcohol **8**. The stereochemistry of the trans isomer **2** was established by the 11.2 Hz coupling between the diaxial methine hydrogens indicating that both the hydroxyl and methyl groups are equatorial.<sup>5</sup> The coupling constant between the methine hydrogens in the cis isomer **8** is 5.2 Hz.<sup>5</sup> The absence of a large coupling constant between the methine hydrogen adjacent to the methyl group and either adjacent methylene hydrogen indicates that in the preferred conformation of **8** the methyl group is axial and the alcohol is equatorial and probably hydrogen bonded to the adjacent carbonyl group.


Reaction of 3-bromopropylamine hydrobromide (**9**), dimedone (**10**), and 2,6-lutidine in EtOH in a sealed tube at 140 °C for 2.5 h gave the known vinylogous amide **11**<sup>6</sup> in 61% yield (see Scheme 3). As expected from the conversion of **1** to **5**, oxidation of the unprotected vinylogous amide **11** with dried Mn(OAc)<sub>3</sub> in benzene at 90 °C for 20 min afforded pyridine **12**<sup>7</sup> in 65% yield. Trifluoroacetylation of **11** afforded **13** (95%), which was oxidized with dried Mn(OAc)<sub>3</sub> in benzene at 120 °C for 96 h to give the desired acetate **14** in 65% yield. The oxidation of **13** is considerably slower than that of **6**, probably as a result of increased steric hindrance due to the second alkyl substituent. Demir recently reported that  $\alpha'$ -acetoxylation of enones in 9:1 benzene/HOAc is much faster than in benzene.<sup>8</sup>  $\alpha'$ -Acetoxylation of **13** in 9:1 benzene/HOAc is complete in 6 h at 120 °C and 21 h at 90 °C, but is not quite as clean as the slower  $\alpha'$ -acetoxylation in benzene at 120 °C.



**Scheme 3**

In 1996 we reported that cyclization of the  $\alpha'$ -keto radical formed by oxidation of enones with Mn(OAc)<sub>3</sub> to a proximal double bond is faster than acetoxylation.<sup>9</sup> We therefore investigated whether this was also

the case with *N*-trifluoroacetyl vinylogous amides. Reaction of 3-bromopropylamine hydrobromide (**9**), cyclohexanedione **15**,<sup>10</sup> and 2,6-lutidine in EtOH in a sealed tube at 140 °C for 8 h gave vinylogous amide **16** in 52% unoptimized yield (see Scheme 4). Trifluoroacetylation afforded **17** (93%), which was oxidized with dried Mn(OAc)<sub>3</sub> in benzene at 130 °C for 132 h to give the expected tricyclic alkene **20** in 61% yield and tertiary acetate **21** in 5% yield. Oxidation presumably generates  $\alpha'$ -oxo radical **18**, which undergoes the expected 5-exo cyclization with the alkene on the convex face to give *cis* fused tertiary radical **19**, which is oxidized to the cation that loses a proton to yield **20** and reacts with acetate to form **21**. The stereochemistry of the major product **20** was confirmed by the NOEs shown in Scheme 4 and by the coupling pattern of the allylic methine hydrogen (ddd,  $J = 11.2, 10.4, 7.2$  Hz), which is consistent with that expected for this isomer, but not the other three diastereomers. The pyrrolidine-catalyzed intramolecular Michael reaction of ethyl (*E*)-5-(1-methyl-3-oxocyclohexyl)-2-pentenoate gave a single diastereomer corresponding to **20** providing further support for the structure assignment.<sup>11</sup>



Scheme 4

In conclusion, we have shown that vinylogous amides **1** and **11** are oxidized by dried Mn(OAc)<sub>3</sub> in benzene at 90 °C to pyridines **5** and **12**, respectively, whereas the analogous *N*-trifluoroacetyl vinylogous amides **6** and **13** are oxidized by dried Mn(OAc)<sub>3</sub> in benzene at 120 °C to give acetates **7** and **14**. The  $\alpha'$ -oxo radicals such as **18** that are generated in this oxidation cyclize to proximal double bonds to give tricyclic alkenes such as **20**. We are currently exploring the elaboration of hydroxy vinylogous amide **2** to sauroine (**4**).

## EXPERIMENTAL

**General Experimental Methods.** Reactions were conducted in flame- or oven-dried glassware under a nitrogen atmosphere and were stirred magnetically. The phrase "concentrated" refers to removal of solvents by means of a rotary-evaporator attached to a diaphragm pump (15-60 Torr) followed by removal of residual solvents at < 1 Torr with a vacuum pump. Flash chromatography was performed on silica gel 60 (230-400 mesh). Et<sub>3</sub>N, pyridine, MeCN and benzene were distilled from CaH<sub>2</sub>. Mn(OAc)<sub>3</sub>•2H<sub>2</sub>O was dried over P<sub>2</sub>O<sub>5</sub> under vacuum for several days.<sup>3</sup> <sup>1</sup>H and <sup>13</sup>C NMR spectra were obtained on a 400 MHz spectrometer in CDCl<sub>3</sub> with TMS as internal standard unless specifically indicated. Chemical shifts are reported in δ (ppm downfield from tetramethylsilane). Coupling constants are reported in Hz with multiplicities denoted as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), and br (broad). IR spectra were acquired on an FT-IR spectrometer and are reported in wave numbers (cm<sup>-1</sup>). High resolution mass spectra were obtained using electrospray ionization (ESI).

**2,3,4,6,7,8-Hexahydro-7-methyl-1-trifluoroacetyl-5(1H)-quinolinone (6).** Trifluoroacetic anhydride (0.56 mL, 4.0 mmol, 2.0 equiv) and NEt<sub>3</sub> (0.67 mL, 5.0 mmol, 2.5 equiv) were added to a solution of **1** (330 mg, 2.0 mmol) in 6 mL of CH<sub>2</sub>Cl<sub>2</sub> at 0 °C. The resulting mixture was stirred at 0 °C for 1 h. The reaction was quenched by addition of water (12 mL). The mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub> (12 mL × 3). The combined organic layers were washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated. Flash chromatography of the residue on silica gel (6:1 hexanes/EtOAc) gave 496 mg (95%) of **6** as a colorless oil: <sup>1</sup>H NMR 3.90 (br ddd, 1, *J* = 12.8, 3.6, 3.6, H-2), 3.43 (ddd, 1, *J* = 12.8, 10.4, 2.8, H-2), 2.75 (dddd, 1, *J* = 17.6, 10.4, 3.2, 3.2, H-8), 2.65 (br d, 1, *J* = 17.6, H-8), 2.54 (dd, 1, *J* = 12.8, 1.8, H-6), 2.43 (dddd, 1, *J* = 18.4, 6.8, 3.4, 3.4, H-4), 2.30 (dddd, 1, *J* = 18.4, 8.4, 8.4, 2.4, 2.4, H-4), 2.24-2.13 (m, 1, H-7), 2.14 (dd, 1, *J* = 12.8, 12.8, H-6), 2.08-1.98 (m, 1, H-3), 1.90-1.78 (m, 1, H-3), 1.08 (d, 3, *J* = 6.4); <sup>13</sup>C NMR 198.5, 155.8 (q, 1, *J* = 36.4, CF<sub>3</sub>CO), 153.2, 124.6, 115.9 (q, 1, *J* = 287, CF<sub>3</sub>), 45.4, 45.2 (q, 1, *J* = 4.0, CF<sub>3</sub>CONCH<sub>2</sub>), 37.3, 30.6, 22.2, 21.0, 20.1; IR 1702, 1663, 1623 (wk), 1202, 1140, 913, 748.

**trans-2,3,4,6,7,8-Hexahydro-6-hydroxy-7-methyl-5(1H)-quinolinone (2) and cis-2,3,4,6,7,8-Hexahydro-6-hydroxy-7-methyl-5(1H)-quinolinone (8).** A solution of **6** (496 mg, 1.9 mmol) in 20 mL of benzene and dried Mn(OAc)<sub>3</sub> (1.98 g, 8.6 mmol, 4.5 equiv) were added to a resealable tube. The reaction mixture was sealed and heated at 120 °C for 30 h. The reaction mixture was cooled to room temperature and 10% aq. NaHSO<sub>3</sub> solution (50 mL) was added. The mixture was extracted with EtOAc (50 mL × 3). The combined organic layers were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give crude acetate **7** as a ~1:1 mixture of isomers.

To a solution of the residue in 12 mL of MeOH was added powdered K<sub>2</sub>CO<sub>3</sub> (1.05 g, 4 equiv). The reaction mixture was stirred at room temperature for 6 h and concentrated. To the residue was added water (30 mL). The mixture was extracted with EtOAc (30 mL × 4). The combined organic layers were

washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash chromatography of the residue on silica gel (100:1:1 EtOAc/MeOH/ $\text{NEt}_3$ ) gave 106 mg (31% from **6**) of **2** as a white solid, followed by 96 mg (28% from **6**) of **8** as a colorless oil.

Data for **2**: mp 212-213 °C;  $^1\text{H}$  NMR 4.88 (br, 1, NH), 4.30 (br, 1, OH), 3.62 (d, 1,  $J = 11.2$ , H-6), 3.36-3.20 (m, 2, H-2), 2.45 (br ddd, 1,  $J = 16.0, 6.0, 6.0$ , H-4), 2.29 (br dd, 1,  $J = 13.6, 4.8$ , H-8), 2.24 (dd, 1,  $J = 13.6, 13.6$ , H-8), 2.20 (br dd, 1,  $J = 16.0, 4.8$ , H-4), 2.04-1.92 (m, 1, H-7), 1.90-1.70 (m, 2, H-3), 1.19 (d, 3,  $J = 6.8$ );  $^{13}\text{C}$  NMR 193.4, 158.9, 101.5, 76.1, 41.5, 36.8, 35.9, 20.6, 18.9, 18.4; IR 3423, 3264, 3108, 1602 (wk), 1570 (wk), 1513, 1264, 978, 749; HRMS (ESI) calcd for  $\text{C}_{10}\text{H}_{16}\text{NO}_2$  ( $\text{MH}^+$ ) 182.1181, found 182.1186.

Data for **8**:  $^1\text{H}$  NMR 4.87 (br, 1, NH), 4.20 (br, 1, OH), 4.18 (d, 1,  $J = 5.2$ , H-6), 3.37-3.21 (m, 2, H-2), 2.78 (br dd, 1,  $J = 16.0, 5.2$ , H-8), 2.65-2.53 (m, 1, H-7), 2.50 (br ddd, 1,  $J = 16.0, 5.6, 5.6$ , H-4), 2.25 (br ddd, 1, 16.0, 6.8, 6.8, H-4), 2.09 (br d, 1,  $J = 16.0$ , H-8), 1.90-1.70 (m, 2, H-3), 0.92 (d, 3,  $J = 6.8$ );  $^{13}\text{C}$  NMR 192.1, 157.0, 101.1, 73.1, 41.5, 34.9, 33.3, 20.6, 18.6, 11.6; IR 3262 (br), 3089, 1569 (wk), 1514, 1343, 1260, 913, 748; HRMS (ESI) calcd for  $\text{C}_{10}\text{H}_{16}\text{NO}_2$  ( $\text{MH}^+$ ) 182.1181, found 182.1179.

**2,3,4,6,7,8-Hexahydro-7,7-dimethyl-5(1H)-quinolinone (11)**. A solution of 5,5-dimethylcyclohexane-1,3-dione (**10**) (630 mg, 4.5 mmol), 3-bromopropylammonium bromide (**9**) (1.02 g, 4.7 mmol, 1.04 equiv), and 2,6-lutidine (1.6 mL, 13.5 mmol, 3.0 equiv) in 4 mL of EtOH was added to a resealable tube, which was sealed and heated at 140 °C for 2.5 h. The reaction mixture was cooled to room temperature and 1 M NaOH solution (30 mL) was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (30 mL  $\times$  3). The combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to about 5 mL. MeCN (30 mL  $\times$  2) was added and the resulting solution was concentrated again. Flash chromatography of the residue on silica gel (100:1:1 EtOAc/MeOH/ $\text{NEt}_3$ ) gave 491 mg (61%) of **11** as a white solid: mp 180-181 °C;  $^1\text{H}$  NMR 4.41 (br s, 1, NH), 3.26 (br t, 2,  $J = 6.4$ , H-2), 2.35 (br t, 2,  $J = 6.4$ , H-4), 2.20 (br s, 2, H-6 or 8), 2.12 (br s, 2, H-6 or 8), 1.80 (tt,  $J = 6.4, 6.4$ , H-3), 1.04 (s, 6);  $^{13}\text{C}$  NMR 193.7, 157.3, 103.4, 50.2, 43.1, 41.6, 32.4, 28.4 (2 C), 21.1, 18.7; IR 3265 (br), 1721, 1668, 1622, 1518 (wk), 1275, 1135, 913, 748. The  $^1\text{H}$  NMR spectral data are identical to those previously reported.<sup>6</sup>

**7,8-Dihydro-7,7-dimethyl-5(6H)-quinolinone (12)**. A solution of **11** (50 mg, 0.28 mmol) in 4 mL of benzene and dried  $\text{Mn}(\text{OAc})_3$  (292 mg, 1.26 mmol, 4.5 equiv) were added to a resealable tube. The resulting mixture was sealed and heated at 90 °C for 20 min. The reaction mixture was cooled to room temperature and 10% aq.  $\text{NaHSO}_3$  solution (15 mL) was added. The mixture was extracted with EtOAc (15 mL  $\times$  3). The combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash chromatography of the residue on silica gel (100:1:1 EtOAc/MeOH/ $\text{NEt}_3$ ) gave 32 mg (65%) of **12** as a pale yellow oil:  $^1\text{H}$  NMR 8.71 (dd, 1,  $J = 4.8, 1.8$ ), 8.27 (dd, 1,  $J = 8.0, 1.8$ ), 7.30 (dd, 1,  $J = 8.0, 4.8$ ), 3.05 (s, 2), 2.58 (s, 2), 1.12 (s, 6);  $^{13}\text{C}$  NMR 198.1, 162.2, 153.8, 134.5, 127.1, 122.1, 51.9, 46.3, 32.9,

28.2 (2 C); IR, 3070, 1686, 1582, 1300, 1282, 916, 821, 796, 742, 718. The  $^1\text{H}$  NMR,  $^{13}\text{C}$  NMR, and IR spectral data are identical to those previously reported.<sup>7</sup>

**2,3,4,6,7,8-Hexahydro-7,7-dimethyl-1-trifluoroacetyl-5(1*H*)-quinolinone (13).** Trifluoroacetic anhydride (0.56 mL, 4.0 mmol, 2.0 equiv) and  $\text{NEt}_3$  (0.67 mL, 5.0 mmol, 2.5 equiv) were added to a solution of **11** (360 mg, 2.0 mmol) in 6 mL of  $\text{CH}_2\text{Cl}_2$  at 0 °C. The resulting mixture was stirred at 0 °C for 1 h. The reaction was quenched by addition of water (12 mL). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (12 mL  $\times$  3). The combined organic layers were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash chromatography of the residue on silica gel (6:1 hexanes/EtOAc) gave 523 mg (95%) of **13** as a colorless oil:  $^1\text{H}$  NMR 3.66 (dd, 2,  $J = 5.6, 5.6$ , H-2), 2.70 (dd, 2,  $J = 2.0, 2.0$ , H-8), 2.38 (dddd, 2,  $J = 6.8, 6.8, 2.0, 2.0$ , H-4), 2.32 (s, 2, H-6), 1.99-1.91 (m, 2, H-3), 1.06 (s, 6);  $^{13}\text{C}$  NMR 198.5, 155.9 (q, 1,  $J = 36.8$ ,  $\text{CF}_3\text{CO}$ ), 151.7, 124.2, 115.9 (q, 1,  $J = 287$ ,  $\text{CF}_3$ ), 50.8, 45.2 (q, 1,  $J = 3.8$ ,  $\text{CF}_3\text{CONCH}_2$ ), 42.8, 33.7, 27.9 (2 C), 22.4, 20.0; IR 1702, 1664, 1627 (wk), 1199, 1143, 993, 913, 751.

**6-Acetoxy-2,3,4,6,7,8-hexahydro-7,7-dimethyl-1-trifluoroacetyl-5(1*H*)-quinolinone (14).** A solution of **13** (523 mg, 1.9 mmol) in 20 mL of benzene and dried  $\text{Mn}(\text{OAc})_3$  (1.98 g, 8.6 mmol, 4.5 equiv) were added to a resealable tube. The reaction mixture was sealed and heated at 120 °C for 96 h. The reaction mixture was cooled to room temperature and 10% aq.  $\text{NaHSO}_3$  solution (50 mL) was added. The mixture was extracted with EtOAc (50 mL  $\times$  3). The combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash chromatography of the residue on silica gel (6:1 hexanes/EtOAc) gave 26 mg (5%) of recovered **13**, followed by 412 mg (65%) of **14** as a white solid: mp 104-105 °C;  $^1\text{H}$  NMR 5.27 (s, 1, H-6), 3.89 (ddd, 1,  $J = 12.8, 4.0, 4.0$ , H-2), 3.45 (ddd, 1,  $J = 12.8, 10.4, 2.4$ , H-2), 3.21 (ddd, 1,  $J = 18.0, 3.4, 3.4$ , H-8), 2.61 (dd, 1,  $J = 18.0, 1.8$ , H-8), 2.46 (dddd, 1,  $J = 18.4, 8.0, 8.0, 2.4, 2.4$ , H-4), 2.32 (dddd, 1,  $J = 18.4, 6.8, 3.6, 3.6$ , H-4), 2.22 (s, 3), 2.09-1.99 (m, 1, H-3), 1.93-1.81 (m, 1, H-3), 1.13 (s, 3), 1.00 (s, 3);  $^{13}\text{C}$  NMR 192.6, 170.5, 156.0 (q, 1,  $J = 37.2$ ,  $\text{CF}_3\text{CO}$ ), 150.7, 123.8, 115.8 (q, 1,  $J = 287$ ,  $\text{CF}_3$ ), 80.1, 45.3 (q, 1,  $J = 3.8$ ,  $\text{CF}_3\text{CONCH}_2$ ), 42.7, 38.2, 27.2, 22.0, 20.6, 20.2, 19.4; IR 1748, 1705, 1681, 1628 (wk), 1204, 1152, 913, 749; HRMS (ESI) calcd for  $\text{C}_{15}\text{H}_{19}\text{F}_3\text{NO}_4$  ( $\text{MH}^+$ ) 334.1266, found 334.1272.

**2,3,4,6,7,8-Hexahydro-7-methyl-7-(4-methyl-3-penten-1-yl)-5(1*H*)-quinolinone (16).** A solution of 5-methyl-5-(4-methyl-3-penten-1-yl)cyclohexane-1,3-dione (**15**)<sup>10</sup> (400 mg, 1.9 mmol), 3-bromopropylammonium bromide (**9**) (440 mg, 2.0 mmol, 1.05 equiv), and 2,6-lutidine (0.66 mL, 5.7 mmol, 3.0 equiv) in 1.4 mL of EtOH was added to a resealable tube, which was sealed and heated at 140 °C for 8 h. The reaction mixture was cooled to room temperature and 1 M NaOH solution (15 mL) was added. The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (15 mL  $\times$  3). The combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated to about 5 mL. MeCN (15 mL  $\times$  2) was added and the resulting solution was concentrated again. Flash chromatography of the residue on silica gel (25:75:1 hexanes/EtOAc/ $\text{NEt}_3$ ) gave 247 mg (52%) of **16** as a pale yellow oil:  $^1\text{H}$  NMR 5.07 (br t, 1,  $J = 6.8$ ), 4.72

(br, 1, NH), 3.27 (br t, 2,  $J = 6.0$ , H-2), 2.37-2.33 (m, 2, H-4), 2.26 (br d, 1,  $J = 16.0$ , H-6 or 8), 2.26 (br d, 1,  $J = 16.0$ , H-6 or 8), 2.18 (br d, 1,  $J = 16.0$ , H-6 or 8), 2.03 (br d, 1,  $J = 16.0$ , H-6 or 8), 1.98-1.91 (m, 2), 1.80 (tt, 2,  $J = 6.0$ , 6.0, H-3), 1.67 (br s, 3), 1.59 (br s, 3), 1.37-1.33 (m, 2), 1.03 (s, 3);  $^{13}\text{C}$  NMR 193.5, 157.2, 131.6, 124.3, 103.5, 48.5, 41.7, 41.6, 41.5, 35.0, 25.6, 24.8, 22.4, 21.1, 18.7, 17.5; IR 3261 (br), 3072, 1722, 1655, 1571 (wk), 1519, 913, 754; HRMS (ESI) calcd for  $\text{C}_{16}\text{H}_{26}\text{NO}$  ( $\text{MH}^+$ ) 248.2014, found 248.2019.

**2,3,4,6,7,8-Hexahydro-7-methyl-7-(4-methyl-3-penten-1-yl)-1-trifluoroacetyl-5(1*H*)-quinolinone (17).**

Trifluoroacetic anhydride (0.17 mL, 1.2 mmol, 2.0 equiv) and  $\text{NEt}_3$  (0.20 mL, 1.5 mmol, 2.5 equiv) were added to a solution of **16** (147 mg, 0.60 mmol) in 3 mL of  $\text{CH}_2\text{Cl}_2$  at 0 °C. The resulting mixture was stirred at 0 °C for 1 h. The reaction was quenched by addition of water (8 mL). The mixture was extracted with  $\text{CH}_2\text{Cl}_2$  (8 mL  $\times$  3). The combined organic layers were washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash chromatography of the residue on silica gel (6:1 hexanes/EtOAc) gave 190 mg (93%) of **17** as a colorless oil:  $^1\text{H}$  NMR 5.05 (br t, 1,  $J = 6.8$ ), 3.73-3.61 (m, 2, H-2), 2.81 (br d, 1,  $J = 17.6$ , H-8), 2.62 (br d, 1,  $J = 17.6$ , H-8), 2.41-2.34 (m, 2, H-4), 2.38 (d, 1,  $J = 16.4$ , H-6), 2.31 (d, 1,  $J = 16.4$ , H-6), 2.02-1.86 (m, 4, H-3 and  $\text{CH}_2\text{CH}=\text{C}$ ), 1.67 (br s, 3), 1.58 (br s, 3), 1.40-1.36 (m, 2), 1.04 (s, 3);  $^{13}\text{C}$  NMR 198.4, 155.9 (q, 1,  $J = 36.4$ ,  $\text{CF}_3\text{CO}$ ), 151.5, 131.2, 124.5, 123.9, 115.9 (q, 1,  $J = 286$ ,  $\text{CF}_3$ ), 49.3, 45.2 (q, 1,  $J = 3.8$ ,  $\text{CF}_3\text{CONCH}_2$ ), 41.1, 40.8, 36.4, 25.6, 24.7, 22.4, 22.3, 20.0, 17.5; IR 3010, 1738, 1704, 1666, 1628 (wk), 1204, 1143, 914, 752.

**(±)-(5*aR*,6*S*,8*aR*)-2,3,4,5*a*,7,8,8*a*,9-Octahydro-8*a*-methyl-6-(1-methylethenyl)-1-trifluoroacetyl-6*H*-cyclopenta[*g*]quinolin-5(1*H*)-one (20) and (±)-(5*aR*,6*S*,8*aR*)-2,3,4,5*a*,7,8,8*a*,9-Octahydro-8*a*-methyl-6-(1-acetoxy-1-methylethyl)-1-trifluoroacetyl-6*H*-cyclopenta[*g*]quinolin-5(1*H*)-one (21).** A solution of **17** (103 mg, 0.30 mmol) in 4 mL of benzene and dried  $\text{Mn}(\text{OAc})_3$  (313 mg, 1.35 mmol, 4.5 equiv) were added to a resealable tube. The reaction mixture was sealed and heated at 130 °C for 132 h. The reaction mixture was cooled to room temperature and 10% aq.  $\text{NaHSO}_3$  solution (10 mL) was added. The mixture was extracted with EtOAc (10 mL  $\times$  3). The combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated. Flash chromatography of the residue on silica gel (9:1 hexanes/EtOAc) gave 62 mg (61%) of **20** as a colorless oil, followed by 8 mg (8%) of recovered **17**, and 6 mg (5%) of **21** as a white solid.

Data for **20**:  $^1\text{H}$  NMR 4.79-4.73 (m, 2), 3.92 (br ddd, 1,  $J = 12.8$ , 4.0, 4.0, H-2), 3.39 (ddd, 1,  $J = 12.8$ , 10.4, 2.6, H-2), 3.16 (ddd, 1,  $J = 17.6$ , 3.0, 3.0, H-9), 2.91 (ddd, 1,  $J = 11.2$ , 10.4, 7.2, H-6), 2.40-2.32 (m, 2, H-4), 2.35 (d, 1,  $J = 11.2$ , H-5*a*), 2.28 (d, 1,  $J = 17.6$ , H-9), 2.15-1.98 (m, 2, H-3, H-7), 1.91-1.79 (m, 1, H-3), 1.77 (br s, 3), 1.74-1.64 (m, 2, H-7, H-8), 1.64-1.52 (m, 1, H-8), 1.14 (s, 3);  $^{13}\text{C}$  NMR 199.3, 156.0 (q, 1,  $J = 36.4$ ,  $\text{CF}_3\text{CO}$ ), 150.4, 145.3, 123.0, 115.9 (q, 1,  $J = 287$ ,  $\text{CF}_3$ ), 111.9, 59.0, 50.6, 45.5, 45.2 (q, 1,  $J = 3.8$ ,  $\text{CF}_3\text{CONCH}_2$ ), 39.1, 37.1, 29.3, 25.4, 22.5, 20.1, 18.9; IR 3070, 3007, 1737 (wk), 1703, 1660,

1628 (wk), 1203, 1145, 913, 750; HRMS (ESI) calcd for  $C_{18}H_{23}F_3NO_2$  ( $MH^+$ ) 342.1681, found 342.1674. A 2D NOESY experiment showed strong NOEs between H-5a at  $\delta$  2.35 and both the alkenyl hydrogens at  $\delta$  4.79-4.73 and the upfield methyl group at  $\delta$  1.14 and a weak NOE between the alkenyl hydrogens and the upfield methyl group.

Data for **21**: mp 91-92 °C;  $^1H$  NMR 3.80 (br ddd, 1,  $J = 13.2, 5.2, 3.0$ , H-2), 3.50 (ddd, 1,  $J = 13.2, 9.2, 2.8$ , H-2), 3.07 (ddd, 1,  $J = 18.0, 3.0, 3.0$ , H-9), 2.88 (ddd, 1,  $J = 10.4, 10.4, 7.2$ , H-6), 2.57-2.45 (m, 1, H-4), 2.41 (br d, 1,  $J = 18.0$ , H-9), 2.33 (d, 1,  $J = 10.4$ , H-5a), 2.25-2.15 (m, 1, H-4), 2.04-1.86 (m, 3, 2 H-3, H-7), 1.92 (s, 3), 1.72-1.56 (m, 2, H-7, H-8), 1.49 (s, 3), 1.47 (s, 3), 1.5-1.4 (m, 1, H-8), 1.13 (s, 3);  $^{13}C$  NMR 201.3, 170.5, 156.0 (q, 1,  $J = 37.2$ ,  $CF_3CO$ ), 149.6, 123.1, 115.9 (q, 1,  $J = 287$ ,  $CF_3$ ), 83.8, 57.4, 51.8, 45.9, 45.2 (q, 1,  $J = 3.4$ ,  $CF_3CONCH_2$ ), 38.7, 36.6, 26.0, 25.3, 24.9, 23.0, 22.5, 22.4, 20.5; IR 1731, 1708, 1666, 1632 (wk), 1205, 1160, 918, 749; HRMS (ESI) calcd for  $C_{20}H_{27}F_3NO_4$  ( $MH^+$ ) 402.1892, found 402.1886.

## REFERENCES

1. M. G. Vallejo, M. G. Ortega, J. L. Cabrera, V. P. Carlini, S. Rubiales de Barioglio, R. S. Almiron, O. A. Ramírez, and A. M. Agnese, *J. Nat. Prod.*, 2009, **72**, 156.
2. (a) H.-Y. Lin and B. B. Snider, *Org. Lett.*, 2011, **13**, 1234; (b) H.-Y. Lin, R. Causey, G. E. Garcia, and B. B. Snider, *J. Org. Chem.*, 2012, **77**, 7143.
3. (a) A. S. Demir, T. Sayrac, and D. S. Watt, *Synthesis*, 1990, 1119; (b) C. Tanyeli, D. Özermirhan, and B. Sezen, *Tetrahedron*, 2002, **58**, 9983; (c) A. S. Demir, Z. Caliskan, and E. Sahin, *J. Mol. Catal. B*, 2007, **44**, 87; (d) A. S. Demir and M. Emrullahoglu, *Curr. Org. Syn.*, 2007, **4**, 223.
4. For related *N*-acylations of vinylogous amides see: (a) Y. Tamura, Y. Kita, and J. Uraoka, *Chem. Pharm. Bull.*, 1972, **20**, 876; (b) H. M. Sklenicka, R. P. Hsung, M. J. McLaughlin, L.-I. Wei, A. I. Gerasyuto, and W. B. Brennessel, *J. Am. Chem. Soc.*, 2002, **124**, 10435; (c) G. Li, L. J. Carlson, I. K. Sagamanova, B. W. Slafer, R. P. Hsung, C. Gilardi, H. M. Sklenicka, and N. Sydorenko, *Synthesis*, 2009, 2905.
5. For spectral data of related 6-hydroxy-5-methyl-2-cyclohexenones, see: (a) G. M. Rubottom and J. M. Gruber, *J. Org. Chem.*, 1978, **43**, 1599; (b) A. K. Musser and P. L. Fuchs, *J. Org. Chem.*, 1982, **47**, 3121.
6. (a) H. Reinshagen, *Angew. Chem., Int. Ed. Engl.*, 1964, **3**, 807; (b) J. C. Rohloff, N. H. Dyson, J. O. Gardner, T. V. Alfredson, M. L. Sparacino, and R. Robinson III, *J. Org. Chem.*, 1993, **58**, 1935.
7. F. Liéby-Muller, C. Allais, T. Constantieux, and J. Rodriguez, *Chem. Commun.*, 2008, 4207.
8. A. S. Demir, Ö. Reis, and A. C. Igdir, *Tetrahedron*, 2004, **60**, 3427.
9. (a) B. McCarthy-Cole, L. Han, and B. B. Snider, *J. Org. Chem.*, 1996, **61**, 7832; (b) B. B. Snider and

- E. Y. Kiselgof, *Tetrahedron*, 1996, **52**, 6073.
10. G. Mehta and M. K. Bera, *Tetrahedron Lett.*, 2008, **49**, 1417.
11. (a) B. Chenera, C.-P. Chuang, D. J. Hart, and C.-S. Lai, *J. Org. Chem.*, 1992, **57**, 2018. See also: (b) A.-C. Guevel and D. J. Hart, *J. Org. Chem.*, 1996, **61**, 465; (c) A.-C. Guevel and D. J. Hart, *J. Org. Chem.*, 1996, **61**, 473; (d) Y.-L. Kuo, M. Dhanasekaran, and C.-K. Sha, *J. Org. Chem.*, 2009, **74**, 2033.