

HETEROCYCLES, Vol. 87, No. 10, 2013, pp. 2047 - 2052. © The Japan Institute of Heterocyclic Chemistry
Received, 11th July, 2013, Accepted, 14th August, 2013, Published online, 30th August, 2013
DOI: 10.3987/COM-13-12773

ALKALOIDS FROM *MELODINUS SUAVEOLENS*

Ting-Ting Zhang,^{a†} Zhi-Wen Liu,^{a†} Wen-Jing Wang,^a Yong-Bin Tong,^a
Fang-Fang Xu,^{b*} Jing-Quan Yuan,^c Bo Liu,^b Xiao-Qi Zhang,^{a*} and Wen-Cai
Ye^a

^a Institute of Traditional Chinese Medicine and Natural Products, College of Pharmacy, Jinan University, Guangzhou 510632, P. R. China

^b Guangdong Provincial Academy of Chinese Medical Sciences, Guangzhou 510120, P. R. China


^c Guangxi Medicinal Herb Garden, Nanning 530023, P. R. China

[†] These authors contributed equally to this work.

* Corresponding authors. Tel: +86-20-85221559; e-mail: xqzhang7401@aliyun.com (X. Q. Zhang), xudubfang@163.com (F. F. Xu)

Abstract – Two new quinoline alkaloids, 14,15- β -epoxyscandine (**3**) and meloscine *N*-oxide (**6**), along with six known alkaloids scandine (**1**), scandine *N*-oxide (**2**), meloscandonine (**4**), meloscine (**5**), tubotaiwine (**7**), polyneuridine (**8**) were isolated from the twigs and leaves of *Melodinus suaveolens*. The structures of the new compounds were elucidated on the basis of spectroscopic methods and circular dichroism experiments.

The genus *Melodinus* (Apocynaceae) comprises 53 species mainly distributed in tropical or subtropical Asia and Australia.¹ This genus has been regarded as a rich source of monoterpenoid indole alkaloids, which originate from the condensation of tryptophan with secologanin.² Up to now, more than 100 monomeric and dimeric indole alkaloids as well as quinoline alkaloids have been isolated from *Melodinus* sp. The interesting chemical significance of the *Melodinus* plants prompted us to initiate a phytochemical study on the twigs and leaves of *Melodinus suaveolens*, which led to the isolation of two new quinoline alkaloids, 14,15- β -epoxyscandine (**3**) and meloscine *N*-oxide (**6**), along with six known alkaloids scandine (**1**),³ scandine *N*-oxide (**2**),³ meloscandonine (**4**),⁴ meloscine (**5**),⁵ tubotaiwine (**7**),⁶ and polyneuridine (**8**).⁷



14,15- β -Epoxy-scandine (**3**) was obtained as yellow oil. The molecular formula of **3** was established to be $C_{21}H_{22}N_2O_4$ by its HR-ESI-MS (m/z 367.1648 $[M+H]^+$, calcd for $C_{21}H_{23}N_2O_4$, 367.1652). The IR spectrum of **3** suggested the presence of NH (3456 cm^{-1}), carbonyl ($1744, 1663\text{ cm}^{-1}$) and aromatic ring ($1596, 1498\text{ cm}^{-1}$). The UV spectrum showed absorption maxima at 341, 326, 287, 257 and 214 nm, suggesting that **3** was a quinoline alkaloid. The ^1H NMR spectrum of **3** displayed signals for an *ortho*-disubstituted phenyl ring [δ_{H} 7.30 (1H, dd, $J = 7.9, 1.3\text{ Hz}$, H-9), 7.12 (1H, td, $J = 7.9, 1.3\text{ Hz}$, H-11), 7.03 (1H, td, $J = 7.9, 1.3\text{ Hz}$, H-10), and 6.74 (1H, dd, $J = 7.9, 1.3\text{ Hz}$, H-12)], a terminal double bond [δ_{H} 5.63 (1H, dd, $J = 17.8, 11.1\text{ Hz}$, H-19), 4.93 (1H, d, $J = 17.8\text{ Hz}$, H-18a) and 4.92 (1H, d, $J = 11.1\text{ Hz}$, H-18b)], and a methoxyl group [δ_{H} 3.63 (3H, s, CO_2Me)]. The ^{13}C NMR spectrum showed twenty-one carbon signals, including two carbonyls, eight olefinic carbons, three quaternary carbons, one methoxyl, three methines and four methylenes. With the aid of ^1H - ^1H COSY, HSQC and HMBC experiments, the ^1H and ^{13}C NMR signals of **3** were assigned as shown in Table 1.

Comparison of the NMR data of **3** with those of the known compound scandine (**1**)³ revealed their structural similarity, except for an epoxy group [δ_{C} 58.2, 55.5; δ_{H} 3.48 (1H, m), 3.03 (1H, d, $J = 3.7\text{ Hz}$)] instead of a double bond [δ_{C} 132.7, 124.2; δ_{H} 5.72 (1H, m, H-14), 5.65 (1H, overlapped, H-15)]. The ^1H - ^1H COSY spectrum of **3** revealed the presence of the spin system (C-3 to C-15) and the HMBC cross peaks between H-3 α /H-3 β /H-15/H-19/H-17 α /H-17 β and C-21 indicated that epoxy group was located at C-14 and C-15 (Figure 1). Based on the above evidence, the planar structure of **3** was elucidated as 14,15-epoxy-scandine.⁴ However, the relative configuration of the epoxy group was not clearly defined in the previous work. In this paper, the relative stereochemistry of **3** was deduced by a ROESY experiment,

in which correlations between H-19 and H-15/H-21 suggested that these protons had the same orientation. Furthermore, the CD spectrum of **3** showed the same Cotton effects as scandine (**1**) (Figure 2), indicating the presence of *S, R, R, R* configurations at C-7, C-16, C-20 and C-21, respectively. Thus, the structure of **3** was elucidated as 14,15- β -epoxyscandine.

Table 1. NMR Data of Compounds **3** and **6** (CD₃OD, δ in ppm, *J* in Hz)^a

No	3 ^b		6 ^c	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}
2		170.8		171.3
3	β	3.43 dd (12.8, 5.7)	β	4.05 dd (14.9, 6.6)
	α	2.59 d (12.8)	α	3.89 m
5	β	3.09 dd (8.0, 8.0)	α	3.99
	α	2.69 m	β	3.32
6	α	2.18 dd (13.6, 5.7)	α	2.51 m
	β	1.93 m	β	2.20
7		58.2		59.0
8		129.9		125.9
9		129.1		129.7
10		124.8		125.6
11		128.6		129.8
12		116.5		117.0
13		135.8		136.6
14		55.5		125.7
15		58.2		136.4
16		66.1		49.2
17	β	3.13 d (12.9)	β	2.23
	α	2.64 dd (12.9, 1.2)	α	1.93 dd (13.7, 12.8)
18	a	4.93 d (17.8)	a	5.16 d (17.2)
	b	4.92 d (11.1)	b	5.04 d (10.4)
19		140.0		141.2
20		49.6		52.7
21		83.9		98.3
CO ₂ Me		172.3		
CO ₂ Me		53.0		

^a Overlapped signals were reported without designating multiplicity.

^b Measured at 600 MHz. ^c Measured at 400 MHz.

Meloscine *N*-oxide (**6**) was isolated as yellow oil with a molecular formula of C₁₉H₂₀N₂O₂, as determined by the HR-ESI-MS at *m/z* 309.1597 [M+H]⁺ (calcd for C₁₉H₂₁N₂O₂, 309.1598). The IR and UV spectra of

6 were similar to those of **3**, suggesting that **6** was also a quinoline alkaloid. The ^1H NMR spectrum of **6** showed signals for an *ortho*-disubstituted phenyl ring [δ_{H} 7.90 (1H, dd, $J = 7.7, 1.4$ Hz, H-9), 7.24 (1H, td, $J = 7.7, 1.4$ Hz, H-11), 7.17 (1H, td, $J = 7.7, 1.3$ Hz, H-10), 6.91 (1H, dd, $J = 7.7, 1.3$ Hz, H-12)], a double bond [δ_{H} 6.21 (1H, m, H-14) and 6.08 (1H, dd, $J = 9.8, 2.8$ Hz, H-15)], and a terminal double bond [δ_{H} 5.79 (1H, dd, $J = 17.2, 10.4$ Hz, H-19), 5.16 (1H, d, $J = 17.2$ Hz, H-18a) and 5.04 (1H, d, $J = 10.4$ Hz, H-18b)]. The ^{13}C NMR spectrum displayed nineteen carbon signals, including one carbonyl, ten olefinic carbons, two quaternary carbons, two methines and four methylenes. Detailed examination of 1D and 2D NMR spectra of **6** and comparison with those of the known compound meloscine (**5**)⁵ revealed that **6** was the N-oxide of compound **5**, in particular the characteristic downfield shifts of the C-21 [δ_{C} 98.3; δ_{H} 4.48 (1H, s)] in **6**. In the HMBC spectrum, the correlations between H-3 α /H-3 β /H-5 α /H-5 β /H-15/H-19/H-17 α /H-17 β and C-21 further confirmed the oxidation of N₄ (Figure 1). In the ROESY spectrum, correlations between H-9 and H-6 α /H-21 suggested that H-6 α and H-21 were α -oriented, whereas the correlations H-16 and H-6 β /H-17 β suggested that H-16 was β -oriented. Furthermore, H-19, H-17 α and H-21 were assigned on the same side on the basis of the ROESY correlations among them. The absolute configuration of **6** was identical to that of meloscine (**5**), since they showed similar Cotton effects in the CD measurement (Figure 2). Accordingly, the structure of **6** was determined as meloscine N-oxide.




Figure 1. Key ^1H - ^1H COSY and HMBC correlations of **3** and **6**




Figure 2. CD spectra of compounds **1**, **3**, **5** and **6**

The structures of the known compounds were identified by the comparison of their spectroscopic data (UV, IR, ^1H NMR, ^{13}C NMR, MS) with the data from the corresponding values in the literature as scandine (**1**),³ scandine N-oxide (**2**),³ meloscandonine (**4**),⁴ meloscine (**5**),⁵ tubotaiwine (**7**),⁶ and polyneuridine (**8**),⁷ respectively.

EXPERIMENTAL

General experimental procedures: Optical rotations were carried out using a Jasco P-1020 digital polarimeter. UV spectra were measured on a Jasco V-550 UV/VIS spectrophotometer with a 1 cm length cell. IR spectra were recorded on a Bruker Equinox 55 infrared spectrometer with KBr disc. CD spectra were measured on a Chirascan spectrometer (Applied Photophysics Ltd) at 25 °C for 200-400 nm with a quartz cell of path length 1 cm. HR-ESI-MS data were measured on an Agilent 6210 ESI/TOF mass spectrometer. NMR experiments were performed on Bruker AV-400 and AV-600 spectrometers. Column chromatography (CC) were performed on silica gel (200-300 mesh; Qingdao Marine Chemical Inc., Qingdao, China) and ODS (YMC, Kyoto, Japan), respectively. Preparative high-performance liquid chromatography (HPLC) was carried on a Agilent 1260 system equipped with a G1310B Iso pump, a G1365D MWD VL detector and a CAPCELL PAK MGII C₁₈ reversed-phase column (20×250 mm, 5 μm, Shiseido Fine Chemicals Ltd, Japan).

Plant material: The leaves and twigs of *M. suaveolens* were collected in Jinxiu, a Yao Autonomous County of Guangxi Province of China, in September 2010, and authenticated by Dr. Jing-Quan Yuan (Guangxi Medicinal Herb Garden). A voucher specimen (No. CP2010093001) is deposited in the herbarium of the College of Pharmacy, Jinan University, Guangzhou, China.

Extraction and isolation: The air-dry leaves and twigs (18 kg) of *M. suaveolens* was powdered and extracted with 95% EtOH (50 L) three times at room temperature, and the solution was concentrated under reduced pressure to afford a brownish residue (1700 g), which was suspended in water, and then successively partitioned with petroleum ether, EtOAc and *n*-BuOH, respectively. The EtOAc extract (326 g) was subjected to a silica gel column with gradient CHCl₃-MeOH (100:0→0:100) to afford 9 fractions (Fr-E1~Fr-E9). Fraction E2 was then subjected to sephadex LH-20 (CHCl₃-MeOH, 1:1) and ODS (MeOH-H₂O, 30:90→90:10) column chromatography (CC) to yield **1** (17.5 mg) and **8** (2.3 mg). The *n*-BuOH extract (306 g) was subjected to a silica gel column with gradient CHCl₃-MeOH (100:0→0:100) to afford 11 fractions (Fr-B1~Fr-B11). Fraction B3 was further separated on silica gel column eluting with CHCl₃-MeOH (100:1→90:10), followed by HPLC (MeOH-H₂O+0.1% Et₂NH, 65:35) to afford **3** (2.4 mg), **4** (3.2 mg) and **5** (13.5 mg). The Fr-B5 was purified by Sephadex LH-20 column (CHCl₃-MeOH, 1:1) and HPLC (MeOH-H₂O+0.1% Et₂NH, 60:40) to afford **2** (2.4 mg), **6** (2.5 mg) and **7** (2.7 mg).

14,15-β-Epoxy scandine (3). Yellow oil; $[\alpha]_D^{25} +109.8^\circ$ (*c* 0.9, MeOH); UV $\lambda_{\max}^{\text{MeOH}}$ (log ϵ): 341 (3.38), 326 (3.43), 287 (3.74), 257 (4.12), 214 (4.56) nm; CD (MeOH, $\Delta\epsilon$) λ_{\max} 257 (+8.5), 221 (-21.4) nm; IR ν_{\max}^{KBr} : 3456, 2924, 1744, 1663, 1634, 1596, 1498, 1384, 1235, 1115, 756 cm⁻¹; HR-ESI-MS *m/z* 367.1648 [M+H]⁺ (calcd for C₂₁H₂₃N₂O₄, 367.1652).

Meloscine N-oxide (6). Yellow oil; $[\alpha]_D^{25} +128.6^\circ$ (MeOH, c 0.6); UV $\lambda_{\max}^{\text{MeOH}}$ ($\log \epsilon$): 252 (4.40), 210 (3.91) nm; CD (MeOH, $\Delta\epsilon$) λ_{\max} 254 (+6.2), 224 (-5.3) nm; IR ν_{\max}^{KBr} : 3455, 2923, 1665, 1594, 1491, 1384, 763 cm^{-1} ; HR-ESI-MS m/z 309.1597 $[\text{M}+\text{H}]^+$ (calcd for $\text{C}_{19}\text{H}_{21}\text{N}_2\text{O}_2$, 309.1598).

ACKNOWLEDGEMENTS

Financial support of this research was provided by grants from National Natural Science Foundation of China (Nos. 81172947, 81202428), Ministry of Science and Technology of China (Nos. 2013DFM30080, 2013BAI11B05, 2012ZX09103201-056), China Postdoctoral Science Foundation (No. 2013M531910), Educational Commission of Guangdong Province (No. gihz1003), and the Program for New Century Excellent Talents in University (No. NCET-12-0676).

REFERENCES

1. C. M. Li, H. L. Zheng, S. G. Wu, and H. D. Sun, *Acta Bot. Yunn.*, 1994, **16**, 315.
2. S. Baassou, H. M. Mehri, A. Rabaron, and M. Plat, *Tetrahedron Lett.*, 1983, **24**, 761; K. Bernauer, G. Englert, W. Vetter, and E. Weiss, *Helv. Chim. Acta*, 1969, **52**, 1886; Y. L. He, W. M. Chen, and X. Z. Feng, *J. Nat. Prod.*, 1994, **57**, 411; M. H. Mehri, A. Rabaron, T. Sevenet, and M. M. Plat, *Phytochemistry*, 1978, **17**, 1451; Y. W. Zhang, R. Yang, Q. Cheng, and K. Ofuji, *Helv. Chim. Acta*, 2003, **86**, 415; Y. P. Liu, Y. Li, X. H. Cai, X. Y. Li, L. M. Kong, G. G. Cheng, and X. D. Luo, *J. Nat. Prod.*, 2012, **75**, 220; M. Kitajima, S. Ohara, N. Kogure, Y. Q. Wu, R. P. Zhang, and H. Takayama, *Heterocycles*, 2012, **85**, 1949; C. R. Hutchinson, *Tetrahedron*, 1981, **37**, 1047.
3. C. J. Zhou, H. Y. Guan, Y. H. Zhang, R. Xu, X. J. Hao, and X. S. Yang, *Zhongchengyao*, 2012, **34**, 85.
4. L. W. Guo and Y. L. Zhou, *Phytochemistry*, 1993, **34**, 563.
5. C. M. Li, H. L. Zheng, S. G. Wu, and H. D. Sun, *Yunnan Zhiwu Yanjiu*, 1994, **16**, 315.
6. C. M. Li, G. D. Tao, S. G. Wu, J. Y. Zhong, and Y. L. Zhou, *Zhiwu Xuebao*, 1989, **31**, 792.
7. R. Jokela and M. Lounasmaa, *Heterocycles*, 1996, **43**, 1015.