

HETEROCYCLES, Vol. 90, No. 1, 2015, pp. 104 - 107. © 2015 The Japan Institute of Heterocyclic Chemistry  
 Received, 6th March, 2014, Accepted, 24th March, 2014, Published online, 24th March, 2014  
 DOI: 10.3987/COM-14-S(K)17

## CRYSTAL STRUCTURES OF 3-METHYLPYRROLO[2,3-*b*]TROPONE AND ITS COPPER(II) COMPLEX<sup>†</sup>


Kanji Kubo,<sup>a\*</sup> Taisuke Matsumoto,<sup>b</sup> Keiko Ideta,<sup>b</sup> and Akira Mori<sup>b</sup>

<sup>a</sup>Department of Life Science and Technology, Faculty of Engineering, Hokkai-Gakuen University, Asahimachi, Toyohira-ku, Sapporo 062-8605, Japan.  
 kubo@lst.hokkai-s-u.ac.jp

<sup>b</sup>Institute for Materials Chemistry and Engineering, Kyushu University, Kasuga-koen, Kasuga, Fukuoka 816-8580, Japan

**Abstract** – The crystal structures of 3-methylpyrrolo[2,3-*b*]tropone (**1**) and its Cu(II) complex (**1**<sub>2</sub>·Cu) were analyzed by X-ray crystallography. The structure of **1** exists in the crystal in its keto form rather than in the enol form (**1'**). Pyrrolo[2,3-*b*]tropone (**1**) formed a 2:1 complex with Cu(II). The Cu(II) ion has a tetragonal environment formed by two tropone O atoms and two pyrrole N atoms.

Pyrroles represent an important class of heterocycles in organic chemistry. They are structural units in many natural products such as porphyrin, which form the complexes with Fe(II) and Cu(II) ions.<sup>1</sup> Troponoids, being a remarkable class of non-benzenoid  $\pi$  conjugated systems, form complexes with metal salts such as CuCl<sub>2</sub>, ZnCl<sub>2</sub>, CdCl<sub>2</sub>, CoCl<sub>2</sub>, MnCl<sub>2</sub>, and HgCl<sub>2</sub>.<sup>2</sup> Pyrrolo[2,3-*b*]tropones (cyclohepta[*b*]pyrrol-8(1*H*)-one), being isoelectronic with 8-quinolinols, are interesting compounds biochemically.<sup>3</sup> They give coloration when treated with metal ion such as Fe(III) and Cu(II). The color fades upon addition of mineral acid.<sup>3,4</sup> This suggests that pyrrolo[2,3-*b*]tropones can be used as an analytical reagent for metal cations. However the crystal structures of the metal complex of pyrrolo[2,3-*b*]tropone have not been elucidated. We now report the structures of 3-methylpyrrolo[2,3-*b*]tropone (**1**) and its Cu(II) complex (**1**<sub>2</sub>·Cu).



<sup>†</sup>Dedicated to Professor Dr. Isao Kuwajima on the occasion of his 77th birthday

Compound (**1**)<sup>5</sup> was prepared by condensation of 2-hydrazinotropone with 1-propanal accompanied by cyclization as reported in a previous paper.<sup>3</sup> Single crystals of **1** were grown in a mixture of chloroform-methanol (1:1 *v/v*) at room temperature. An ORTEP drawing<sup>6</sup> of molecule of **1** is shown in Figure 1a. The structure of **1** contains two crystallographically independent molecules (**1a**) and (**1b**). The pyrrolotropone moiety of **1** is approximately planar; the deviations from the least-squares plane defined by C1-C9/O1/N1 do not exceed 0.1 Å. The C-C bond lengths of seven-membered ring show bond alternation typical of tropone; the C1-O1 bond length (1.248(3) Å) is similar to that observed in the structure of unsubstituted tropone (1.259 Å).<sup>7</sup> The C-C and C-N bond lengths in the pyrrole ring are close to those found in unsubstituted pyrrole.<sup>8</sup> The objective location of the H1 atom bond to N1, rather than O1, as well as the tropone-like bond-length distribution, makes assignment of the keto form (**1**) rather than the enol form (**1'**), unambiguous.




Figure 1. (a) ORTEP structure of **1**, showing 50% probability ellipsoids and (b) intermolecular N-H...O,  $\pi$ ... $\pi$ , and CH... $\pi$  interactions in **1**. Symmetry codes: (i) 1-x, -y, 2-z; (ii) 1-x, 1-y, 2-z, (iii) 3-x, 2-y, 3-z.

Table 1. Selected bond lengths (Å) of **1** and **1**<sub>2</sub>·Cu

Bond Lengths	<b>1a</b>	<b>1b</b>	<b>1</b> <sub>2</sub> ·Cu
C1-O1 (C11-O2)	1.248(3)	1.243(3)	1.285(4)
C2-N1 (C12-N2)	1.347(3)	1.369(3)	1.355(4)
C1-C2 (C11-C12)	1.459(4)	1.427(4)	1.423(4)
C2-C3 (C12-C13)	1.409(4)	1.415(4)	1.420(4)
C3-C4 (C13-C14)	1.402(4)	1.393(4)	1.405(4)
C4-C5 (C14-C15)	1.354(4)	1.377(4)	1.379(4)
C5-C6 (C15-C16)	1.410(4)	1.403(4)	1.410(5)
C6-C7 (C16-C17)	1.364(4)	1.369(4)	1.369(5)
C7-C1 (C17-C11)	1.435(4)	1.432(4)	1.425(4)
C8-C9 (C18-C19)	1.376(4)	1.355(4)	1.428(4)
C9-C3 (C19-C13)	1.442(4)	1.449(4)	1.387(5)
N1-C8 (N2-C18)	1.362(3)	1.367(3)	1.360(4)
C9-C10 (C19-C20)	1.481(4)	1.492(4)	1.489(5)
Cu1-O1	-	-	1.9860(19)
Cu1-N1	-	-	1.912(3)

The pyrrole NH group participates in an intermolecular hydrogen bond of the N-H $\cdots$ O type as shown in Figure 1b and Table 2. An intermolecular  $\pi\cdots\pi$  interaction is observed. The distance

Table 2. Hydroge-bond geometry ( $\text{\AA}$ ,  $^\circ$ ) of **1**

D—H $\cdots$ A	D—H	H $\cdots$ A	D $\cdots$ A	D—H $\cdots$ A
N1—H1 $\cdots$ O2 <sup>i</sup>	0.860	1.985	2.815(3)	162
N2—H10 $\cdots$ O1 <sup>i</sup>	0.860	1.941	2.768(3)	161
C10—H7 $\cdots$ C1 <sup>ii</sup>	0.989	2.839	3.475(4)	123

Symmetry codes: (i) 1-x, -y, 2-z; (ii) 1-x, 1-y, 2-z.

between intermolecular pyrrolotropone planes is 3.424(4)  $\text{\AA}$  for C9 $\cdots$ C13, which is within the range associated with  $\pi\cdots\pi$  interactions.<sup>4,9</sup> There is an intermolecular C-H $\cdots$  $\pi$  interaction with distance for this type of interaction (2.8–3.1  $\text{\AA}$ ).<sup>4,10</sup> The combination of intermolecular N-H $\cdots$ O,  $\pi\cdots\pi$ , and C-H $\cdots$  $\pi$  interactions in **1** builds up a three-dimensional network.

Single crystals of Cu(II) complex of **1** (**1**<sub>2</sub>·Cu) were grown in a mixture of chloroform-methanol (1:1 v/v) containing of **1** (2 eq.) and copper(II) acetate (1eq.) at room temperature. An ORTEP drawing<sup>11</sup> of molecule of **1**<sub>2</sub>·Cu is shown in Figure 2a. The pyrrolotropone (**1**) forms 2:1 complex with Cu(II) ion. The complex has a tetragonal planar Cu(II). The Cu atom is positioned at the center of symmetry, coordinated with two N atoms of the pyrrole ring and two O atoms of the tropone ring. The Cu-O bond distance (1.9860(19)  $\text{\AA}$ ) is shorter than that (2.036  $\text{\AA}$ )<sup>12</sup> of bis(2-acetylpyrrolate)copper(II) and longer than that (1.913, 1.915  $\text{\AA}$ )<sup>13</sup> of bis(tropolonato)copper(II). The Cu-N bond distance is similar to that (1.912  $\text{\AA}$ )<sup>13</sup> of bis(2-acetylpyrrolate)copper(II). The C-O bond length is longer than that of **1** and agree with those (1.27-1.31  $\text{\AA}$ )<sup>13,14</sup> in other metal tropolonates.

Intermolecular  $\pi\cdots\pi$  interactions in the crystal structure of **1**<sub>2</sub>·Cu are observed in Figure 2b. The distance between intermolecular pyrrolotropone planes is 3.454(4)  $\text{\AA}$  for C1 $\cdots$ C8<sup>iv</sup> (symmetry codes: (iv) x, y-1, z), which is within the range associated with  $\pi\cdots\pi$  interactions.<sup>4,9</sup> Intermolecular C-H $\cdots$  $\pi$  and C-H $\cdots$ O interactions (Figure 2b and Table 3) are observed with distances typical for these type of interactions:




Figure 2. (a) ORTEP structure of **1**<sub>2</sub>·Cu, showing 50% probability ellipsoids and (b) intermolecular  $\pi\cdots\pi$ , CH $\cdots$  $\pi$ , and C-H $\cdots$ O interactions in **1**<sub>2</sub>·Cu. Symmetry codes: (i) 3/2-x, y-1/2, 1/2-z; (ii) 1/2-x, y-1/2, 1/2-z; (iii) 1/2-x, 1/2+y, 1/2-z; (iv) x, y-1, z.

C-H... $\pi$ =2.8–3.1 Å and C-H...O=2.5–2.7 Å.<sup>4,10</sup> The combination of intermolecular C-H... $\pi$  and C-H...O interactions and  $\pi$ ... $\pi$  stacking interactions in  $\mathbf{1}_2\cdot\text{Cu}$  build up a three-dimensional network.

Table 3. Hydroge-bond geometry (Å, °) of  $\mathbf{1}_2\cdot\text{Cu}$ 

D—H...A	D—H	H...A	D...A	D—H...A
C4—H4...C4 <sup>i</sup>	0.950	2.847	3.701(4)	160
C6—H6...O1 <sup>ii</sup>	0.950	2.696	3.412(4)	133
C7—H7...C6 <sup>iii</sup>	0.950	2.847	3.641(4)	142
C10—H10B...C8 <sup>iv</sup>	0.980	2.719	3.687(4)	170

Symmetry codes: (i) 3/2-x, y-1/2, 1/2-z; (ii) 1/2-x, y-1/2, 1/2-z; (iii) 1/2-x, 1/2+y, 1/2-z; (iv) x, y-1, z.

## ACKNOWLEDGEMENTS

This work was performed under the Cooperative Research Program of "Network Joint Research Center for Materials and Devices".

## REFERENCES AND NOTES

- B. A. Trofimov, L. N. Sobenina, A. P. Demenev, and A. I. Mikhaleva, *Chem. Rev.*, 2004, **104**, 2481.
- T. Nozoe, *Bull. Chem. Soc. Jpn.*, 1936, **11**, 295; H. Erdtmann and J. Gripenberg, *Nature*, 1948, **161**, 719; T. Asao and Y. Kikuchi, *Chem. Lett.*, 1972, 413.
- T. Nozoe and K. Kikuchi, 'Dai Yuki Kagaku,' Vol. 13, ed. by M. Kotake, Asakura Syoten, Tokyo, 1959, pp. 572-575; G. Sunagawa, N. Soma, H. Nakano, and Y. Matsumoto, *Yakugaku Zasshi*, 1961, **81**, 1799; Y. Miura, N. Okamoto, and A. Shimamune, *Seikagaku*, 1960, **32**, 744.
- K. Kubo, T. Tsujimoto, and A. Mori, *Acta Cryst.*, 2001, **E57**, o225; K. Kubo, T. Tsujimoto, N. Kato, and A. Mori, *Acta Cryst.*, 2001, **E57**, o370.
- <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>)  $\delta$  = 2.36 (3H, s), 6.89 (1H, dd,  $J$  = 10.6, 8.6 Hz), 7.19 (1H, d,  $J$  = 12.0 Hz), 7.24 (1H, s), 7.37 (1H, dd,  $J$  = 12.0, 8.6 Hz), 7.66 (1H, d,  $J$  = 10.6 Hz), and 10.60 (1H, brs). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>)  $\delta$  = 9.8, 119.7, 123.0, 124.3, 128.0, 131.3, 134.0, 136.3, 138.4, and 176.3.
- Crystal data of **1**, C<sub>10</sub>H<sub>9</sub>NO,  $M_r$  = 159.19, Monoclinic,  $P2_1/n$ ,  $a$  = 11.307(6) Å,  $b$  = 9.990(5) Å,  $c$  = 14.040(8) Å,  $\beta$  = 101.64(3)°,  $V$  = 101.64(3) Å<sup>3</sup>,  $Z$  = 8,  $D_x$  = 1.361 Mg m<sup>-3</sup>,  $T$  = 293 K,  $\mu$  = 7.127 cm<sup>-1</sup>,  $wR(F^2)$  = 0.2280,  $R$  = 0.0753.
- M. J. Barrow, O. S. Mills, and G. Filippini, *J. Chem. Soc., Chem. Commun.*, 1973, 66.
- R. Goddard, O. Heinemann, and C. Krüger, *Acta Cryst.*, 1997, **C53**, 1846.
- C. K. Prout, T. M. Orley, I. J. Tickle, and J. D. Wright, *J. Chem. Soc., Perkin Trans. 2*, 1973, 523.
- K. Kubo, T. Matsumoto, K. Katahira, and A. Mori, *Acta Cryst.*, 2005, **E61**, o2750; K. Kubo, T. Matsumoto, K. Ideta, H. Takechi, and H. Takahashi, *Heterocycles*, 2012, **84**, 315.
- Crystal data of  $\mathbf{1}_2\cdot\text{Cu}$ , C<sub>20</sub>H<sub>16</sub>CuN<sub>2</sub>O<sub>2</sub>,  $M_r$  = 379.90, Monoclinic,  $P2_1/n$ ,  $a$  = 9.904(3) Å,  $b$  = 5.2142(10) Å,  $c$  = 15.092(4) Å,  $\beta$  = 96.938(12)°,  $V$  = 773.6(3) Å<sup>3</sup>,  $Z$  = 2,  $D_x$  = 1.631 Mg m<sup>-3</sup>,  $T$  = 123 K,  $\mu$  = 14.282 cm<sup>-1</sup>,  $wR(F^2)$  = 0.1073,  $R$  = 0.0434.
- W. J. Birdsall, B. A. Weber, and M. Parvez, *Inorg. Chim. Acta*, 1992, **196**, 213.
- J. E. Berg and A. M. Pilotti, *Acta Cryst.*, 1978, **B34**, 3071.
- M. C. Barret, M. F. Mahon, K. C. Molloy, P. Wright, and J. E. Creeth, *Polyhedron*, 2002, **21**, 1761.