

HETEROCYCLES, Vol. 92, No. 11, 2016, pp. 2059 - 2065. © 2016 The Japan Institute of Heterocyclic Chemistry  
 Received, 17th August, 2016, Accepted, 20th September, 2016, Published online, 13th October, 2016  
 DOI: 10.3987/COM-16-13556

## TWO NEW ATISINE-TYPE C<sub>20</sub>-DITERPENOID ALKALOIDS FROM *ACONITUM LEUCOSTOMUM*

Wenliang Xu,<sup>a,†</sup> Lin Chen,<sup>a,b,c,†</sup> Lianhai Shan,<sup>a</sup> Feng Gao,<sup>a</sup> Shuai Huang,<sup>a,\*</sup>  
 and Xianli Zhou<sup>a,b,\*</sup>

<sup>a</sup>School of Life Science and Engineering, Southwest Jiaotong University, Chengdu 610031 China; <sup>b</sup>Key Laboratory of Advanced Technologies of Materials, Ministry of Education, School of Material Science and Engineering, Southwest Jiaotong University, Chengdu 610031 China; <sup>c</sup>School of Chemistry and Chemical Engineering, China West Normal University, Nanchong 637002 China. Email: shuaih@swjtu.edu.cn; zhoulx@swjtu.edu.cn. <sup>†</sup>These authors contributed equally to this work and should be considered co-first authors.

**Abstract** – Two new atisine-type C<sub>20</sub>-diterpenoid alkaloids, leucostomine A (**1**) and leucostomine B (**2**) were isolated from the root of *Aconitum leucostomum*. Their structures were determined by spectroscopic methods including 1D-, 2D-NMR, and HR-ESI-MS.


*Aconitum leucostomum* Worosch. (Ranunculaceae) is a perennial herb distributed in the Gansu Province and Xinjiang Uygur Autonomous Region of China. It has been used as a folk medicine for the treatment of rheumatoid arthritis, cancer, and various types of pains for a long time.<sup>1</sup> The previous phytochemical studies on this herb have led to the isolation of more than a dozen diterpenoid alkaloids.<sup>2</sup> Lappaconitine, one of the main diterpenoid alkaloids, have been developed into a new non-narcotic analgesics drug.<sup>3</sup>

Our further chemical investigation of this plant led to the isolation of two new atisine-type C<sub>20</sub>-diterpenoid alkaloids, named leucostomine A (**1**) and leucostomine B (**2**) (Figure 1). This paper reports the isolation and structural elucidation of these alkaloids.

The 90% EtOH extract of roots of *A. leucostomum*

was treated in the usual procedures<sup>4</sup> to give CH<sub>2</sub>Cl<sub>2</sub>-soluble alkaloid and water-soluble alkaloid fractions.


The water-soluble alkaloid fraction was sequentially subjected to column chromatography over D101



**Figure 1.** Structures of compounds **1** and **2**

macroporous resin, aluminum oxide, and silica gel to give two new compounds.


Compound **1** was obtained as a white solid and gave a positive reaction to Dragendorff's reagent. Its molecular formula was determined as  $C_{22}H_{34}NO_3$  by analysis of its HR-ESI-MS spectrum ( $m/z$  360.2544  $[M]^+$ , calcd. for  $C_{22}H_{34}NO_3$  360.2539). The IR spectrum showed absorption bands for hydroxyl group ( $3411\text{ cm}^{-1}$ ) and an imine bond ( $1706\text{ cm}^{-1}$ ). The  $^1\text{H}$  NMR spectrum showed signals characteristic for an angular methyl group ( $\delta_{\text{H}}$  1.08 s) and an exocyclic methylene group ( $\delta_{\text{H}}$  5.25 and 5.33, br s). Its  $^{13}\text{C}$ -NMR and DEPT spectra showed twenty-two carbon signals including one methyl ( $\delta_{\text{C}}$  26.5), eleven methylenes ( $\delta_{\text{C}}$  21.4, 21.6, 25.2, 26.1, 32.1, 37.3, 42.8, 59.9, 61.8, 66.5, 117.4), six methines ( $\delta_{\text{C}}$  45.9, 46.7, 52.5, 71.8, 77.0, 185.4) and four typical non-oxygenated quaternary carbons ( $\delta_{\text{C}}$  35.6, 40.3, 48.4, 151.2), suggesting compound **1** might be an atisine-type alkaloid.<sup>5</sup> In addition, the resonances at  $\delta_{\text{C}}$  35.6 (s, C-4), 48.4 (s, C-10) and 185.4 (d, C-20) strongly indicated the existence of a quaternary-N center in the molecule,<sup>6</sup> which was also supported by its deshielding effects on N-hydroxyethyl ( $\delta_{\text{H}}$  4.07 and 4.19), H-20 ( $\delta_{\text{H}}$  8.52) and H-19 ( $\delta_{\text{H}}$  3.72 and 3.84, ABq,  $J = 18\text{ Hz}$ ) compared with the values of free atisine-type alkaloids.<sup>7</sup> From the above-mentioned evidence, compound **1** could be assigned as an atisine-type alkaloid possessing a quaternary nitrogen atom. Comparison of the 1D-NMR data (Table) of **1** with those of the known compound atisinium chloride indicated that there was a hydroxyl group at C-11 in **1** instead of the hydrogen group in atisinium chloride,<sup>8</sup> which was also confirmed by the difference of 16 mass units between the two compounds. Furthermore, the existence of hydroxyl group at C-11 was supported by the cross-peaks between H-11 ( $\delta_{\text{H}}$  4.05) and C-9 ( $\delta_{\text{C}}$  52.5), C-10 ( $\delta_{\text{C}}$  48.4) and C-16 ( $\delta_{\text{C}}$  151.2) in the HMBC spectrum. The complete planar structure of **1** was further verified by analysis of the HMBC and  $^1\text{H}$ - $^1\text{H}$  COSY spectra (Figure 2).



**Figure 2.** Key  $^1\text{H}$ - $^1\text{H}$ COSY, HMBC (a) and NOESY (b) correlations of **1**

The configuration of compound **1** was deduced from the coupling constants (Table) and NOESY spectrum (Figure 2). The hydroxyl group at C-11 was assigned a  $\beta$ -orientation based on the coupling constants of H-11 at  $\delta_{\text{H}}$  4.05 (1H, d,  $J = 4.2\text{ Hz}$ ) in the NMR spectra (when the hydroxyl group at C-11 is

$\beta$ -oriented, the coupling of H-11 with H-9 is almost nil and the coupling with H-12 is  $\sim 5$  Hz; in contrast, H-11 show a large coupling ( $\sim 9$ -10 Hz) with H-9 and  $\sim 4$ -5 Hz coupling with H-12<sup>9</sup>). The NOESY correlations of H-11 with H-20 and H-11 with H $_{\alpha}$ -1 also proved a  $\beta$ -orientation of hydroxyl group at C-11. The hydroxyl group at C-15 took  $\beta$ -orientation, since the resonance of C-15 exhibited nearly identical NMR data to those of the known alkaloids, atisium chloride<sup>10</sup> and brunonine<sup>11</sup> whose structure was unambiguously determined by X-ray crystallography. This conclusion was also supported by the cross-peak between H-15 with H $_{\beta}$ -14 in the NOESY experiment. Since the absolute configuration of the atisine nucleus was repeatedly confirmed by the X-ray crystallographic analysis of analogues isolated from species of the same genus,<sup>12</sup> it was proposed that the absolute configuration of the atisine nucleus was retained in **1**. Therefore, the structure of compound **1** was characterized as shown in Figure 1, named as leucostomine A.



**Figure 3.** Key  $^1\text{H}$ - $^1\text{H}$ COSY, HMBC (a) and NOESY (b) correlations of **2**

Compound **2**, a white solid, possessed the molecular formula  $\text{C}_{22}\text{H}_{34}\text{NO}_4$ , which was determined by the analysis of HR-ESI-MS ( $m/z$  376.2494  $[\text{M}]^+$ , calcd. for  $\text{C}_{22}\text{H}_{34}\text{NO}_4$  376.2488). The IR spectrum showed absorption bands for hydroxyl group ( $3377\text{ cm}^{-1}$ ) and an imine bond ( $1672\text{ cm}^{-1}$ ). It was readily recognized that compound **2** was also an atisine-type quaternary ammonium salt alkaloid<sup>13</sup> according to the NMR spectra (Table), which showed the presence of a methyl group ( $\delta_{\text{H}}$  1.10 s,  $\delta_{\text{C}}$  24.9 q), an exocyclic double bond ( $\delta_{\text{H}}$  5.19 and 5.34, br s;  $\delta_{\text{C}}$  115.0 t, 150.7 s), an N-hydroxyethyl group ( $\delta_{\text{H}}$  3.99 and 4.15;  $\delta_{\text{C}}$  58.4 t, 65.0 t) and four typical non-oxygenated quaternary carbon signals ( $\delta_{\text{C}}$  34.2, 44.3, 46.7 and 150.7). The characteristic NMR spectra signals of compound **2** were similar to those of compound **1**, all the differences being due to the hydroxyl group at C-7 in compound **2**. This conclusion was supported by the appearance of a new methine signal at 68.5 ppm, the absence of a methylene signal at 32.1 ppm (compared the  $^{13}\text{C}$ -NMR data with compound **1**), and the downfield shift at C-6 ( $\Delta\delta_{\text{C}}$  7.4 ppm) and C-8 ( $\Delta\delta_{\text{C}}$  4.0 ppm). Moreover, the existence of hydroxyl group at C-7 was confirmed by the cross-peaks between H-7 ( $\delta_{\text{H}}$  3.97) with C-5 ( $\delta_{\text{C}}$  44.0), C-14 ( $\delta_{\text{C}}$  18.5) and C-15 ( $\delta_{\text{C}}$  70.8) in the HMBC spectrum

(Figure 3). The configuration of **2** was determined to be identical to that of **1**, this being supported not only by their almost similar  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR data (Table), but also by the NOESY data (Figure 3). The correlations among H-7, H-5, and H-9 in the NOESY spectrum indicated that H-7 was  $\beta$ -oriented. Furthermore, the upfield shift<sup>14</sup> of C-14 ( $\Delta \delta_{\text{C}}$  7.6 ppm) and C-15 ( $\Delta \delta_{\text{C}}$  6.2 ppm) due to steric compression by the hydroxyl group at C-7 also give the proof of  $\text{H}_{\beta}$ -7. Accordingly, the structure of compound **2** was established to be leucostomine B.

**Table.**  $^1\text{H}$ - and  $^{13}\text{C}$ -NMR data (600 and 150 MHz, resp.) of compounds **1** and **2**

Position	1 in D <sub>2</sub> O		2 in CD <sub>3</sub> OD	
	$\delta_{\text{H}}$ ( <i>J</i> in Hz)	$\delta_{\text{C}}$	$\delta_{\text{H}}$ ( <i>J</i> in Hz)	$\delta_{\text{C}}$
1	$\beta$ 1.78 <sup>a</sup> $\alpha$ 2.22 d (15)	37.3 t	$\beta$ 1.71 m $\alpha$ 2.28 d (13.8)	35.7 t
2	$\alpha$ 1.28 <sup>a</sup> $\beta$ 1.78 <sup>a</sup>	21.4 t	1.75 m	20.1 t
3	$\beta$ 1.57 m $\alpha$ 1.78 <sup>a</sup>	42.8 t	$\beta$ 1.56 <sup>a</sup> $\alpha$ 1.73 m	41.6 t
4	—	35.6 s	—	34.2 s
5	$\beta$ 1.49 <sup>a</sup>	46.7 d	$\beta$ 1.56 <sup>a</sup>	44.0 d
6	$\alpha$ 1.05 m $\beta$ 1.78 <sup>a</sup>	21.6 t	$\alpha$ 1.14 q (13.2) $\beta$ 1.88 dt (3.6, 13.2)	29.0 t
7	$\beta$ 1.28 <sup>a</sup> $\alpha$ 1.83 m	32.1 t	$\beta$ 3.97 br s	68.5 d
8	—	40.3 s	—	44.3 s
9	$\beta$ 1.88 d (1.8)	52.5 d	$\beta$ 1.84 d (1.8)	51.3 d
10	—	48.4 s	—	46.7 s
11	$\alpha$ 4.05 d (4.2)	71.8 d	$\alpha$ 3.98 d (3.6)	70.3 d
12	2.50 dd (1.8, 4.2)	45.9 d	2.41 dd (1.8, 3.6)	45.4 d
13	1.72 m	25.2 t	1.65 m	23.4 t
14	$\alpha$ 0.90 m $\beta$ 1.49 <sup>a</sup>	26.1 t	$\alpha$ 1.27 m $\beta$ 1.35 m	18.5 t
15	$\alpha$ 3.84 br s	77.0 d	$\alpha$ 4.20 br s	70.8 d
16	—	151.2 s	—	150.7 s
17	a 5.25 br s b 5.33 br s	117.4 t	a 5.19 br s b 5.34 br s	115.0 t
18	1.08 s	26.5 q	1.10 s	24.9 q
19	3.72 ABq (18) 3.84 ABq (18)	61.8 t	3.75 ABq (18) 3.82 ABq (18)	60.3 t
20	8.52 br s	185.4 d	8.62 br s	183.8 d
21	4.19 t (4.8)	66.5 t	4.15 t (4.8)	65.0 t
22	4.07 m	59.9 t	3.99 m	58.4 t

<sup>a</sup> Overlapped signals.

## EXPERIMENTAL

### General experimental procedure

Optical rotations were measured on a Perkin-Elmer 341 polarimeter. 1D and 2D NMR spectra were recorded on a Bruker AV 600 NMR spectrometer, and IR spectrum on a ThermoFisher Nicolet 6700 spectrometer (KBr discs,  $\text{cm}^{-1}$ ). HR-ESI-MS were carried out on a Q-TOF micro mass spectrometer (Waters, USA). Silica gel (Qingdao Haiyang Chemical Co., Ltd., China, 200–300 mesh), D-101 macroporous resin (Rohm & Haas), and alumina (Qingdao Haiyang Chemical Co., Ltd., China, 100–200 mesh) were used for column chromatography (CC). TLC plates precoated with silica gel GF254 (Qingdao Haiyang Chemical Co., Ltd., China) were visualized under a UV lamp at 254 nm or by spraying the Dragendorff's reagent or by iodine.

### Plant Material

*A. leucostomum* were collected in Nileke county, Xinjiang Uygur Autonomous Region of China, in August 2014, and were identified by Prof. Qing-Er Yang of the Institute of Botany, Chinese Academy of Sciences. A voucher specimen (C. Ren & L. Wang 735) was deposited in the School of Life Science and Engineering, Southwest Jiaotong University.

### Extraction and isolation

Dried and powder roots of *A. leucostomum* (6.5 kg) were extracted with 95% EtOH three times at room temperature, with each soaking process lasting three days. Evaporation of the solvent under reduced pressure provided a 95% EtOH extract (280.4 g). The extract was treated with 0.5 N hydrochloric acid (2 L) and successively extracted with light petroleum (4×1 L) and EtOAc (4×1 L) to remove non-alkaloid constituents. Then, 28% aqueous ammonia soln. (2 L) was added to the acidic solution to adjust to pH 10. The solutions were extracted with  $\text{CH}_2\text{Cl}_2$  (4×1 L) to afford the  $\text{CH}_2\text{Cl}_2$ -soluble alkaloid. Then, the aqueous layer was purified using chromatography with D101 macroporous resin, and the column contents were eluted with  $\text{H}_2\text{O}$ , 40% EtOH, 60% EtOH, and 95% EtOH, using a gradient. The 60% EtOH elute was concentrated *in vacuo* to afford a residue (13.7 g), which was subjected to alumina column chromatography, using a  $\text{CHCl}_3$ -MeOH (50:1-0:1) as the eluent to separate it into four fractions, A-D. Fraction B (314.6 mg) was chromatographed on silica gel column and eluted with  $\text{CHCl}_3$ /MeOH/concentrated  $\text{NH}_4\text{OH}$  (3:1:1, v/v/v) to afford **1** (32.3 mg) and **2** (25.0 mg) successively.

### Leucostomine A (1)

White solid;  $[\alpha]_{\text{D}}^{20}$  +117 (c 0.45, MeOH); IR (KBr)  $\nu_{\text{max}}$  3411, 2936, 2878, 1706, 1672, 1639, 1452, 1409, 1376, 1110, 1076, 1046, 1018, 864, 580  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (600 MHz,  $\text{D}_2\text{O}$ ) and  $^{13}\text{C}$  NMR (150 MHz,  $\text{D}_2\text{O}$ ): see Table; HR-ESI-MS (m/z): 360.2544  $[\text{M}]^+$ , calcd. for  $\text{C}_{22}\text{H}_{34}\text{NO}_3$  360.2539.

### Leucostomine B (2)

White solid;  $[\alpha]_{\text{D}}^{20}$  +50 (c 1.5, MeOH); IR (KBr)  $\nu_{\text{max}}$  3377, 2935, 2875, 1672, 1637, 1431, 1385, 1112,

1076, 1050, 1017, 987, 857, 794, 636  $\text{cm}^{-1}$ .  $^1\text{H}$  NMR (600 MHz,  $\text{CD}_3\text{OD}$ ) and  $^{13}\text{C}$  NMR (150 MHz,  $\text{CD}_3\text{OD}$ ): see Table; HR-ESI-MS ( $m/z$ ): 376.2494  $[\text{M}]^+$ , calcd. for  $\text{C}_{22}\text{H}_{34}\text{NO}_4$  376.2488.

## ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (81402803), the Research Foundation for Educational Commission of Sichuan Province (15TD0048 and 15ZB0140), and the Fundamental Research Funds for Central Universities (2682015ZT04).

## REFERENCES

1. C.-S. Qian and H.-Y. Chen, 'Flora of China,' Vol. 27, ed. by W.-C. Wang and M. Warnock, Science Press, Beijing, 1979, p 171.
2. (a) J.-M. Yue, J. Xu, Q.-S. Zhao, H.-D. Sun, and Y.-Z. Chen, *J. Nat. Prod.*, 1996, **59**, 277; (b) V. A. Telnov, M. S. Yunusov, N. D. Abdullaev, and M. G. Zhamierashvili, *Khim. Prir. Soedin.*, 1988, **4**, 556; (c) X.-Y. Wei, B.-Y. Wei, and J. Zhang, *Acta Bot. Sin.*, 1996, **38**, 995.
3. V. A. Tel'nov, M. S. Yunusov, and S. Y. Yunusov, *Chem. Nat. Compd.*, 1970, **6**, 598.
4. L. Gao, X. Wei, X. Jin, and L. Yang, *Heterocycles*, 2004, **63**, 1181.
5. (a) N. V. Mody and S. W. Pelletier, *Tetrahedron*, 1978, **34**, 2421; (b) M. S. Yunusov, *Nat. Prod. Rep.*, 1993, **10**, 471; (c) F.-P. Wang, Q.-H. Chen, and X.-Y. Liu, *Nat. Prod. Rep.*, 2010, **27**, 529.
6. (a) S. W. Pelletier and T. N. Oeltmann, *Tetrahedron*, 1968, **24**, 2019; (b) H. K. Desai, B. S. Joshi, S. W. Pelletier, B. Sener, F. Bingöl, and T. Baykal, *Heterocycles*, 1993, **36**, 1081; (c) B.-G. Wang, L. Li, X.-S. Yang, Z.-H. Chen, and X.-J. Hao, *Heterocycles*, 2000, **53**, 1343.
7. (a) X.-J. Hao, M. Node, T. Taga, Y. Miwa, J. Zhou, S. Chen, and K. Fuji, *Chem. Pharm. Bull.*, 1987, **35**, 1670; (b) F. Sun, X.-T. Liang, and D.-q. Yu, *Heterocycles*, 1987, **26**, 19.
8. S. W. Pelletier and N. V. Mody, *J. Am. Chem. Soc.*, 1979, **101**, 492.
9. (a) I. A. Bessonova and S. A. Saidkhodzhaeva, *Chem. Nat. Compd.*, 2000, **36**, 419; (b) A. Ulubelen, H. K. Desai, S. K. Srivastava, B. P. Hart, J. C. Park, B. S. Joshi, S. W. Pelletier, A. H. Mericli, F. Mericli, and R. Ilarslan, *J. Nat. Prod.*, 1996, **59**, 360.
10. P. Wangchuk, J. B. Bremner, Samten, B. W. Skelton, A. H. White, R. Rattanajak, and S. Kamchonwongpaisan, *J. Ethnopharmacol.*, 2010, **130**, 559.
11. W. Deng and W. L. Sung, *Heterocycles*, 1986, **24**, 869.
12. (a) S. W. Pelletier, W. H. De Camp, and N. V. Mody, *J. Am. Chem. Soc.*, 1978, **100**, 7976; (b) S. Ahmad, H. Ahmad, H. U. Khan, A. Shahzad, E. Khan, S. A. A. Shah, M. Ali, A. Wadud, M. Ghufraan, H. Naz, and M. Ahmad, *J. Mol. Struct.*, 2016, **1123**, 441; (c) B. T. Salimov, B. Tashkhodzhaev, M. S. Kodirova, F. M. Tursunkhodzhaeva, and F. N. Dzhakhangirov. *Chem. Nat.*

*Compd.*, 2011, **47**, 261.

13. S. W. Pelletier, F. M. Harraz, M. M. Badawi, S. Tantiraksachai, F.-p. Wang, and S.-y. Chen, *Heterocycles*, 1986, **24**, 1853.
14. S. I. Sakai, T. Okazaki, K. Yamaguchi, H. Takayama, and N. Aimi, *Chem. Pharm. Bull.*, 1987, **35**, 2615.