

HETEROCYCLES, Vol. 105, No. 1, 2022, pp. 352 - 357. © 2022 The Japan Institute of Heterocyclic Chemistry
 Received, 14th March, 2022, Accepted, 1st April, 2022, Published online, 12th April, 2022
 DOI: 10.3987/COM-22-S(R)18


SYNTHESIS AND STEREOCHEMICAL ANALYSIS OF PLANAR CHIRAL NINE-MEMBERED AZA-ORTHO CYCLOPHYNE

Yuuya Kawasaki, Sumire Tanaka, Kazunobu Igawa, and Katsuhiko Tomooka*

Institute for Materials Chemistry and Engineering, Department of Molecular and Material Sciences, and IRCCS Kyushu University, Kasuga, Fukuoka 816-8580, Japan. E-mail: ktomooka@cm.kyushu-u.ac.jp


Abstract – Aza-orthocyclophylene **3b** with an oxy-substituent on the 1⁶ position of the benzene ring was synthesized, and it was revealed that the stereochemical stability of **3b** is consistent with the order of 1,3-steric repulsion between the C2 methylene protons and the oxy-substituent.

Cyclophanes with heteroatom-embedded ansa chains, namely, heterocyclophane **1**, have attracted considerable interest because of their unusual chemical and conformational properties caused by the deformation and strain of the molecule.¹ Among heterocyclophanes, the *ortho*-isomer has received less attention than the *meta*- and *para*-isomers owing to its relatively simple topology and lack of planar chirality in general.² As an exceptional example of planar chiral orthoheterocyclophane, we found that a nine-membered (*E*)-aza-orthocyclophane (*E*)-**2**, that is orthocyclophane with an ansa chain containing a nitrogen functionality along with the (*E*)-alkene, has isolable enantiomers at ambient temperature.³ The key factor




of the stable planar chirality of (*E*)-**2** is the presence of the (*E*)-alkene. Indeed, the diastereomer (*Z*)-**2** has no detectable planar chirality.⁴ Furthermore, we did not observe the presence of isolable enantiomers of nine-membered aza-orthocyclophynes **3a**,⁵ an alkyne analogue of **2**; its chirality would be very labile, unlike that of (*E*)-**2**. In our continuous study on this field of chemistry, quite recently, we have found the significant substituent effect on the stability of planar chirality of the aza-orthocyclophynes; cyclophynes **3b** with oxy-substituent (OR) on the 1⁶ position of the benzene ring has more stable planar chirality than **3a**. Herein, we wish to report the detailed synthesis and stereochemical analysis of **3b**.

We planned to construct the strained nine-membered skeleton of **3b** via intramolecular C–N bond formation of **i**, as shown in Scheme 1. Key intermediate **i** can be prepared from 2,6-dihydroxybenzoic acid (**4**) through stepwise construction of the pentynyl alcohol moiety via i) Migita–Kosugi–Stille coupling, ii) Seyferth–Gilbert reaction, and iii) carboxylation followed by reduction.



Scheme 1. Retrosynthetic analysis of **3b**

At the outset, we synthesized **3b** as shown in Scheme 2. The protection of the carboxylic acid and hydroxyl groups of **4** by acetonide formation, followed by trifluoromethanesulfonylation of the hydroxyl group, afforded **5** in 74% yield (two steps). The Migita–Kosugi–Stille coupling of **5** and allyltributyltin catalyzed by Pd(PPh₃)₄ afforded **6**. Alcoholysis followed by MOMCl treatment of **6** afforded **7** in 93% yield (three steps from **5**). Reduction of the ester moiety using LiAlH₄, followed by the Mitsunobu reaction with TsNHCO₂Me, afforded **8** in 72% yield (two steps). Hydroboration of the alkene moiety of **8**, followed by oxidation of the resulting alcohol into aldehyde, and the reaction with Ohira–Bestmann reagent afforded alkyne **9** in 74% yield (three steps from **8**). Hydroxymethylation of the terminal alkyne of **9** was performed using a carboxylation and reduction sequence in 52% yield over two steps. The resulting alcohol **10** was converted into a methanesulfonyloxy derivative, and cyclization was performed through treatment with K₂CO₃ in DMF to afford the desired **3ba** in 58% yield (two steps). Derivatives of **3ba** with different oxy-functionalities at the 1⁶ position, **3bb** (R = H), **3bc** (R = Ac), and **3bd** (R = Piv) were prepared through the conversion of the MOM-ether moiety of **3ba**.⁶



Scheme 2. Reagents and conditions: (a) acetone, SOCl_2 , cat. DMAP, $(\text{MeOCH}_2)_2$, 0°C to rt; (b) Tf_2O , pyridine, CH_2Cl_2 , 0°C ; (c) allyltributyltin, cat. $\text{Pd}(\text{PPh}_3)_4$, LiCl , THF, 65°C ; (d) K_2CO_3 , MeOH, rt; (e) MOMCl, Hünig base, CH_2Cl_2 , 0°C to rt; (f) LiAlH_4 , Et_2O , 0°C ; (g) TsNHCO_2Me , PPh_3 , bis(2-methoxyethyl) azodicarboxylate, THF, 0°C to rt; (h) $\text{BH}_3 \cdot \text{THF}$ then 38% aq. H_2O_2 , 3 M aq. NaOH, THF, 0°C to rt; (i) Dess-Martin periodinane, CH_2Cl_2 , 0°C ; (j) dimethyl (1-diazo-2-oxopropyl)phosphonate, K_2CO_3 , MeOH, 0°C ; (k) *n*-BuLi then ClCO_2Me , THF, -78°C ; (l) DIBAL, CH_2Cl_2 , -78°C ; (m) MsCl, NEt_3 , CH_2Cl_2 , 0°C ; (n) K_2CO_3 , DMF, 0°C to rt; (o) TMSCl , MeCN/ CH_2Cl_2 , rt; (p) Ac_2O , NEt_3 , cat. DMAP, CH_2Cl_2 , 0°C ; (q) PivCl , NEt_3 , cat. DMAP, CH_2Cl_2 , 0°C .

The ^1H NMR spectrum of **3ba** in CDCl_3 at ambient temperature exhibited two sets of sharp signals of non-equivalent geminal methylene protons, which is in sharp contrast to a similar analysis of **3a** exhibiting broad signals (Figure 1). This result suggests that the planar chirality of **3ba** is more stable than that of **3a**.




Figure 1. ^1H NMR spectra (300 MHz in CDCl_3 at ambient temperature) of **3ba** and **3a**

However, our attempts to separate the enantiomers of **3bb** and **3ba** through HPLC analysis using a chiral stationary phase failed; even with cooling of the column at -20°C , only broad peaks and coalesced plateau-shaped peaks were observed, as shown in Figure 2.⁷ In contrast, similar attempts of **3bc** and **3bd** exhibited

excellent results; the existence of isolable enantiomers was revealed through the HPLC analysis. As shown in Figure 2, both enantiomers of **3bc** were successfully separated at $-20\text{ }^{\circ}\text{C}$. Moreover, both enantiomers of **3bd** were partially separated at $10\text{ }^{\circ}\text{C}$, and baseline separation and measurement of CD signs were accomplished at $-20\text{ }^{\circ}\text{C}$. These results are the first direct observation of the existence of enantiomers of nine-membered aza-orthocyclophynes and suggest that the trend of stereochemical stability would be $\mathbf{3bb} < \mathbf{3ba} < \mathbf{3bc} < \mathbf{3bd}$.⁸




Figure 2. HPLC analysis using a CHIRALPAK IB (a) UV chart of **3ba-3bd** at 10 to $-20\text{ }^{\circ}\text{C}$; (b) CD chart of **3bd** at $-20\text{ }^{\circ}\text{C}$.

Furthermore, **3bd** afforded suitable single crystals to elucidate its detailed structure in the solid state through X-ray crystallographic analysis (Figure 3).⁹ The analysis reveals the packing of two similar conformers of **3bd**; the alkyne-containing ansa chains are located outside the plane of the benzene ring. Therefore, **3bd** possesses planar chirality in the solid state. The average bond angles of C4–C5–C6 and C5–C6–C7 are 161° and 162° , respectively. It means that the alkyne moiety of **3bd** is more bent than that of the previously synthesized **3a**: the C4–C5–C6 and C5–C6–C7 bond angles of **3a** are 166° and 164° , respectively.




Figure 3. Structure of **3**; (a) X-ray analysis of **3bd** (Hydrogen atoms are omitted for clarity. Ellipsoid set at 50% of probability level.); (b) bond angles of **3bd**; (c) bond angles of **3a**.

The former primitive density functional theory calculations revealed that the transition state for the racemization of **3** involves flipping of the benzene ring.¹⁰ The energy barrier for the racemization of **3b** should be higher than that of **3a** because of the substantial 1,3-steric repulsion between the C2 methylene protons and the OR on the 1⁶ position of the benzene ring, as shown in Figure 4. As above mentioned, the trend of stereochemical stability of **3b** (**3bb** < **3ba** < **3bc** < **3bd**) was consistent with the order of 1,3-steric repulsion (OH < OMOM < OAc < OPiv).




Figure 4. 1,3-Steric repulsion between the C2 methylene protons and the OR on the 1⁶ position of benzene ring.

In conclusion, we described the synthesis of a novel nine-membered aza-orthocyclophene **3b** and revealed the presence of isolable enantiomers thereof. Further detailed studies on the relationship between the structures and stereochemical stability of aza-orthocyclophenes, and the preparation of their enantioenriched form by DYASIN¹¹ and synthetic applications are in progress.

SUPPORTING INFORMATION

Supplementary (synthesis of all compounds, ¹H, ¹³C NMR, IR, and HRMS, etc.) data associated with this article can be found, in the online version, at URL: <https://www.heterocycles.jp/newlibrary/downloads/PDFsi/27626/105/1>.

ACKNOWLEDGEMENTS

This research was supported by JSPS KAKENHI Grants (JP20H05677). We thank H. Toshima (Kyushu University) for assistance with the HRMS measurements.

REFERENCES AND NOTES

1. For pioneering works on heterocyclophane, see: R. Fourneau and P. M. Baranger, *Bull. Soc. Chim.*, 1931, **49**, 1161; L. Ruzicka, J. B. Buijs, and M. Stoll, *Helv. Chim. Acta*, 1932, **15**, 1220; K. Ziegler and A. Lüttringhaus, *Justus Liebigs Ann. Chem.*, 1934, **511**, 1; E. W. Spanagel and W. H. Carothers, *J. Am. Chem. Soc.*, 1935, **57**, 935; A. Lüttringhaus, *Justus Liebigs Ann. Chem.*, 1937, **528**, 185. For representative review on heterocyclophane, see: F. Vögtle, G. Pawlitzki, and U. Hahn, 'Modern Cyclophane Chemistry', ed. by R. Gleiter and H. Hopf, Wiley-VCH, Weinheim, 2004, pp. 41–80.
2. For pioneering work on medium-sized orthocyclophynes, see: A. C. Cope and M. W. Fordice, *J. Am. Chem. Soc.*, 1967, **89**, 6187. For representative studies on medium-sized orthocyclophynes, see: G. Seitz, L. Pohl, and R. Pohlke, *Angew. Chem., Int. Ed. Engl.*, 1969, **8**, 447; H. N. C. Wong, P. J. Garratt, and F. Sondheimer, *J. Am. Chem. Soc.*, 1974, **96**, 5604; A. Krebs, J. Odenthal, and H. Kimling, *Tetrahedron Lett.*, 1975, **16**, 4663.
3. K. Tomooka, C. Iso, K. Uehara, M. Suzuki, R. Nishikawa-Shimono, and K. Igawa, *Angew. Chem. Int. Ed.*, 2012, **51**, 10355.
4. K. Machida, Y. Yoshida, K. Igawa, and K. Tomooka, *Chem. Lett.*, 2018, **47**, 186.
5. K. Igawa, T. Kawabata, R. Ni, and K. Tomooka, *Chem. Lett.*, 2013, **42**, 1374.
6. All new compounds were fully characterized by ^1H , ^{13}C NMR, IR, and HRMS analysis: see, Supporting Information for details.
7. Analytical conditions; column: CHIRALPAK IB (0.46 cm \times 25 cm), eluent: hexane/IPA = 30:70, flow rate: 0.5 mL/min, detection: UV and CD (270 nm): see, Supporting Information for details.
8. For representative studies on the shape change of HPLC charts depending on analytical temperature, see: C. Roussel, N. Vanthuyne, M. Bouchekara, A. Djafri, J. Elguero, and I. Alkorta, *J. Org. Chem.*, 2008, **73**, 403; F. Maier and O. Trapp, *Angew. Chem. Int. Ed.*, 2012, **51**, 2985. For representative review on the shape change of HPLC charts depending on analytical temperature, see: C. Wolf, *Chem. Soc. Rev.*, 2005, **34**, 595.
9. Deposition number for compound **3bd** is CCDC 2156588. Selected crystallographic data: triclinic, $P\bar{1}$, $a = 11.50586(17)$ Å, $b = 12.5371(2)$ Å, $c = 17.1225(2)$ Å, $\alpha = 83.5826(14)^\circ$, $\beta = 79.8978(12)^\circ$, $\gamma = 69.6621(16)^\circ$, $V = 2276.59(7)$ Å³, $Z = 2$, $R_1 = 0.0623$, $wR_2 = 0.1725$.: see, Supporting Information for details.
10. K. Igawa, T. Kawabata, K. Uehara, and K. Tomooka, *Heterocycles*, 2015, **90**, 901.
11. K. Igawa, Y. Kawasaki, Y. Ano, T. Kashiwagi, K. Ogawa, J. Hayashi, R. Morita, Y. Yoshioka, K. Uehara, and K. Tomooka, *Chem. Lett.*, 2019, **48**, 726; Y. Kawasaki, R. Kamikubo, Y. Kumegawa, K. Ogawa, T. Kashiwagi, Y. Ano, K. Igawa, and K. Tomooka, *Chem. Commun.*, 2022, **58**, 1605.