


## Supporting Information

# TOTAL SYNTHESIS OF (+)-HETEROPLEXISOLIDE E<sup>‡</sup>

Noriki Kutsumura\*, Akito Kiriseko, Takao Saito\*

Department of Chemistry, Faculty of Science, Tokyo University of Science,  
Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan  
nkutsu@rs.kagu.tus.ac.jp; tsaito@rs.kagu.tus.ac.jp

<sup>‡</sup>This paper is dedicated to Professor Ei-ichi Negishi on celebration of his 77<sup>th</sup> birthday.



## Photocopies of <sup>1</sup>H and <sup>13</sup>C NMR Spectra

S2	(+)-5
S3	(+)-6
S4	(+)-11
S5	(+)-12
S6	(+)-13
S7	(+)-14
S8	(-)-15
S9	11:1 mixture of <b>16</b> and <b>17</b> ( <sup>1</sup> H NMR)
S10	(+)-19
S11	(+)-20
S12	(+)-21
S13	(+)-Heteroplexisolide E ( <b>1</b> ) (CDCl <sub>3</sub> )
S14	(+)-Heteroplexisolide E ( <b>1</b> ) (acetone-d <sub>6</sub> )
S15	(+)-22

