

Supporting Information

SYNTHESIS AND SYNTHESIS-BASED STRUCTURAL ELUCIDATION OF (–)-MACROSPHELIDES J AND K

Hongchan An,^a Seung-Hee Kim,^a Heegyung Kim,^b Kyeojin Kim,^a
Jaehoon Sim,^a Jaebong Jang,^a Seung-Mann Paek,^c Young-Ger Suh,^a
and Hwayoung Yun^{b*}

^a College of Pharmacy, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 151-742, Korea

^b College of Pharmacy, Pusan National University, Geumjeong-gu, Busan 609-735, Korea

^c College of Pharmacy, Research Institute of Pharmaceutical Sciences, Gyeongsang National University, Jinju daero, Jinju, Gyeongnam 660-751, Korea

Email: hyun@pusan.ac.kr

List of the Contents

Experimental procedures ----- S2 – S11

¹H and ¹³C NMR copies of the compounds ----- S12 – S35

General experimental

Unless noted otherwise, all starting materials and reagents were obtained from commercial suppliers and were used without further purification. Tetrahydrofuran and Et₂O were distilled from sodium benzophenone ketyl. Dichloromethane, chloroform, triethylamine, acetonitrile and pyridine were freshly distilled from calcium hydride. All solvents used for routine isolation of products and chromatography were reagent grade and glass distilled. Reaction flasks were dried at 100 °C. Air and moisture sensitive reactions were performed under argon atmosphere. Flash column chromatography was performed using silica gel 60 (230-400 mesh, Merck) with the indicated solvents. Thin-layer chromatography was performed using 0.25 mm silica gel plates (Merck). Optical rotations were measured with JASCO DIP-1000 digital polarimeter at ambient temperature using 100 nm cell of 2 mL capacity. Infrared spectra were recorded on a Perkin-Elmer 1710 FT-IR spectrometer. Mass spectra were obtained with VG Trio-2 GC-MS instrument. High resolution mass spectra were obtained with JEOL JMS-AX 505WA instrument. ¹H and ¹³C NMR spectra were recorded on either a JEOL JNM-GCX 400 or JEOL JNM-LA 300 spectrometer as solutions in deuteriochloroform (CDCl₃). Chemical shifts are expressed in parts per million (ppm, δ) downfield from tetramethylsilane and are referenced to the deuterated solvent (CHCl₃). ¹H-NMR data were reported in the order of chemical shift, multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet and/or multiple resonance), number of protons, and coupling constant in hertz (Hz).

Aldoxime (6). To a solution of aldehyde **8** (6.11 g, 31.5 mmol) in EtOH (50 mL) were added NaOAc (6.20 g, 75.6 mmol) and NH₂OH·HCl (2.63 g, 37.8 mmol) subsequently at ambient temperature. After the reaction mixture was vigorously stirred for 3 h at the same temperature, the mixture was concentrated under reduced pressure. Brine was added to the residue and then extracted with EtOAc. The combined organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 4) to afford 6.30 g (96%) of aldoxime **6** as a white solid: FT-IR (thin film, neat) ν_{\max} 3334, 2934, 1612, 1586, 1514, 1464, 1373, 1303 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 7.96 (s, 1H), 7.33 (d, 1H, *J* = 7.5 Hz), 7.24 (d, 2H, *J* = 8.3 Hz), 6.86 (d, 2H, *J* = 8.6 Hz), 4.43 (m, 2H), 4.07 (m, 1H), 3.78 (s, 3H), 1.33 (d, 3H, *J* = 6.6 Hz); ¹³C-NMR (CDCl₃, 75 MHz) δ 159.3, 155.3, 129.9, 129.5, 129.5, 113.8, 113.8, 71.7, 70.3, 55.3, 19.4; LR-MS (FAB+) *m/z* 210 (M+H⁺).

Isoxazoline (5). To a solution of aldoxime **6** (772 mg, 3.69 mmol) and acrylate **7** (663 mg, 2.46 mmol) in toluene (50 mL) were added aqueous NaOCl (2.93 mL, 4.92 mmol, c.a. 10 %) at ambient temperature. The reaction mixture was stirred for 1 h at the same temperature and diluted with H₂O. The aqueous layer was extracted with EtOAc and the combined organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 2) to afford 922 mg (79%) of isoxazoline **5** as a pale yellow foamy solid: $[\alpha]_{\text{D}}^{20}$ +144 (*c* 2.34, CHCl₃); FT-IR (thin film, neat) ν_{\max} 2959, 1698, 1613, 1514, 1460, 1392, 1335 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 7.27 (d, 2H, *J* = 8.4 Hz), 6.86 (d, 2H, *J* = 8.6 Hz), 5.55 (dd, 1H, *J* = 10.8, 6.4 Hz), 4.44 (q, 1H, *J* = 6.6 Hz), 4.38 (m, 2H), 3.91 (dd, 1H, *J* = 7.9, 5.0 Hz), 3.78 (s, 3H), 3.48 (m, 2H), 3.34 (m, 2H), 2.14 (m, 1H), 2.07 (m, 2H), 1.89 (m, 2H), 1.40 (m, 2H), 1.36 (d, 3H, *J* = 6.6 Hz), 1.18 (s, 3H), 0.97 (s, 3H); ¹³C-NMR (CDCl₃, 100 MHz) δ 168.5, 159.5, 159.3, 129.8, 129.8, 129.6, 113.8, 113.8, 76.8, 70.3, 68.8, 65.2, 55.2, 52.9,

49.0, 47.8, 44.6, 38.1, 37.0, 32.8, 26.4, 20.9, 19.8, 19.0; LR-MS (FAB+) m/z 477 ($M+H^+$); HR-MS (FAB+) calcd for $C_{24}H_{33}N_2O_6S$ ($M+H^+$) 477.2059; found 477.2057.

Allyl ester (4). To a solution of isoxazoline **5** (902 mg, 1.89 mmol) in the presence of 4 Å molecular sieves (473 mg, 250 mg/mmol) in allyl alcohol (50 mL, 0.0379 M), was added $Ti(Oi-Pr)_4$ (1.66 mL, 5.67 mmol) at 130 °C. After the reaction mixture was vigorously stirred for 1 h at same temperature, the mixture was quenched with saturated aqueous NH_4Cl , and then extracted with EtOAc. The combined organic layers were dried over $MgSO_4$ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 3) to afford 508 mg (84%) of allyl ester **4** as a yellow oil: $[\alpha]_D^{20} +17.2$ (*c* 2.18, $CHCl_3$); FT-IR (thin film, neat) ν_{max} 2938, 1743, 1613, 1514, 1458, 1375 cm^{-1} ; 1H -NMR ($CDCl_3$, 300 MHz) δ 7.23 (d, 2H, $J = 8.8$ Hz), 6.85 (d, 2H, $J = 8.8$ Hz), 5.90 (m, 1H), 5.37 - 5.22 (m, 2H), 5.04 (t, 1H, $J = 8.7$ Hz), 4.66 (dt, 2H, $J = 5.9, 1.3$ Hz), 4.45 (q, 1H, $J = 6.6$ Hz), 4.35 (m, 2H), 3.78 (s, 3H), 3.29 (d, 2H, $J = 8.6$ Hz), 1.36 (d, 3H, $J = 6.6$ Hz); ^{13}C -NMR ($CDCl_3$, 100 MHz) δ 169.8, 159.5, 159.3, 131.2, 129.6, 129.6, 129.4, 119.2, 113.8, 113.8, 77.1, 70.4, 68.9, 66.2, 55.2, 36.9, 18.9; LR-MS (FAB+) m/z 320 ($M+H^+$); HR-MS (FAB+) calcd for $C_{17}H_{22}NO_5$ ($M+H^+$) 320.1498; found 320.1502.

PMB deprotected alcohol (9). To a solution of allyl ester **4** (637 mg, 2.00 mmol) in MeCN (80 mL) and H_2O (20 mL) was added ammonium cerium(IV) nitrate (3.29 g, 6.00 mmol) at ambient temperature. After stirring for 3 h at the same temperature, the reaction mixture was quenched with H_2O . The aqueous layer was extracted with EtOAc and the combined organic layers were dried over $MgSO_4$ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 356 mg (90%) of PMB deprotected alcohol **9** as a pale yellow oil: $[\alpha]_D^{20} +128$ (*c*

0.653, CHCl₃); FT-IR (thin film, neat) ν_{\max} 3432, 2982, 1744, 1647, 1439, 1376 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 5.91 (m, 1H), 5.37 - 5.25 (m, 2H), 5.05 (dd, 1H, J = 9.9, 8.3 Hz), 4.71 (m, 1H), 4.67 (dt, 2H, J = 5.9, 1.3 Hz), 3.33 (m, 2H), 2.13 (d, 1H, J = 4.2 Hz), 1.43 (d, 3H, J = 6.6 Hz); ¹³C-NMR (CDCl₃, 100 MHz) δ 170.0, 161.2, 131.1, 119.4, 77.4, 66.4, 63.5, 37.6, 20.7; LR-MS (FAB+) m/z 200 (M+H⁺); HR-MS (FAB+) calcd for C₉H₁₄NO₄ (M+H⁺) 200.0923; found 200.0924.

PMB protected β -hydroxy ester (11). To a solution of acid **10** (529 mg, 2.36 mmol) and alcohol **9** (312 mg, 1.57 mmol) in CH₂Cl₂ (50 mL) were added EDCI (602 mg, 3.14 mmol), DMAP (383 mg, 3.14 mmol) at ambient temperature. After stirring for 2 h at the same temperature, the reaction mixture was quenched with H₂O. The aqueous layer was extracted with CH₂Cl₂ and the combined organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 3) to afford 584 mg (92%) of ester **11** as a yellow oil: $[\alpha]_{\text{D}}^{20}$ +49.6 (*c* 1.01, CHCl₃); FT-IR (thin film, neat) ν_{\max} 2937, 1742, 1613, 1514, 1451, 1376 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 7.20 (d, 2H, J = 8.4 Hz), 6.84 (d, 2H, J = 8.6 Hz), 5.90 (m, 1H), 5.67 (q, 1H, J = 6.6 Hz), 5.36 - 5.24 (m, 2H), 4.88 (dd, 1H, J = 11.0, 7.5 Hz), 4.64 (dt, 2H, J = 5.9, 1.3 Hz), 4.43 (m, 2H), 3.99 (m, 1H), 3.77 (s, 3H), 3.15 (m, 2H), 2.52 (m, 2H), 1.48 (d, 3H, J = 6.6 Hz), 1.25 (d, 3H, J = 6.2 Hz); ¹³C-NMR (CDCl₃, 100 MHz) δ 170.4, 169.4, 159.2, 157.7, 131.2, 130.3, 129.3, 129.3, 119.2, 113.7, 113.7, 77.7, 71.7, 70.5, 66.3, 65.4, 55.2, 42.1, 38.6, 19.6, 17.6; LR-MS (FAB+) m/z 406 (M+H⁺); HR-MS (FAB+) calcd for C₂₁H₂₈NO₇ (M+H⁺) 406.1866; found 406.1872.

β -Hydroxy ester (12). To a solution of PMB ether **11** (379 mg, 0.935 mmol) in CH₂Cl₂ (30 mL) and phosphate buffer solution (3 mL, pH 7.0) was added DDQ (638 mg, 2.81 mmol) at ambient temperature. The reaction mixture was stirred for 3 h and the reaction

mixture was diluted with CH₂Cl₂, filtered under reduced pressure. The organic layer was washed with H₂O and aqueous layer was extracted with CH₂Cl₂. The organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 250 mg (94%) of alcohol **12** as a yellow oil: [α]_D²⁰ +90.4 (*c* 1.51, CHCl₃); FT-IR (thin film, neat) ν_{\max} 3540, 3087, 2978, 1740, 1644, 1452, 1377 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 5.82 (m, 1H), 5.61 (q, 1H, *J* = 6.5 Hz), 5.28 - 5.15 (m, 2H), 4.97 (dd, 1H, *J* = 11.0, 6.9 Hz), 4.57 (d, 2H, *J* = 5.9 Hz), 4.11 (m, 1H), 3.21 (m, 2H), 3.06 (s, 1H), 2.39 (m, 2H), 1.42 (d, 3H, *J* = 6.8 Hz), 1.14 (d, 3H, *J* = 6.2 Hz); ¹³C-NMR (CDCl₃, 100 MHz) δ 171.3, 169.3, 157.6, 131.0, 119.0, 77.5, 66.1, 65.5, 64.2, 43.0, 38.1, 22.5, 17.6; LR-MS (FAB+) *m/z* 286 (M+H⁺); HR-MS (FAB+) calcd for C₁₃H₂₀NO₆ (M+H⁺) 286.1291; found 286.1300.

MEM protected γ -hydroxy α,β -unsaturated ester (14). To a solution of acid **13** (655 mg, 1.85 mmol) and alcohol **12** (376 mg, 1.32 mmol) in CH₂Cl₂ (40 mL) were added EDCI (506 mg, 2.64 mmol), DMAP (322 mg, 2.64 mmol) at ambient temperature. After stirring for 3 h at the same temperature, the reaction mixture was quenched with H₂O. The aqueous layer was extracted with CH₂Cl₂ and the combined organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 523 mg (64%) of ester **14** as a yellow oil: [α]_D²⁰ +25.2 (*c* 1.74, CHCl₃); FT-IR (thin film, neat) ν_{\max} 2937, 1743, 1656, 1613, 1514, 1453, 1377 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 7.18 (d, 2H, *J* = 8.6 Hz), 6.81 (m, 1H), 6.80 (d, 2H, *J* = 8.6 Hz), 5.95 (dd, 1H, *J* = 15.8, 1.5 Hz), 5.86 (m, 1H), 5.61 (q, 1H, *J* = 6.6 Hz), 5.32 - 5.20 (m, 2H), 5.28 (m, 1H), 4.98 (dd, 1H, *J* = 10.6, 7.7 Hz), 4.68 (m, 2H), 4.61 (dt, 2H, *J* = 5.9, 1.3 Hz), 4.45 (s, 2H), 4.27 (m, 1H), 3.74 (s, 3H), 3.72 (m, 1H), 3.57 (m, 2H), 3.44 (m, 2H), 3.30 (s, 3H), 3.19 (m, 2H), 2.59 (m, 2H), 1.41 (d, 3H, *J* = 6.6 Hz), 1.28 (d, 3H, *J* = 6.2 Hz), 1.11 (d, 3H, *J* = 6.2 Hz); ¹³C-NMR (CDCl₃, 100 MHz) δ 169.3, 168.9,

165.0, 159.0, 157.3, 145.6, 131.1, 130.3, 129.1, 129.1, 122.9, 119.1, 113.6, 113.6, 93.9, 92.4, 92.2, 77.6, 76.7, 71.5, 70.7, 67.1, 66.2, 65.7, 58.9, 55.2, 40.7, 38.3, 19.8, 17.7, 15.5; LR-MS (FAB+) m/z 622 (M+H⁺); HR-MS (FAB+) calcd for C₃₁H₄₄NO₁₂ (M+H⁺) 622.2864; found 622.2871.

PMB deprotected secondary alcohol (15). To a solution of PMB ether **14** (523 mg, 0.842 mmol) in CH₂Cl₂ (50 mL) and phosphate buffer solution (5 mL, pH 7.0) was added DDQ (574 mg, 2.52 mmol) at ambient temperature. The reaction mixture was stirred for 2 h and the reaction mixture was diluted with CH₂Cl₂, filtered under reduced pressure. The organic layer was washed with H₂O and aqueous layer was extracted with CH₂Cl₂. The organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 3 : 2) to afford 332 mg (79%) of alcohol **15** as a yellow oil: $[\alpha]_D^{20} +19.0$ (*c* 0.753, CHCl₃); FT-IR (thin film, neat) ν_{\max} 3464, 2936, 1744, 1654, 1454, 1377 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 6.83 (dd, 1H, *J* = 15.7, 6.2 Hz), 5.99 (dd, 1H, *J* = 15.6, 1.4 Hz), 5.90 (m, 1H), 5.65 (q, 1H, *J* = 6.5 Hz), 5.36 - 5.24 (m, 2H), 5.32 (m, 1H), 5.04 (dd, 1H, *J* = 11.1, 7.4 Hz), 4.72 (m, 2H), 4.66 (dt, 2H, *J* = 5.9, 1.3 Hz), 4.19 (m, 1H), 3.92 (m, 1H), 3.74 (m, 2H), 3.53 (t, 2H, *J* = 4.5 Hz), 3.36 (s, 3H), 3.24 (m, 2H), 2.87 (d, 1H, *J* = 5.9 Hz), 2.62 (m, 2H), 1.46 (d, 3H, *J* = 6.6 Hz), 1.32 (d, 3H, *J* = 6.2 Hz), 1.12 (d, 3H, *J* = 6.6 Hz); ¹³C-NMR (CDCl₃, 75 MHz) δ 169.4, 168.9, 164.9, 157.3, 144.4, 131.2, 123.6, 119.1, 94.4, 80.8, 77.7, 71.6, 68.9, 67.5, 67.2, 66.3, 65.8, 58.9, 40.8, 38.4, 19.9, 19.8, 17.7; LR-MS (FAB+) m/z 524 (M+Na⁺); HR-MS (FAB+) calcd for C₂₃H₃₅NO₁₁Na (M+Na⁺) 524.2108; found 524.2092.

16-Membered macrolactone (3). To a solution of alcohol **15** (191 mg, 0.381 mmol) in THF (7 mL) were added Pd(PPh₃)₄ (440 mg, 0.381 mmol) and morpholine (70.0 μ L, 0.762 mmol) at ambient temperature. The mixture was stirred for 12 h at the same temperature

and quenched with 1N HCl and extracted with EtOAc three times. The organic layers were dried over MgSO₄ and concentrated *in vacuo*. This crude mixture was used for next step without further purification. To a solution of acid in toluene (10 mL) were added Et₃N (1.59 mL, 11.4 mmol) and 2,4,6-trichlorobenzoyl chloride (1.19 mL, 7.62 mmol) at 0 °C. The mixture was warmed to ambient temperature and stirred for 2 h. The residue was redissolved in toluene (10 mL) and added slowly to a rapidly stirred solution of DMAP (1.86 g, 15.2 mmol) in toluene (90 mL). The mixture was stirred for 24 h and quenched with H₂O. The reaction mixture was extracted with EtOAc and the organic layers were dried over MgSO₄ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 2 : 1) to afford 102 mg (60%) of ester **3** as a pale yellow oil.

β-Hydroxy ketone (16). To a solution of Ti(Oi-Pr)₄ (0.495 mL, 1.69 mmol) in Et₂O (25 mL) was added EtMgBr (0.450 mL, 1.35 mmol, 3.0 M solution in Et₂O) at ambient temperature. The initially colorless solution turned black-brown and heterogeneous over the course of EtMgBr addition. The reaction mixture was stirred for 2 h at the same temperature. To a solution of isoxazoline **3** (25.0 mg, 56.4 μmol) in Et₂O (2 mL) was added the black-brown reagent at ambient temperature. After stirring for 3 h at the same temperature, quenched with H₂O, the reaction mixture was filtered with silica gel under reduced pressure. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 2 : 1) to afford 16 mg (64%) of ester **16** as a pale yellow oil: $[\alpha]_D^{20} - 73.7$ (*c* 0.520, CH₃OH); FT-IR (thin film, neat) ν_{\max} 3482, 2935, 1731, 1661, 1453, 1372 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 6.67 (dd, 1H, *J* = 15.8, 8.2 Hz), 5.98 (d, 1H, *J* = 15.8 Hz), 5.33 (m, 1H), 5.18 (q, 1H, *J* = 6.7 Hz), 4.94 (m, 1H), 4.68 (m, 2H), 4.55 (m, 1H), 4.07 (m, 1H), 3.68 (m, 2H), 3.52 (t, 2H, *J* = 4.6 Hz), 3.36 (s, 3H), 2.99 (s, 1H), 2.70 (m, 2H), 2.65 (m, 2H), 1.38 (d, 6H, *J* = 6.6 Hz), 1.35 (d, 3H, *J* = 6.4 Hz); ¹³C-NMR (CDCl₃, 75

MHz) δ 204.2, 172.7, 169.9, 164.2, 144.9, 125.6, 93.7, 78.1, 75.9, 72.2, 71.6, 68.6, 67.5, 66.3, 59.0, 42.3, 41.4, 20.1, 17.7, 15.6; LR-MS (FAB+) m/z 447 ($M+H^+$); HR-MS (FAB+) calcd for $C_{20}H_{31}O_{11}$ ($M+H^+$) 447.1866; found 447.1860.

β -Methoxy ketone (17). To a solution of β -hydroxy ketone **16** (12.0 mg, 26.9 μ mol) in $CHCl_3$ (1 mL) were added 2,6-di-*t*-butylpyridine (0.181 mL, 0.807 mmol) and MeOTf (88.5 μ L, 0.807 mmol) at ambient temperature. The mixture was warmed to 80 $^{\circ}C$ and stirred for 12 h. The mixture was quenched with saturated aqueous NH_4Cl , and then extracted with CH_2Cl_2 . The combined organic layers were dried over $MgSO_4$ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 9.0 mg (73%) of allyl ester **17** as a colorless oil: FT-IR (thin film, neat) ν_{max} 2933, 1731, 1662, 1454, 1372 cm^{-1} ; 1H -NMR ($CDCl_3$, 300 MHz) δ 6.69 (dd, 1H, $J = 15.8, 8.2$ Hz), 5.97 (d, 1H, $J = 15.8$ Hz), 5.30 (m, 1H), 5.19 (q, 1H, $J = 6.7$ Hz), 4.98 (m, 1H), 4.68 (m, 2H), 4.19 (dd, 1H, $J = 9.0, 3.9$ Hz), 4.05 (t, 1H, $J = 8.5$ Hz), 3.69 (m, 2H), 3.52 (t, 2H, $J = 4.5$ Hz), 3.36 (s, 3H), 3.36 (s, 3H), 2.67 (m, 2H), 2.66 (m, 2H), 1.37 (d, 6H, $J = 6.2$ Hz), 1.34 (d, 3H, $J = 6.4$ Hz); LR-MS (FAB+) m/z 483 ($M+Na^+$); HR-MS (FAB+) calcd for $C_{21}H_{32}O_{11}Na$ ($M+Na^+$) 483.1842; found 483.1822.

β -Ethoxy ketone (18). To a solution of β -hydroxy ketone **16** (14.0 mg, 31.4 μ mol) in $CHCl_3$ (1 mL) were added 2,6-di-*t*-butylpyridine (0.212 mL, 0.942 mmol) and EtOTf (0.122 mL, 0.942 mmol) at ambient temperature. The mixture was warmed to 80 $^{\circ}C$ and stirred for 12 h. The mixture was quenched with saturated aqueous NH_4Cl , and then extracted with CH_2Cl_2 . The combined organic layers were dried over $MgSO_4$ and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 8.0 mg (54%) of allyl ester **18** as a colorless oil: 1H -NMR ($CDCl_3$, 300 MHz) δ 6.69 (dd, 1H, $J = 15.8, 8.1$ Hz), 5.97 (d, 1H, $J = 15.9$ Hz),


5.29 (m, 1H), 5.20 (q, 1H, $J = 6.7$ Hz), 4.98 (m, 1H), 4.68 (m, 2H), 4.28 (dd, 1H, $J = 9.2$, 3.5 Hz), 4.05 (t, 1H, $J = 8.6$ Hz), 3.68 (m, 2H), 3.52 (t, 2H, $J = 4.5$ Hz), 3.52 (m, 2H), 3.36 (s, 3H), 2.68 (m, 2H), 2.66 (m, 2H), 1.37 (d, 3H, $J = 6.8$ Hz), 1.36 (d, 3H, $J = 6.2$ Hz), 1.34 (d, 3H, $J = 6.4$ Hz), 1.14 (t, 3H, $J = 7.1$ Hz); LR-MS (FAB+) m/z 497 (M+Na⁺); HR-MS (FAB+) calcd for C₂₂H₃₄O₁₁Na (M+Na⁺) 497.1999; found 497.2018.

(–)-Macrosphelide J (1). To a solution of **17** (4.10 mg, 8.91 μmol) in CH₂Cl₂ (0.50 mL) was added trifluoroacetic acid (0.5 mL) at ambient temperature. After stirring for 12 h, the reaction mixture was concentrated *in vacuo* and purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 3.2 mg (97%) of (–)-macrosphelide **J 1** as a colorless oil: $[\alpha]_{\text{D}}^{20} -40.4$ (*c* 0.100, CH₃OH), the natural **1** $[\alpha]_{\text{D}}^{20} -41$ (*c* 0.200, CH₃OH); FT-IR (thin film, neat) ν_{max} 3478, 2929, 1730, 1454, 1374 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 6.79 (dd, 1H, $J = 15.6$, 4.2 Hz), 6.10 (dd, 1H, $J = 15.8$, 1.8 Hz), 5.43 (m, 1H), 5.20 (q, 1H, $J = 6.8$ Hz), 4.94 (m, 1H), 4.25 (dd, 1H, $J = 9.4$, 3.2 Hz), 4.22 (m, 1H), 3.38 (s, 3H), 2.72 (m, 2H), 2.69 (m, 2H), 1.43 (d, 3H, $J = 6.6$ Hz), 1.38 (d, 3H, $J = 6.8$ Hz), 1.35 (d, 3H, $J = 6.4$ Hz); ¹³C-NMR (CDCl₃, 75 MHz) δ 202.7, 172.9, 169.8, 164.2, 144.2, 123.3, 76.6, 75.4, 74.7, 74.6, 68.5, 58.8, 42.0, 41.0, 20.0, 18.2, 15.0; LR-MS (FAB+) m/z 395 (M+Na⁺); HR-MS (FAB+) calcd for C₁₇H₂₄O₉Na (M+Na⁺) 395.1318; found 395.1308.


(–)-Macrosphelide K (2). To a solution of **18** (4.0 mg, 8.44 μmol) in CH₂Cl₂ (0.5 mL) was added trifluoroacetic acid (0.5 mL) at ambient temperature. After stirring for 12 h, the reaction mixture was concentrated *in vacuo* and purified by flash column chromatography on silica gel (EtOAc : *n*-hexane = 1 : 1) to afford 3.2 mg (98%) of (–)-macrosphelide **K 2** as a colorless oil: $[\alpha]_{\text{D}}^{20} -55.8$ (*c* 0.167, CH₃OH), the natural **2** $[\alpha]_{\text{D}}^{20} -59$ (*c* 0.200, CH₃OH); FT-IR (thin film, neat) ν_{max} 3396, 2924, 2853, 1736, 1540, 1459, 1374 cm⁻¹; ¹H-NMR (CDCl₃, 300 MHz) δ 6.79 (dd, 1H, $J = 15.9$, 4.0 Hz), 6.11 (dd, 1H, $J = 15.6$, 1.4 Hz), 5.42

(m, 1H), 5.21 (q, 1H, $J = 6.7$ Hz), 4.93 (m, 1H), 4.34 (dd, 1H, $J = 9.1, 3.4$ Hz), 4.20 (m, 1H), 3.53 (m, 2H), 2.72 (m, 2H), 2.70 (m, 2H), 1.42 (d, 3H, $J = 6.8$ Hz), 1.38 (d, 3H, $J = 6.6$ Hz), 1.35 (d, 3H, $J = 6.2$ Hz), 1.15 (t, 3H, $J = 6.9$ Hz); ^{13}C -NMR (CDCl_3 , 75 MHz) δ 202.7, 173.3, 169.7, 164.2, 144.2, 123.3, 76.6, 75.4, 74.8, 73.0, 68.5, 66.9, 42.1, 41.0, 19.9, 18.2, 15.0, 15.0; LR-MS (FAB+) m/z 409 ($\text{M}+\text{Na}^+$); HR-MS (FAB+) calcd for $\text{C}_{18}\text{H}_{26}\text{O}_9\text{Na}$ ($\text{M}+\text{Na}^+$) 409.1475; found 409.1462. (FAB+) m/z 428 ($\text{M}+\text{Na}^+$); HR-MS (FAB+) calcd for $\text{C}_{21}\text{H}_{27}\text{NO}_7\text{Na}$ ($\text{M}+\text{Na}^+$) 428.1685; found 428.1673.


▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ ¹H-NMR (CDCl₃, 300 MHz)




▼ ¹³C-NMR (CDCl₃, 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ ¹H-NMR (CDCl₃, 300 MHz)




▼ ¹H-NMR (CDCl₃, 300 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)




▼ ¹H-NMR (CDCl₃, 300 MHz)




▼ ¹³C-NMR (CDCl₃, 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 400 MHz)




▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)






▼ ¹H-NMR (CDCl₃, 300 MHz)


▼ ¹³C-NMR (CDCl₃, 75 MHz)




▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)



▼ $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)



▼ $^1\text{H-NMR}$ (CDCl_3 , 300 MHz)



▼ $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz)

