

TWO NEW 2-ARYLBENZOFURANS FROM THE STEMS OF *NICOTIANA TABACUM* AND THEIR BIOACTIVITIES

Pei-Song Yang,^{1,2} Wei Zhang,² Xiao-Feng Shen,² Xin-Lin Wang,² Chao Li,² Xiao-Wei Gong,²
Xu-Dong Zheng,² Dong-Lai Zhu,^{2*} and Jia-Qiang Wang,^{2,*}

¹ School of Chemical Science and Technology, Yunnan University, Kunming 650091, P.R. China; ² Key Laboratory of Tobacco Chemistry of Yunnan Province, China Tobacco Yunnan Industrial Co., Ltd, Kunming 650231, P.R. China; E-mail: 16594939@qq.com; jszxtg_2015@163.com.

Contents of Supporting Information

No.	Contents:	Pages:
Figure S1-S2	^{13}C NMR and ^1H NMR spectra of (1)	3-4
Figure S3-S4	^{13}C NMR and ^1H NMR spectra of (2)	5-6

Figure S1 ^{13}C NMR spectrum of 4-(7-methoxy-3,5-dimethylbenzofuran-2-yl)phenol (**1**)




Figure S2 ^1H NMR spectrum of 4-(7-methoxy-3,5-dimethylbenzofuran-2-yl)phenol (**1**)




Figure S3 ¹³C NMR spectrum of 4-(6-methoxy-3,5-dimethylbenzofuran-2-yl)phenol (2)




Figure S4 Figure S4 ¹H NMR spectrum of 4-(6-methoxy-3,5-dimethylbenzofuran-2-yl)phenol (2)

