

Supporting information for

**Oxidative synthesis of isoxazoline-N-oxide from optically active nitro alcohols**

Takaaki Moriyama,<sup>1</sup> Takuji Kawamoto,<sup>1</sup> Hidemitsu Uno,<sup>2</sup> and Akio Kamimura\*<sup>1</sup>


*<sup>1</sup>Department of Applied Molecular Bioscience, Graduate School of Medicine,  
Yamaguchi University, Ube 755-8611 Japan*



*<sup>2</sup>Department of Chemistry, Graduate School of Science and Engineering, Ehime  
University, Matsuyama 790-8577, Japan*


e-mail: [ak10@yamaguchi-u.ac.jp](mailto:ak10@yamaguchi-u.ac.jp)


Contents



<sup>1</sup> H and <sup>13</sup> C NMR for compounds <b>3a</b> , <b>3b</b> , <b>4a</b> , <b>4b</b> , <b>5a</b> , <b>5b</b> , <b>5c</b> , <b>6a</b> , <b>6b</b> , <b>7a</b> , <b>7b</b> , <b>7c</b> , <b>8a</b> , and <b>9c</b>	S2
ORTEP chart for compound <b>7b</b>	S16
ORTEP chart for compound <b>8a</b>	S16






















































ORTEP chart for compound **7b**ORTEP chart for compound **8a**