

# Supporting Information

## Phosphinic Acid-Promoted Addition Reaction of Isocyanides to 1-Methoxyisochroman Derivatives

Takahiro Soeta\*, Syunsuke Matsuzaki, Yutaka Ukaji\*

*Division of Material Chemistry, Graduate School of Natural Science and Technology,  
Kanazawa University, Kakuma, Kanazawa 920-1192, Japan.*

soeta@se.kanazawa-u.ac.jp; [ukaji@staff.kanazawa-u.ac.jp](mailto:ukaji@staff.kanazawa-u.ac.jp)

### List of Contents

<sup>1</sup> H NMR of <b>3aa</b>	S3
<sup>13</sup> C NMR of <b>3aa</b>	S4
<sup>1</sup> H NMR of <b>3ab</b>	S5
<sup>13</sup> C NMR of <b>3ab</b>	S6
<sup>1</sup> H NMR of <b>3ac</b>	S7
<sup>13</sup> C NMR of <b>3ac</b>	S8
<sup>1</sup> H NMR of <b>3ad</b>	S9
<sup>13</sup> C NMR of <b>3ad</b>	S10
<sup>1</sup> H NMR of <b>3ae</b>	S11
<sup>13</sup> C NMR of <b>3ae</b>	S12
<sup>1</sup> H NMR of <b>3af</b>	S13
<sup>13</sup> C NMR of <b>3af</b>	S14
<sup>1</sup> H NMR of <b>3ag</b>	S15
<sup>13</sup> C NMR of <b>3ag</b>	S16
<sup>1</sup> H NMR of <b>3ah</b>	S17
<sup>13</sup> C NMR of <b>3ah</b>	S18
<sup>1</sup> H NMR of <b>3ba</b>	S19
<sup>13</sup> C NMR of <b>3ba</b>	S20
<sup>1</sup> H NMR of <b>3ca</b>	S21

<sup>13</sup> C NMR of <b>3ca</b>	S22
<sup>1</sup> H NMR of <b>3da</b>	S23
<sup>13</sup> C NMR of <b>3da</b>	S24
<sup>1</sup> H NMR of <b>3ea</b>	S25
<sup>13</sup> C NMR of <b>3ea</b>	S26
<sup>1</sup> H NMR of <b>3fa</b>	S27
<sup>13</sup> C NMR of <b>3fa</b>	S28
<sup>1</sup> H NMR of <b>3ga</b>	S29
<sup>13</sup> C NMR of <b>3ga</b>	S30
<sup>1</sup> H NMR of <b>3ha</b>	S31
<sup>13</sup> C NMR of <b>3ha</b>	S32



























































