

Supporting information

**THREE NEW ANTI-TOBACCO MOSAIC VIRUS PRENYL
CHROMONE DERIVATIVES FROM CASSIA NOMAME**

**Yong Xu,¹ Qi-Li Mi,¹ Yin-Ke Li,^{1,2} Wan-Li Zeng,¹ Qian Gao,¹ Chun-Man Song,¹ Hai-Tao Huang,¹
Hai-Ying Xiang,¹ Xue-Mei Li,¹ Guang-Yu Yang,^{1,2} Qiu-Fen Hu,^{2*} and Jin Wang^{1*}**

¹Key Laboratory of Tobacco Chemistry of Yunnan Province, China Tobacco Yunnan Industrial Co., Ltd, 650231, Kunming, P. R. China. ²Key Laboratory of Chemistry in Ethnic Medicinal Resources, Yunnan Minzu University, Kunming 650031, P. R. China. E-mail: jszxtg_2015@163.com, huqiufena@aliyun.com

Contents of Supporting Information

No.	Contents:	Pages:
Figure S1-S4	¹³ C NMR, ¹ H NMR, HSQC, and HMBC spectra of siamchromone R (1)	3-6
Figure S5- S6	¹³ C NMR and ¹ H NMR spectra of siamchromone S (2)	7-8
Figure S7- S8	¹³ C NMR and ¹ H NMR spectra of siamchromone T (3)	9-10
Figure S9	The antiviral inhibition rates tested by half-leaf method for compounds 1-3	11
Figure S10	The protective effects of compound 3 on TMV	12

Figure S2 Figure S1 ¹H NMR spectrum of siamchromone R (1)




Figure S3 HSQC NMR spectrum of siamchromone R (1)




Figure S4 HMBC NMR spectrum of siamchromone R (1)




Figure S5 ^{13}C NMR spectrum of siamchromone S (2)




Figure S6 ¹H NMR spectrum of siamchromone S (2)




Figure S7 ¹³C NMR spectrum of siamchromone T (3)




Figure S8 ¹H NMR spectrum of siamchromone T (3)




Figure S9 The antiviral inhibition rates tested by half-leaf method for compounds **1-3**

[**a.** siamchromone R (**1**); **b.** siamchromone S (**2**); **c.** siamchromone T (**3**)]




Figure S10 The protective effects of compound **3** on TMV
(**a.** control; **b.** pretreating the tobacco plant with compound **3**)

