

Supporting Information for

Mn(III)-Based Oxidative Cyclization of 2-((2-Arylamino)ethyl)malonates: Synthesis of Quinolines via Dihydroquinolinedicarboxylates

Takayuki Nagashimada, Masahiro Morikawa, Kengo Ohki, and Hiroshi Nishino*

Department of Chemistry, Graduate School of Science, Kumamoto University,

Kurokami 2-39-1, Chûou-Ku, Kumamoto 860-8555, Japan

Fax: +81-96-342-3374; E-mail: nishino@kumamoto-u.ac.jp

Table of Contents

Title and Authors -----	S1
¹ H, ¹³ C, and DEPT Spectra of 1a-t -----	S2–S36
Including	
¹ H, ¹³ C, and DEPT Spectra of 1 (<i>N</i> -Et, <i>N</i> -Bu, 4-NO ₂ C ₆ H ₄) -----	S15–S19
¹ H, ¹³ C, and DEPT Spectra of 2a-s -----	S36–S73
Including	
COSY, HMQC, and HMBC Spectra of 2a -----	S38–S41
COSY, HMQC, and HMBC Spectra of 2k -----	S55–S58
¹ H, ¹³ C, and DEPT Spectra of 3n -----	S63–S64
¹ H, ¹³ C, and DEPT Spectra of 4t -----	S73–S74
¹ H, ¹³ C, and DEPT Spectra of 5-10 -----	S75–S84

























































































































































