

HETEROCYCLES, Vol. 77, No. 1, 2009, pp. 187 - 192. © The Japan Institute of Heterocyclic Chemistry
 Received, 13th May, 2008, Accepted, 27th June, 2008, Published online, 30th June, 2008.
 DOI: 10.3987/COM-08-S(F)17

TRIFLIC IMIDE CATALYZED [3+2] CYCLOADDITION OF ALDIMINES WITH α,α -DIMETHYLALLYLSILANE

Naoya Shindoh,¹ Hidetoshi Tokuyama,² Yoshiji Takemoto,^{1,*} and Kiyosei Takasu^{1,*}


¹Graduate School of Pharmaceutical Sciences, Kyoto University, Yoshida, Sakyo-ku, Kyoto 606-8501, Japan

E-mail: kay-t@pharm.kyoto-u.ac.jp

²Graduate School of Pharmaceutical Sciences, Tohoku University, Aobayama, Sendai 980-8578, Japan


Abstract – Tf₂NH-catalyzed [3+2] cycloaddition of *N*-aryl imines with α,α -dimethylallylsilane to give substituted pyrrolidines is described. We have found the mode of cycloaddition depends upon α -substituents of allylsilanes.

Triflic imide (Tf₂NH), which is recognized as a super Brønsted acid,¹ catalyzes several classes of C-C bond formation reactions.² We have shown Tf₂NH efficiently activates imines in hetero Diels-Alder reaction of imines with 2-siloxydienes.^{3,4} Moreover, we have recently reported Tf₂NH-catalyzed cascade hetero Diels-Alder and hydrogen transfer reaction.⁵ Namely, treatment of *N*-aryl imine (**1**) with allylsilane (**2**) in the presence of a catalytic amount of Tf₂NH afforded substituted quinoline (**4**) in a single operation along with amine (**5**). Notably, in the cascade reaction Tf₂NH activates two mechanistically distinct reactions, such as hetero Diels-Alder reaction of **1** with **2** and hydrogen transfer between produced tetrahydroquinoline (**3**) and imine (**1**) (Scheme 1). During the course of our study, we observed that reaction of **1** with α,α -dimethylallylsilane in the presence of Tf₂NH furnished, unexpectedly, not



Scheme 1. Cascade hetero Diels-Alder and hydrogen transfer reaction catalyzed by Tf₂NH

4). Treatment of a mixture of **8**, *p*-tolylaldehyde (**12**) and *p*-trifluoromethylaniline (**13**) (molar ratio = 1.2 : 1 : 1) with 10 mol% of Tf₂NH in toluene furnished the desired pyrrolidine (**10a**) in 78% yield as a 3 : 2 mixture of diastereomers.



Scheme 3. [3+2] Cycloaddition of **1a** (X = CF₃, R = *p*-tolyl) with **8** in the presence of Tf₂NH

Table 1. Tf₂NH-catalyzed [3+2] cycloaddition of **1** with **8**^a

entry	imine (R, X)	%yield of 10 ^b	dr (<i>cis</i> : <i>trans</i>) ^c
1	1a (<i>p</i> -MeC ₆ H ₄ , CF ₃)	84	3 : 2
2 ^d	1a	76	1 : 1
3 ^e	1b (<i>o</i> -NO ₂ C ₆ H ₄ , CF ₃)	61	3 : 2
4	1c (2-furyl, CF ₃)	44	3 : 2
5	1d (2-thienyl, CF ₃)	60	3 : 2
6 ^f	1d	70	3 : 2
7	1e (2-pyridyl, CF ₃)	0	—
8	1f (<i>p</i> -MeC ₆ H ₄ , NO ₂)	94	1 : 1
9	1g (<i>p</i> -MeC ₆ H ₄ , Br)	47	3 : 2
10	1h ((<i>E</i>)-CH=CH-Ph, CF ₃)	complex mixture	—

^aStandard conditions: **1** (1.0 equiv.), **8** (1.2equiv.), Tf₂NH (10 mol%), in toluene, at 60 °C, for 24 h. ^bYields were calculated based on **1**. ^cDiastereomeric ratio was determined by ¹H NMR.

^dReaction was carried out in 1,2-dichloroethane.

^eHomoallylamine (**11b**) was obtained in 27% yield. ^f20 Mol% of Tf₂NH was used.







Figure 1. Crystal structure of *cis*-**10a** (ORTEP drawing)



Scheme 4. Multicomponent reaction catalyzed by Tf_2NH

In sharp contrast to our previous results,⁵ the mode of cycloaddition depends upon α -substitution of the allylsilane. A plausible mechanism for reactions of imines with allylsilanes is outlined in Figure 2. With α -nonsubstituted allylsilane (**2**), [4+2] cycloaddition took place to give tetrahydroquinolines **3** through a stepwise manner. Namely, $\text{S}_{\text{E}}2'$ reaction of **2** to imine (**1**) in the presence of Tf_2NH would afford β -silyl cation intermediate (**14**), and then intramolecular addition of the aromatic carbon of **14** would promote to furnish [4+2] cycloadduct (**3**) (mode a). If intramolecular addition of the nitrogen atom of **14** takes place, azetidines (**15**) would be produced (mode b). In contrast, in the reaction with α,α -disubstituted allylsilane (**8**), intermediate (**14**) would transform into more stable β -silyl cation (**16**) or siliranium cation (**17**) by 1,2-silyl shift or silacyclopropanation, respectively.¹¹ Then, intramolecular addition of the nitrogen atom of **16** or **17** would afford [3+2] cycloadduct (**10**) (mode c).



Although several studies on cycloaddition reactions of imines with allylsilanes have been reported,^{12–15} studies to control modes of the cycloadditions are quite limited. Akiyama and his co-workers described [3+2] cycloaddition of *N*-sulfonyl imines with triisopropylsilylpropene in the presence of a stoichiometric amount of BF₃-OEt₂,^{13d} whereas *N*-acyl and *N*-aryl imines took place [2+2]¹⁴ and [4+2] cycloadditions,¹⁵ respectively. They concluded *N*-substituent of imines would be a control factor in the selective formation of [2+2], [3+2], or [4+2] cycloadducts. Our abovementioned study indicates α -substitution of allylsilane is one of factors to control the mode of cycloaddition of *N*-aryl imines with allylsilanes.

In conclusion, reaction of imines with α,α -dimethylallylsilane in the presence of Tf₂NH provides substituted pyrrolidines by [3+2] cycloaddition. We found the mode of cycloaddition depends upon α -substituents of allylsilanes. It is noteworthy that, to the best of our knowledge, it is the first example to achieve the catalytic [3+2] cycloaddition of imines with allylsilanes.

ACKNOWLEDGEMENTS

This work was financially supported by a Grant-in-Aid for Scientific Research and Targeted Proteins Research Program from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

REFERENCES

1. H. Yamamoto and M. B. Boxer, *Chimia*, 2007, **61**, 279.
2. Recent representative examples, *see*: (a) K. Ishihara, Y. Hiraiwa, and H. Yamamoto, *Synlett*, 2001, 1851. (b) J. Cossy, F. Lutz, V. Alauze, and C. Meyer, *Synlett*, 2002, 45. (c) L. Zhang and S. A. Kozmin, *J. Am. Chem. Soc.*, 2004, **126**, 10204. (d) K. Inanaga, K. Takasu, and M. Ihara, *J. Am. Chem. Soc.*, 2005, **127**, 3668. (e) K. Takasu, S. Nagao, and M. Ihara, *Adv. Synth. Catal.*, 2006, **348**, 2376. (f) K. Takasu, N. Hosokawa, K. Inanaga, and M. Ihara, *Tetrahedron Lett.*, 2006, **47**, 6053. (g) M. B. Boxer and H. Yamamoto, *J. Am. Chem. Soc.*, 2006, **128**, 48. (h) M. E. Jung and D. G. Ho, *Org. Lett.*, 2007, **9**, 375. (i) K. Takasu, *J. Syn. Org. Chem., Jpn.*, 2008, **66**, 554.
3. A review of C-C bond formation reactions in the presence of a catalytic amount of Brønsted acids, *see*: T. Akiyama, *Chem. Rev.*, 2007, **107**, 5744.
4. K. Takasu, N. Shindoh, H. Tokuyama, and M. Ihara, *Tetrahedron*, 2006, **62**, 11900.
5. N. Shindoh, H. Tokuyama, and K. Takasu, *Tetrahedron Lett.*, 2007, **48**, 4749.
6. L. F. Tietze, T. Neumann, M. Kajino, and M. Pretor, *Synthesis*, 1995, 1003.
7. A. Simion, C. Simion, T. Kanda, S. Nagashima, Y. Mitoma, T. Yamada, K. Mimura, and M. Tashiro, *J. Chem. Soc., Perkin Trans. 1*, 2001, 2071.
8. Spectral data for *cis*-**10a**; Mp 169–170 °C (colorless pillars from MeOH); IR (KBr) 2953, 2854, 1613, 1523, 1324 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.23 (d, *J* = 8.0 Hz, 2H), 7.10 (d, *J* = 8.0 Hz,

- 2H), 7.05 (d, $J = 8.0$ Hz, 2H), 6.67 (d, $J = 8.0$ Hz, 2H), 4.74 (dd, $J = 10.3, 6.3$ Hz, 1H), 2.41 (ddd, $J = 12.3, 6.3, 5.5$ Hz, 1H), 2.40 (s, 3H), 1.78 (ddd, $J = 14.9, 12.3, 10.3$ Hz, 1H), 1.64 (s, 3H), 1.54 (dd, $J = 14.9, 5.5$ Hz, 1H), 0.92 (s, 9H), 0.19 (s, 3H), 0.02 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 147.8, 140.9, 136.4, 129.4, 125.9, 125.2 (q, $^3J_{\text{C,F}} = 3.6$ Hz), 124.0, 123.0 (q, $^1J_{\text{C,F}} = 269.9$ Hz), 117.7 (q, $^2J_{\text{C,F}} = 32.4$ Hz), 116.3, 66.7, 65.6, 39.7, 38.5, 27.5, 27.1, 26.8, 21.1, 17.2, -4.7, -6.0; LRMS (FAB) m/z 447 (M^+), for *trans*-**10b**; ^1H NMR (400 MHz, CDCl_3) δ 7.29 (d, $J = 8.8$ Hz, 2H), 7.09 (m, 4H), 6.63 (d, $J = 8.8$ Hz, 2H), 4.74 (m, 1H), 2.42 (m, 1H), 2.31 (s, 3H), 1.88 (dd, $J = 12.4, 5.1$ Hz, 1H), 1.81 (s, 3H), 1.66 (m, 1H), 1.49 (s, 3H), 0.75 (s, 9H), 0.18 (s, 3H), 0.03 (s, 3H).
9. Crystal data for *cis*-**10a**. $\text{C}_{26}\text{H}_{36}\text{F}_3\text{NSi}$, *monoclinic*, space group $P2_1/n$, $a = 13.8584(3)$ Å, $b = 9.9621(2)$ Å, $c = 17.7342$ Å, $\beta = 91.7284(9)^\circ$, $V = 2447.25(8)$ Å³, $Z = 4$, $D_{\text{calc}} = 1.215$ g/cm³, $R = 0.050$, $R_w = 0.064$, GOF = 1.416.
 10. Y. Yamamoto and N. Asao, [Chem. Rev., 1993, 93, 2207](#).
 11. We have reported [3+2] cycloaddition of acrylates with allylsilanes through formation of similar intermediates, *see*: ref 2f.
 12. For reviews of cycloaddition of allylsilanes, *see*: (a) C. E. Masse and J. S. Panek, [Chem. Rev., 1995, 95, 1293](#). (b) H.-J. Knölker, [J. Prakt. Chem., 1997, 339, 304](#).
 13. (a) J. S. Panek and F. Jain, [J. Org. Chem., 1994, 59, 2674](#). (b) A. Stahl, E. Steckhan, and M. Nieger, [Tetrahedron Lett., 1994, 35, 7371](#). (c) J. V. Schaus, N. Jain, and J. S. Panek, [Tetrahedron, 2000, 56, 10263](#). (d) T. Akiyama, M. Sugano, and H. Kagoshima, [Tetrahedron Lett., 2001, 42, 3889](#).
 14. T. Uyehara, M. Yuuki, H. Masaki, M. Matsumoto, M. Ueno, and T. Sato, [Chem. Lett., 1995, 24, 789](#).
 15. T. Akiyama, M. Suzuki, and H. Kagoshima, [Heterocycles, 2000, 52, 529](#).