

HETEROCYCLES, Vol. 78, No. 8, 2009, pp. 1977 - 1991. © The Japan Institute of Heterocyclic Chemistry  
Received, 10th February, 2009, Accepted, 13th April, 2009, Published online, 13th April, 2009.  
DOI: 10.3987/COM-09-11681

## REACTIVITY OF ARYLNITRILE OXIDES AND C-AROYL-N-PHENYLNITRONES WITH 3-METHYLENEDIHYDRO-(3H)-FURAN-2-ONE AND ITACONIC ANHYDRIDE

Christophe Roussel,<sup>a†</sup> Kabula Ciamala,<sup>a\*</sup> Joël Vebrel,<sup>a</sup> and Claude Riche<sup>b</sup>

<sup>a</sup>Institute UTINAM UMR CNRS 6213, Faculty of Science and Technology, University of Franche-Comté, 16 Route de Gray, F-25030 Besançon

<sup>b</sup>Laboratory of Crystallochemistry, Institute of Chemistry of Natural Products (ICSN) CNRS, 1, Avenue de la terrasse, F-91198 Gif sur Yvette

<sup>†</sup> Current address: Institute of Chemical Sciences and Engineering, Station 6, Swiss Federal Institute of Technology, CH-1015 Lausanne, Switzerland

E-mail: kabula.ciamala@univ-fcomte.fr, christophe.roussel@epfl.ch


**Abstract** - 1,3-dipolar cycloaddition is a powerful route for the synthesis of five-membered heterocycles. The [3+2] cycloadditions of some  $\alpha$ -methylene- $\gamma$ -butyrolactones, namely 3-methylenedihydro-(3H)-furan-2-one (**1**) and itaconic anhydride (**2**) were studied. Their reactions with aryl nitrile oxides (**3**) and C-aroil-N-phenylnitrones (**7**) proceed with complete regioselectivity. From a stereochemical point of view, the addition of aryl nitrile oxides (**3**) leads to the unique spiroheterocycles (**4-5**). Actually, a single stereocenter is generated during the reaction. In the particular case of *p*-nitrophenyl nitrile oxide (**3d**), only the diacidic form (**10d**) of the spiroheterocycle (**5d**) was isolated. In contrast, the addition of C-aroil-N-phenylnitrones (**7**) produces a couple of diastereoisomers, since two stereocenters are generated simultaneously. Nevertheless, the reaction is regioselective and stereospecific leading also to the single spiroheterocycles (**19-20**). The proposed stereochemistry of spiranic compounds (**19d**) and (**20d**) has been corroborated by two single crystal X-ray crystallographic analysis.

## INTRODUCTION

Pharmaceutical industry is much interested in molecules presenting biological activities. Specific syntheses of  $\alpha$ -methylene- $\gamma$ -butyrolactones with cytotoxic, phytotoxic, antitumoral and bactericidal properties have been reviewed.<sup>1</sup> In addition, simple chemical modifications performed at the methylene


group have been carried out and the biological activity of the resulting compounds has been compared to that of the original molecules.<sup>2,3,4,5</sup> For example, the transformation of dehydroleucodine into 11,13-dihydro-dehydroleucodine ( $=\text{CH}_2 \rightarrow \text{CH}_3$ ) decreases cytotoxic effects without significant alteration of the original therapeutic properties.<sup>2</sup> One of the possibilities to chemically modify the methylene group is the [3+2] cycloaddition, affording bicyclic spiranic heterocycles.<sup>6,7,8,9,10,11,12</sup> As an example, cyclopropyl spiranic lactones presenting anticonvulsant properties have been prepared from the reaction of unsaturated lactones with diazomethane.<sup>11</sup>

Since several years we have focused on the reactivity of methylene- $\gamma$ -butyrolactones towards several 1,3-dipoles such as nitrones,<sup>13,14</sup> nitrile oxides<sup>15,16,17</sup> and diazoalkanes.<sup>18,19</sup> We particularly paid attention to the role played by the substituents of the dipolar entities on both chemical reactivity and stereochemistry. We have demonstrated that the reactions of linear C-N-O 1,3-dipoles such as aryl nitrile oxides with achiral  $\gamma$ -methylene- $\gamma$ -butyrolactones proceed with complete regioselectivity. The [3+2] cycloadditions lead to unique spiroheterocycles,<sup>15,16,17</sup> only one stereocenter being generated from the prostereogenic carbon atom of the methylene group (Scheme 1). To the best of our knowledge, except our work,<sup>15</sup> only two references relate the reactivity of aryl nitrile oxides with simple  $\alpha$ -methylene- $\gamma$ -butyrolactones. The [3+2] cycloadditions of phenylnitrile oxide (**3a**) and *p*-tolynitrile oxide (**3b**) with 3-methylenedihydro-(3*H*)-furan-2-one (**1**)<sup>20</sup> and itaconic anhydride (**2**)<sup>21</sup> have been reported (Scheme 1).



**Scheme 1.** [3+2] Cycloadditions of phenyl- (**3a**) and *p*-tolynitrile oxide (**3b**) with 3-methylenedihydro-(3*H*)-furan-2-one (**1**)<sup>15,20</sup> and itaconic anhydride (**2**).<sup>21</sup> a) in refluxing benzene for 5 h,<sup>20</sup> b) in Et<sub>2</sub>O at room temperature for 48 h,<sup>15</sup> c) in Et<sub>2</sub>O at room temperature for few minutes and refluxed for 30 min.<sup>21</sup> Note that recrystallisation of (**5a**) in water affords the diacidic form of the anhydride moiety (**10a**). \* represents a center of chirality.

When going towards bended C-N-O 1,3-dipoles like nitrones, the reactions proceed differently as each reactant possesses a prostereogenic carbon atom. For example, the reactions of  $\gamma$ -methylene- $\gamma$ -butyrolactones with *C*-aroyl-*N*-phenylnitrones (**7**) also present complete regioselectivities but lead to two diastereoisomers (**8**) and (**9**) as depicted in Scheme 2.<sup>13</sup>



**Scheme 2.** [3+2] Cycloadditions of *C*-aryl-*N*-phenylnitrones (**7a-d**) with 3-methylenephthalide (**6**). 24 h in EtOAc at room temperature. The amount of each diastereoisomer depends on Ar. \*represents a chiral center.

In these examples, the diastereoselectivity of the reactions was explained by the different interactions occurring during the establishment of the transition states. Similar studies involving (**7a**) and several  $\gamma$ -methylene- $\gamma$ -butyrolactams (derived from (**6**)) showed the same behaviour.<sup>22</sup> In contrast, the [3+2] cycloadditions of (**7a**) with various  $\alpha$ -methylene- $\gamma$ -butyrolactams (derived from (**1**)) were found to be stereospecific,<sup>23</sup> pointing out the important role played by these secondary interactions.

Herein, we report a reinvestigation of the reactivity of substituted aryl nitrile oxides (**3a-d**), bearing either electron donating or withdrawing groups, with 3-methylenedihydro-(3*H*)-furan-2-one (**1**) and itaconic anhydride (**2**) to complete and generalise the previously reported studies. In addition, we extended our study on the reactivity of *C*-aryl-*N*-phenylnitrones (**7**) towards the dipolarophiles (**1**) and (**2**) with the following aims : (i) to figure out the factors governing the stereoselectivity of the reactions and (ii) to compare the stereochemistry of the [3+2] adducts with those resulting from the reactions of similar  $\alpha$ -methylene- $\gamma$ -butyrolactams with the same 1,3-dipoles.<sup>23</sup>


## RESULTS AND DISCUSSION

### Cycloaddition of aryl nitrile oxides

The [3+2] cycloadditions of 3-methylenedihydro-(3*H*)-furan-2-one (**1**) and itaconic anhydride (**2**) with aryl nitrile oxides (**3a-d**) led to unique adducts corresponding to the expected five-membered spiroheterocycles (**4-5**), except for the reaction involving (**2**) and (**3d**). The reactions proceed with 100% regioselectivity and spiroheterocycles (**4-5**) were isolated as racemates (Scheme 3).

The regiochemistry depicted in Scheme 3 was ascertained by <sup>13</sup>C NMR spectroscopy. The chemical shift of the spiranic carbon atom C<sup>3,5'</sup> (~84 ppm/TMS for (**4**) and ~87 ppm/TMS for (**5**)) is in agreement with vicinity of the isoxazolinic oxygen atom, as already reported.<sup>20</sup>

The reactions were carried out at room temperature in the presence of hydroquinone as reported previously.<sup>13,14,16,17,18,19</sup> The reaction of (1) with aryl nitrile oxides (3) proceeded with good yields (except for (4d)) compared to that in refluxing benzene (71% instead of 41% for (4a)).<sup>20</sup>



**Scheme 3.** [3+2] Cycloadditions of aryl nitrile oxides (3a-d) with 3-methylenedihydro-(3H)-furan-2-one (1) and itaconic anhydride (2). a) 24 h in Et<sub>2</sub>O at room temperature, b) 24 h in EtOAc at room temperature. ((2) is not soluble in Et<sub>2</sub>O). \*represents a center of chirality.


The classic procedure for generating aryl nitrile oxides (3) *in situ* from aryl hydroximoyl chlorides (by addition of triethylamine)<sup>24</sup> used for the reaction involving (1) was not suitable with itaconic anhydride (2). After addition of the base, a degradation occurred, even with DABCO or substituted pyridines (pyridine catalyzes aryl nitrile oxide dimerization).<sup>25</sup> To overcome any degradation, aryl nitrile oxides (3a-d) were first generated with triethylamine in a separate flask and then added to itaconic anhydride (2). This procedure, different from the thermal generation of aryl nitrile oxides from aryl hydroximoyl chlorides,<sup>21</sup> afforded also the adducts (5) in good yields (similar yield for (5a), Schemes 1 and 3), except for (5d).

Another important point concerns the reaction of (2) with *p*-nitrophenyl nitrile oxide (3d) and especially the structure of the resulting product. Compared to the isolated adducts (5a-c), the IR analyses of the obtained product revealed the disappearance of the vibrations at ~1870 and 1790 cm<sup>-1</sup> (C=O anhydride) and the appearance of vibrations in the range between 3305 - 2720 cm<sup>-1</sup> (associated OH) and at 1710 cm<sup>-1</sup> (carboxylic C=O). The <sup>1</sup>H NMR spectrum displayed two exchangeable protons with D<sub>2</sub>O and MS analyses (EI at 70 eV) gave a molecular weight of 294 g/mol, corresponding to the diacidic form of the expected adduct (5d) (276 g/mol). Further comparisons of these spectroscopic data with those of the hydrolysis products of (5a-c), confirmed that (10d) arose from (5d) by an *in situ* opening of the anhydride moiety.

Cycloadditions of *C*-aroyl-*N*-phenylnitrones


Goti and co-workers have reported the reaction of the *C,N*-diphenylnitronone with  $\alpha$ -methylene- $\gamma$ -butyrolactone (**1**) in refluxing benzene or toluene.<sup>12</sup> The cycloaddition afforded two diastereoisomers 5-spirosubstituted isoxazolidines with high selectivity (90:10) or (80:20) respectively (Scheme 4).

The 1,3-dipolar cycloaddition of *C*-benzoyl-*N*-phenylnitronone with  $\alpha$ -methylene- $\gamma$ -butyrolactams exclusively gave a single cycloaddition in toluene *via* an-*endo* transition state<sup>23</sup> (Scheme 4).



**Scheme 4.** [3+2] Cycloadditions of a) *C,N*-diphenylnitronone with 3-methylenedihydro-(3*H*)-furan-2-one<sup>12</sup> (**1**) and 1-methyl-3-methylenepyrrolidin-2-one (**14**).<sup>23</sup> b) *C*-benzoyl-*N*-phenylnitronone (**7a**) with 1-methyl-3-methylenepyrrolidin-2-one (**14**).<sup>23</sup>


For comparison, the reactions of *C*-aroyl-*N*-phenylnitrones (**7a-d**) with 3-methylenedihydro-(3*H*)-furan-2-one (**1**) and itaconic anhydride (**2**) were also investigated. Independently of the nature of the cycloaddition partners, the reactions proceeded with complete regioselectivity and stereospecificity.<sup>26,27</sup> The spiranic adducts (**19**) and (**20**) were isolated as racemates and the stereochemical features were established by NMR and X-ray crystallographic analysis (Scheme 5).



**Scheme 5.** [3+2] Cycloadditions of *C*-aroyl-*N*-phenylnitrones (**7a-d**) with 3-methylenedihydro-(3*H*)-furan-2-one (**1**) and itaconic anhydride (**2**). 24 h in EtOAc at room temperature.


Again, the reactions were carried out at room temperature in the presence of hydroquinone. Whereas the [3+2] cycloaddition of itaconic anhydride (**2**) with *C*-aroyl-*N*-phenylnitrones (**7**) proceeded in good yields, 3-methylenedihydro-(3*H*)-furan-2-one (**1**) remained less reactive, as already observed with ethyl diazoacetate.<sup>18,19</sup>

The regioselectivity of these [3+2] cycloadditions were accessed by <sup>1</sup>H NMR spectroscopy. The <sup>1</sup>H NMR spectra of (**19**) and (**20**) display an ABX system corresponding to the protons attached on the isoxazolidinic ring (H<sup>4'a</sup>, H<sup>4'b</sup> and H<sup>3'</sup>). This pattern is in favour of formation of regioisomers (**19**) and (**20**). For the hypothetical regioisomers (**19'**) and (**20'**), the proton H<sup>3'</sup> should appear as a singlet. Further confirmation of the regiochemistry came from <sup>13</sup>C-<sup>1</sup>H HSQC NMR. As shown in Figure 1, the spiranic carbon atom C<sup>3,5'</sup> was easily identified at 80 ppm/TMS as it is the sole *sp*<sup>3</sup> quaternary carbon atom which therefore, do not present any correlation with protons. Extended <sup>13</sup>C NMR studies gave a chemical shift for the spiranic carbon atoms between ~82 and ~84 ppm for (**19**) and (**20**), in agreement with the proximity of the isoxazolidinic oxygen atom<sup>9,12,13</sup> (Figure 2). In addition, <sup>13</sup>C-<sup>1</sup>H HSQC NMR performed on spiroheterocycles (**19**) allows the correct attribution of the chemical shifts of C<sup>4</sup> and C<sup>4'</sup>. Whereas these chemical shifts are rather similar, the main difference between C<sup>4</sup> and C<sup>4'</sup> stems from the spin multiplicity of their respective born protons. Actually, C<sup>4</sup> bears the AB part of an ABXX' system (H<sup>4a</sup>, H<sup>4b</sup> and H<sup>5a</sup>, H<sup>5b</sup>) and C<sup>4'</sup>, the AB part of an ABX system (H<sup>4'a</sup>, H<sup>4'b</sup> and H<sup>3'</sup>). Based on figure 1, the respective chemical shifts of C<sup>4</sup> and C<sup>4'</sup> were found to be ~34 and ~37 ppm/TMS respectively. In contrast, for spiroadducts (**20**), C<sup>4</sup> and C<sup>4'</sup> could be easily distinguished by <sup>13</sup>C NMR. Since C<sup>4</sup> is directly linked to the anhydride carbonyl, it is therefore more deshielded (C<sup>4</sup> and C<sup>4'</sup> at ~41.0 and ~37 ppm/TMS respectively).




**Figure 1.** <sup>13</sup>C-<sup>1</sup>H HSQC NMR spectrum of (**19b**)

From a stereochemical point of view, the reactions generate two stereocenters  $C^{3,5'}$  and  $C^{3'}$  from the two prostereogenic carbon atoms of (**1-2**) and (**7**), respectively. Indeed, two possible diastereoisomers as racemates (the two dipolarophilic faces are equivalent) could be envisaged if the stereochemistry of the nitrones (**7**) is conserved. As confirmed by the X-ray studies performed on both (**19d**) and (**20d**), the retention of the original relative (*Z*) configuration of the nitrones (**7**)<sup>28,29</sup> is in favour of a concerted mechanism<sup>30</sup> (Figure 2). Thus, according to the Woodward-Hoffmann rules,<sup>31</sup> the stereospecificity of the reaction could be figured out from *endo*- $C=O$  and *exo*- $C=O$  transition states as depicted in scheme 6.<sup>12,13</sup>



**Figure 2.** Ortep drawing of molecule (**19d**) and (**20d**), drawn by program *PLATON*.<sup>32</sup> Displacement ellipsoids are shown at the 20% probability level. Comparison of torsion angle values shows that the isoxazolidinic ring adopts the same half-chair conformation, in which atoms N2 and C3 are respectively deviated, by  $-0.347(3)$  and  $0.293(3)$  Å from the mean plane of the other three atoms in (**19d**), by  $-0.288(2)$  and  $0.329(2)$  Å, in (**20d**).



**Scheme 6.** *Endo*- $C=O$  and *exo*- $C=O$  transition states for the reaction of *C*-aroyle-*N*-phenylnitrones (**7a-d**) with 3-methylenedihydro-(3*H*)-furan-2-one (**1**) and itaconic anhydride (**2**).

As shown in Figure 2, the *anti*-periplanar relationship between the dipolar and lactonic carbonyl groups is in favour of the *endo*-C=O approach, scheme 6. Whereas the *endo*-C=O transition state provides some secondary orbital interactions between the *N*-phenyl group of the nitron and the carbonyl group of the lactone, the *exo*-C=O transition state does not afford such stabilising effects.<sup>12</sup> Nevertheless, the difference between these two transition states could not explain the stereospecificity of the reaction. For example, the [3+2] cycloaddition of *C,N*-diphenylnitron with (1), is stereoselective, the *endo*-C=O product being the major one.<sup>12</sup> Indeed, as the two dipolarophiles under study lead to the same diastereoisomers, the stereospecificity of our reactions should stem from the presence of the supplementary C=O born by the *C*-aroyl-*N*-phenylnitron compared to *C,N*-diphenylnitron. As shown in scheme 6, the *exo*-C=O transition state presents also the electrostatic repulsion of the two carbon-oxygen dipoles born by the reactants. This electrostatic repulsion could then be considered as the main factor that orientates the reaction through the generation of the *endo*-C=O spiroadducts (19) and (20) only.

## CONCLUSION

We revisited the reactivity of aryl nitrile oxides with simple  $\alpha$ -methylene- $\gamma$ -butyrolactones, using substituted 1,3-dipoles to complete and generalise the previous reported studies. The reactions involving both 3-methylene-(3*H*)-furan-2-one and itaconic anhydride lead to the expected spiroheterocycles, except for the reaction of itaconic anhydride with *p*-nitrophenylnitrile oxide which affords directly the diacidic form of the spiranic compound. Whatever the dipolarophile, the 1,3-dipolar cycloadditions are entirely regioselective, the observed regioselectivity being in total agreement with other studies reported previously. The [3+2] cycloadditions of the same dipolarophiles (1) and (2) with *C*-aroyl-*N*-phenylnitrones are again fully regioselective and stereospecific. Based on X-ray diffraction studies, the stereospecificity of the reaction was explained by the repulsion of the lactonic and dipolar oxygen atoms during the establishment of the transition state.

## EXPERIMENTAL

The cycloadditions were carried out under nitrogen atmosphere using standard Schlenk techniques. TLC plates, DC-Alufolien Kieselgel 60 F<sub>254</sub>, were from Merck. Melting points were measured on an Electrothermal IA 9200 and are not corrected. <sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on Bruker AC 200 [200 MHz (<sup>1</sup>H) and 50 MHz (<sup>13</sup>C)]. HSQC NMR spectra were recorded on a Bruker Avance 300 spectrometer [300 MHz (<sup>1</sup>H) and 75 MHz (<sup>13</sup>C)]. Chemical shifts were measured relative to TMS. IR spectra were recorded on a Bio-Rad FTS-7 spectrometer. Mass spectrometry experiments were run on a NERMAG R 1010 H apparatus under electronic impact at 70 eV. Elemental analyses (C, H, N) were conducted on a Leco Elemental CHN 900; values were in satisfactory agreement with the calculated ones (0.30%).

### Starting materials.

3-Methylenedihydro-(3*H*)-furan-2-one (**1**) was obtained from dihydro-(3*H*)-furan-2-one following the literature procedure<sup>33</sup> (xylene was replaced by anhydrous toluene in the second step). (**1**) was further purified by column chromatography<sup>34</sup> or by distillation under reduced pressure.<sup>35</sup> Itaconic anhydride (**2**) was purchased from Aldrich. Arylnitrile oxides (**3a-d**), were obtained *in situ* by addition of triethylamine to arylhydroximoyl chlorides,<sup>24</sup> synthesised according to the literature.<sup>36,37</sup> *C*-Aroyl-*N*-phenylnitrones (**7a-d**) were obtained from *p*-substituted-2-bromoacetophenone pyridinium salts.<sup>38</sup> *p*-Substituted-2-bromoacetophenone pyridinium salts were synthesised as follows: To 1 equiv. of *p*-substituted-2-bromoacetophenones dissolved in 100 mL of dry acetone was slowly added 1 equiv. of pyridine. The resulting mixture was stirred vigorously for 10 min. The formed precipitate was filtered, washed with 50 mL of dry acetone, dried and used without further purification (yields: 77%, 70%, 55%, 89% for (**a-d**) respectively).

### Cycloaddition of 3-methylenedihydro-(3*H*)-furan-2-one (**1**) with aryl nitrile oxides (**3a-d**)

In a Schlenk tube containing 3-methylenedihydro-(3*H*)-furan-2-one (**1**) (0.49 g, 5 mmol) and 0.05 g (0.45 mmol) of hydroquinone in anhydrous Et<sub>2</sub>O (20 mL), was added under stirring, arylhydroximoyl chlorides (7 mmol). After cooling the mixture at 0 °C, 1 mL (7 mmol) of Et<sub>3</sub>N was added dropwise and after purging with dry N<sub>2</sub>, the reaction was carried out at room temperature for 24 h. After reaction, the solution was first filtered on a glass frit to get rid of triethylammonium chloride (resulting from the dechlorination of arylhydroximoyl chlorides). The filtrate was evaporated under reduced pressure leading to a solid. The two collected solids were dissolved in 20 mL of CH<sub>2</sub>Cl<sub>2</sub> and washed with distilled water (3×20 mL). The organic phase was dried on Na<sub>2</sub>SO<sub>4</sub> and evaporated under reduced pressure; the crude products were purified by recrystallisation from EtOH.

### 3'-Phenylspiroisoxazolino[5':3]dihydro-(3*H*)-furan-2-one (**4a**)

White solid; yield 0.76 g (71%, lit., 41%<sup>20</sup>); mp 132-134 °C (lit., 132-133 °C<sup>20</sup>); IR (KBr) 1775, 1580 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ/TMS 2.30-2.80 (m, 2H), 3.30-4.00 (AB, 2H *J*=16.9 Hz), 4.35-4.60 (m, 2H), 7.30-7.75 (m, 5H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ/TMS 35.1, 41.9, 65.8, 84.2, 126.7-130.5, 155.9, 173.9; Anal. Calcd for C<sub>12</sub>H<sub>11</sub>NO<sub>3</sub>: C, 66.35; H, 5.10; N, 6.45. Found: C, 66.58; H, 4.91; N, 6.57.

### 3'-(4-Methylphenyl)spiroisoxazolino[5':3]dihydro-(3*H*)-furan-2-one (**4b**)

White solid; yield 0.74 g (64%, lit., 56%<sup>15</sup>); mp 128 °C (lit., 128 °C<sup>15</sup>); IR (KBr) 1775, 1580 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ/TMS 2.40 (s, 3H), 2.30-2.80 (m, 2H), 3.30-4.00 (AB, 2H, *J*=16.8 Hz), 4.30-4.60 (m, 2H), 7.20 (d, 2H, *J*=8.0 Hz), 7.55 (d, 2H, *J*=8.0 Hz); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ/TMS 21.3, 35.0, 42.0, 65.8, 84.0, 125.3-140.8, 155.8, 174.0; Anal. Calcd for C<sub>13</sub>H<sub>13</sub>NO<sub>3</sub>: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.69; H, 5.61; N, 6.12.

### 3'-(4-Methoxyphenyl)spiroisoxazolino[5':3]dihydro-(3*H*)-furan-2-one (**4c**)

Colorless solid; yield 0.62 g (50%); mp 137-139 °C; IR (KBr) 1775, 1600 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ/TMS

2.30-2.80 (m, 2H), 3.85 (s, 3H), 3.25-4.00 (AB, 2H,  $J=16.8$  Hz), 4.35-4.60 (m, 2H), 6.95 (d, 2H,  $J=8.8$  Hz), 7.60 (d, 2H,  $J=8.8$  Hz);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  35.0, 42.1, 55.2, 65.8, 83.9, 114.0-161.3, 155.5, 174.1; Anal. Calcd for  $\text{C}_{13}\text{H}_{13}\text{NO}_4$ : C, 63.15; H, 5.30; N, 5.67. Found: C, 62.89; H, 5.41; N, 5.79.

### **3'-(4-Nitrophenyl)spiroisoxazolino[5':3]dihydro-(3H)-furan-2-one (4d)**

Yellow solid; yield 0.34 g (26%); mp 208-209 °C, IR (KBr) 1785, 1610  $\text{cm}^{-1}$ ,  $^1\text{H}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  2.60-2.90 (m, 2H), 3.70-4.10 (AB, 2H,  $J=17.5$  Hz), 4.40-4.70 (m, 2H), 8.00 (d, 2H,  $J=8.8$  Hz), 8.35 (d, 2H,  $J=8.8$  Hz);  $^{13}\text{C}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  35.2, 42.0, 66.8, 86.8, 124.8-135.9, 156.3, 174.6; Anal. Calcd for  $\text{C}_{12}\text{H}_{10}\text{N}_2\text{O}_5$ : C, 54.97; H, 3.84; N, 10.68. Found: C, 55.23; H, 3.89; N, 10.53.

### *Cycloaddition of itaconic anhydride (2) with aryl nitrile oxides (3a-d)*

To a magnetically stirred solution of 3.5 mmol of arylhydroximoyl chlorides in 15 mL of EtOAc at 0 °C, was added 0.4 mL of  $\text{Et}_3\text{N}$  (2.9 mmol). After 5 min, the solution was added dropwise through a glass fritted disk (to get ride of the formed  $\text{Et}_3\text{NHCl}$ ) to a solid mixture of 0.28 g (2.5 mmol) of itaconic anhydride (**2**) and 0.05 g (0.45 mmol) of hydroquinone contained in a Schlenk tube cooled at -15 °C. The glass fritted disk was rinsed with 5 mL of EtOAc. After purging with nitrogen, the reaction was stirred at room temperature for 24 h. After reaction, the solvent was evaporated under reduced pressure; the crude products were purified by recrystallisation from  $\text{CH}_2\text{Cl}_2$ - $\text{Et}_2\text{O}$  (1:1).

### **3'-Phenylspiroisoxazolino[5':3]succinic anhydride (5a)**

Colorless solid; yield 0.42 g (73%, lit., 75%<sup>21</sup>); mp 146-148 °C (lit., 147-149 °C<sup>21</sup>); IR (KBr) 1865, 1800, 1605  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  3.45-3.90 (AB, 2H,  $J=19.2$  Hz), 3.90-4.30 (AB, 2H,  $J=17.9$  Hz), 7.45-7.85 (m, 5H);  $^{13}\text{C}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  42.5, 45.0, 87.5, 128.5-132.3, 158.3, 169.4, 172.9; Anal. Calcd for  $\text{C}_{12}\text{H}_9\text{NO}_4$ : C, 62.34; H, 3.92; N, 6.06. Found: C, 62.48; H, 4.19; N, 5.81.

### **3'-(4-Methylphenyl)spiroisoxazolino[5':3]succinic anhydride (5b)**

Colorless solid; yield 0.39 g (64%); mp 170-172 °C; IR (KBr) 1870, 1785, 1600  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  2.40 (s, 3H), 3.45-3.85 (AB, 2H,  $J=19.1$  Hz), 3.85-4.25 (AB, 2H,  $J=17.8$  Hz), 7.35 (d, 2H,  $J=7.9$  Hz), 7.65 (d, 2H,  $J=7.9$  Hz);  $^{13}\text{C}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  22.1, 42.5, 45.1, 87.3, 127.1-142.6, 158.2, 169.4, 172.9; Anal. Calcd for  $\text{C}_{13}\text{H}_{11}\text{NO}_4$ : C, 63.67; H, 4.52; N, 5.71. Found: C, 63.60; H, 4.61; N, 5.69.

### **3'-(4-Methoxyphenyl)spiroisoxazolino[5':3]succinic anhydride (5c)**

White solid; yield 0.39 g (60%); mp 169-171 °C; IR (KBr) 1875, 1795, 1605  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  3.35-3.80 (AB, 2H,  $J=19.0$  Hz), 3.85 (s, 3H), 3.80-4.20 (AB, 2H,  $J=18.0$  Hz), 7.00 (d, 2H,  $J=8.9$  Hz), 7.70 (d, 2H,  $J=8.9$  Hz);  $^{13}\text{C}$  NMR (*acetone- $d_6$* )  $\delta/\text{TMS}$  42.5, 45.2, 56.5, 87.2, 115.8-163.3, 157.8, 169.5, 173.0; Anal. Calcd for  $\text{C}_{13}\text{H}_{11}\text{NO}_5$ : C, 59.77; H, 4.24; N, 5.36. Found: C, 59.94; H, 4.19; N, 5.42.

*Alkaline treatment of 3'-arylspiroisoxazolino[5':3]succinic anhydride (5a-c)*

To a solution of spiroadducts **5a-c** (0.25 mmol) in 4.1 mL of EtOH was added 0.9 mL of NaOH (2 M). The mixture was refluxed for 30 min, poured into an ice-water mixture (100 mL) and acidified with hydrochloric acid (or sulfuric acid). After extraction with CH<sub>2</sub>Cl<sub>2</sub> (20 mL), the organic phase was washed with distilled water (2×20 mL), dried (Na<sub>2</sub>SO<sub>4</sub>) and evaporated *in vacuo*. The crude products were purified by recrystallization from distilled water.

**Isoxazolino-3'-phenyl-5'-carboxy-5'-ethanoic acid (10a)**

Beige solid; yield 0.057 g (92%); mp 206-207 °C (lit., 197 °C<sup>21</sup>); IR (KBr) 2720-3285, 1705, 1605 cm<sup>-1</sup>; <sup>1</sup>H NMR (*acetone-d*<sub>6</sub>) δ/TMS 3.00-3.40 (AB, 2H, *J*=16.7 Hz), 3.50-4.20 (AB, 2H, *J*=17.2 Hz), 7.30-7.90 (m, 5H), 9.30 (s, 2H); <sup>13</sup>C NMR (*acetone-d*<sub>6</sub>) δ/TMS 41.0, 44.0, 86.6, 127.5-131.0, 157.4, 171.2, 171.3; Anal. Calcd for C<sub>12</sub>H<sub>11</sub>NO<sub>5</sub>: C, 57.83; H, 4.45; N, 5.62. Found: C, 57.99; H, 4.33; N, 5.73.

**Isoxazolino-3'-(4-methylphenyl)-5'-carboxy-5'-ethanoic acid (10b)**

Beige solid; yield 0.062 g (94%); mp 192-194 °C; IR (KBr) 2810-3280, 1705, 1610 cm<sup>-1</sup>; <sup>1</sup>H NMR (*acetone-d*<sub>6</sub>) δ/TMS 2.40 (s, 3H), 3.00-3.40 (AB, 2H, *J*=16.8 Hz), 3.50-4.20 (AB, 2H, *J*=17.5 Hz), 7.30 (d, 2H, *J*= 8.0 Hz), 7.65 (d, 2H, *J*= 8.0 Hz), 8.35 (s, 2H); <sup>13</sup>C NMR (*acetone-d*<sub>6</sub>) δ/TMS 21.3, 41.0, 44.1, 86.4, 127.5-141.2, 157.3, 171.2, 171.3; HRMS (EI, 70 eV) *m/z* 263 (I= 0.50%) M<sup>+</sup>, 245 (I= 4.24%) M<sup>+</sup>-(H<sub>2</sub>O), 218 (I= 0.50) M<sup>+</sup>-(COOH), 201 (I= 4.40%) M<sup>+</sup>-(COOH) -(OH), 160 (I= 100%) M<sup>+</sup>-(COOH) -(OH) -(CH=C=O), 131 (I= 26,03%) M<sup>+</sup>-(COOH) -(OH) -(CH=C=O) -(CHO)<sup>a</sup>; Anal. Calcd for C<sub>13</sub>H<sub>13</sub>NO<sub>5</sub>: C, 59.31; H, 4.98; N, 5.32. Found: C, 59.16; H, 5.07; N, 5.39.

a) determined according to the literature<sup>39</sup>

**Isoxazolino-3'-(4-methoxyphenyl)-5'-carboxy-5'-ethanoic acid (10c)**

Beige solid; yield 0.065 g (93%); mp 199-200 °C; IR (KBr) 2760-3280, 1705, 1605 cm<sup>-1</sup>; <sup>1</sup>H NMR (*acetone-d*<sub>6</sub>) δ/TMS 3.00-3.40 (AB, 2H, *J*=16.9 Hz), 3.85 (s, 3H), 3.50-4.20 (AB, 2H, *J*=17.4 Hz), 7.00 (d, 2H, *J*= 8.8 Hz), 7.70 (d, 2H, *J*= 8.8 Hz), 9.10 (s, 2H); <sup>13</sup>C NMR (*acetone-d*<sub>6</sub>) δ/TMS 41.0, 44.2, 55.7, 86.2, 114.9-162.1, 156.9, 171.2, 171.4; Anal. Calcd for C<sub>13</sub>H<sub>13</sub>NO<sub>6</sub>: C, 55.91; H, 4.69; N, 5.02. Found: C, 56.10; H, 4.71; N, 4.97.

**Isoxazolino-3'-(4-nitrophenyl)-5'-carboxy-5'-ethanoic acid (10d)<sup>b</sup>**

Yellow solid; yield 0.15 g (20%); mp 110-112 °C; IR (KBr) 2720-3305, 1710, 1605 cm<sup>-1</sup>; <sup>1</sup>H NMR (*acetone-d*<sub>6</sub>) δ/TMS 3.10-3.45 (AB, 2H, *J*=16.9 Hz), 3.70-4.30 (AB, 2H, *J*=17.5 Hz), 8.05 (d, 2H, *J*= 8.9 Hz), 8.40 (d, 2H, *J*= 8.9 Hz); <sup>13</sup>C NMR (*acetone-d*<sub>6</sub>) δ/TMS 41.7, 44.2, 88.5, 123.6-136.7, 157.3, 171.9; Anal. Calcd for C<sub>12</sub>H<sub>10</sub>N<sub>2</sub>O<sub>7</sub>: C, 48.99; H, 3.43; N, 9.52. Found: C, 49.26; H, 3.49; N, 9.46.

b) by *in situ* ring opening during cycloaddition

*Cycloadditions of 3-methylenedihydro-(3H)-furan-2-one (1) and itaconic anhydride (2) with C-royl-N-phenylnitrones (7a-d)*

In a Schlenk tube were added 5 mmol of methylene-lactone (**1-2**), 5 mmol of C-royl-N-phenylnitrones (**7a-d**), 0.05 g (0.45 mmol) of hydroquinone and 20 mL of EtOAc. The mixture was stirred at room temperature for 24 h under nitrogen atmosphere and the solvent removed under reduced pressure leading to a crude oil. For spiroheterocycles (**19**), EtOH (20 mL) was added to the crude oil and the mixture subjected to ultrasonication. The resulting solids were recrystallised from EtOH. For spiroheterocycles (**20**), Et<sub>2</sub>O (20 mL) was added to the crude oil and the mixture was subjected to ultrasonication. The resulting solids were recrystallised from a mixture of CH<sub>2</sub>Cl<sub>2</sub>-Et<sub>2</sub>O (1:1).

**3'-Benzoyl-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]dihydro-(3H)-furan-2-one (19a)**

Beige solid; yield 0.70 g (43%); mp 159-160 °C; IR (KBr) 1775, 1685 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ/TMS 2.40-2.75 (m, 2H), 2.85-3.10 (AB part of an ABX, 2H, *J*<sub>AX</sub>=4.2, *J*<sub>BX</sub>=7.5, *J*<sub>AB</sub>=12.4 Hz), 4.30-4.55 (m, 2H), 5.55 (X part of an ABX, 1H, *J*<sub>AX</sub>=4.2, *J*<sub>BX</sub>=7.5), 6.95-8.20 (m, 10H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ/TMS 34.4, 37.2, 65.9, 69.0, 82.5, 115.1-148.9, 174.6, 195.3; Anal. Calcd for C<sub>19</sub>H<sub>17</sub>NO<sub>4</sub>: C, 70.58; H, 5.30; N, 4.33. Found: C, 70.87; H, 5.41; N, 4.34.

**3'-(4-Methylbenzoyl)-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]dihydro-(3H)-furan-2-one (19b)**

Beige solid; yield 0.52 g (31%); mp 142-143 °C; IR (KBr) 1775, 1680 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ/TMS 2.40-2.75 (m, 2H), 2.45 (s, 3H), 2.80-3.10 (AB part of an ABX, 2H, *J*<sub>AX</sub>=4.4, *J*<sub>BX</sub>=7.5, *J*<sub>AB</sub>=12.5 Hz), 4.30-4.55 (m, 2H), 5.50 (X part of an ABX, 1H, *J*<sub>AX</sub>=4.4, *J*<sub>BX</sub>=7.5), 6.95-8.00 (m, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ/TMS 21.4, 34.0, 37.4, 65.7, 68.8, 82.3, 114.9-149.0, 174.6, 194.8; Anal. Calcd for C<sub>20</sub>H<sub>19</sub>NO<sub>4</sub>: C, 71.20; H, 5.68; N, 4.15. Found: C, 70.97; H, 5.76; N, 4.21.

**3'-(4-Methoxybenzoyl)-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]dihydro-(3H)-furan-2-one (19c)**

Colorless solid; yield 0.58 g (33%); mp 136-138 °C; IR (KBr) 1775, 1680 cm<sup>-1</sup>; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ/TMS 2.35-2.75 (m, 2H), 2.85-3.05 (AB part of an ABX, 2H, *J*<sub>AX</sub>=4.5, *J*<sub>BX</sub>=7.0, *J*<sub>AB</sub>=12.4 Hz), 3.95 (s, 3H), 4.30-4.55 (m, 2H), 5.45 (X part of an ABX, 1H, *J*<sub>AX</sub>=4.5, *J*<sub>BX</sub>=7.0), 6.90-8.15 (m, 9H); <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ/TMS 34.3, 37.2, 56.4, 65.9, 69.0, 82.6, 110.9-160.0, 174.7, 193.0; Anal. Calcd for C<sub>20</sub>H<sub>19</sub>NO<sub>5</sub>: C, 67.98; H, 5.42; N, 3.96. Found: C, 68.13; H, 5.51; N, 3.89.

**3'-(4-Nitrobenzoyl)-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]dihydro-(3H)-furan-2-one (19d)**

Yellow solid; yield 0.77 g (42%); mp 180-181 °C, IR (KBr) 1775, 1700 cm<sup>-1</sup>, <sup>1</sup>H NMR (*acetone-d*<sub>6</sub>) δ/TMS 2.40-2.80 (m, 2H), 2.80-3.20 (AB part of an ABX, 2H, *J*<sub>AX</sub>=7.1, *J*<sub>BX</sub>=2.9, *J*<sub>AB</sub>=12.7 Hz), 4.30-4.55 (m, 2H), 5.95 (X part of an ABX, 1H, *J*<sub>AX</sub>=7.1, *J*<sub>BX</sub>=2.9), 6.90-8.50 (m, 9H); <sup>13</sup>C NMR (*acetone-d*<sub>6</sub>) δ/TMS 35.1, 37.0, 66.6, 70.3, 84.2, 115.7-151.2, 175.4, 196.6; Anal. Calcd for C<sub>19</sub>H<sub>16</sub>N<sub>2</sub>O<sub>6</sub>: C, 61.95; H, 4.38; N, 7.61. Found: C, 62.14; H, 4.33; N, 7.67.

**3'-Benzoyl-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]succinic anhydride (20a)**

White solid; yield 1.55 g (92%); mp 135-136 °C; IR (KBr) 1875, 1795, 1680  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  2.95-3.20 (AB part of an ABX, 2H,  $J_{\text{AX}}=6.7$ ,  $J_{\text{BX}}=2.6$ ,  $J_{\text{AB}}=12.7$  Hz), 3.20-3.55 (AB, 2H,  $J_{\text{AB}}=19.6$  Hz), 5.65 (X part of an ABX, 1H,  $J_{\text{AX}}=6.7$ ,  $J_{\text{BX}}=2.6$ ), 7.00-8.10 (m, 10H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  37.6, 41.0, 68.1, 84.4, 115.4-147.7, 166.7, 170.7, 194.4; Anal. Calcd for  $\text{C}_{19}\text{H}_{15}\text{NO}_5$ : C, 67.65; H, 4.48; N, 4.15. Found: C, 67.93; H, 4.52; N, 4.13.

**3'-(4-Methylbenzoyl)-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]succinic anhydride (20b)**

White solid; yield 1.36 g (77%); mp 133-134 °C; IR (KBr) 1875, 1790, 1680  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  2.45 (s, 3H), 2.95-3.15 (AB part of an ABX, 2H,  $J_{\text{AX}}=6.5$ ,  $J_{\text{BX}}=2.7$ ,  $J_{\text{AB}}=12.8$  Hz), 3.15-3.55 (AB, 2H,  $J_{\text{AB}}=19.5$  Hz), 5.65 (X part of an ABX, 1H,  $J_{\text{AX}}=6.5$ ,  $J_{\text{BX}}=2.7$ ), 7.00-8.00 (m, 9H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  21.6, 37.8, 40.9, 68.1, 84.3, 115.4-147.8, 166.9, 170.9, 194.1; Anal. Calcd for  $\text{C}_{20}\text{H}_{17}\text{NO}_5$ : C, 68.37; H, 4.88; N, 3.99. Found: C, 68.59; H, 4.97; N, 4.03.

**3'-(4-Methoxybenzoyl)-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]succinic anhydride (20c)**

White solid; yield 1.45 g (79%); mp 132-133 °C; IR (KBr) 1880, 1805, 1680  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  2.95-3.20 (AB part of an ABX, 2H,  $J_{\text{AX}}=6.8$ ,  $J_{\text{BX}}=2.5$ ,  $J_{\text{AB}}=12.5$  Hz), 3.20-3.55 (AB, 2H,  $J_{\text{AB}}=19.5$  Hz), 3.90 (s, 3H), 5.60 (X part of an ABX, 1H,  $J_{\text{AX}}=6.8$ ,  $J_{\text{BX}}=2.5$ ), 6.80-8.10 (m, 9H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  37.8, 40.9, 55.4, 67.9, 84.3, 113.9-164.1, 166.9, 170.9, 192.8; Anal. Calcd for  $\text{C}_{20}\text{H}_{17}\text{NO}_6$ : C, 65.39; H, 4.66; N, 3.81. Found: C, 65.15; H, 4.81; N, 4.01.

**3'-(4-Nitrobenzoyl)-2'-phenyl-2',3'-dihydrospiroisoxazolino[5':3]succinic anhydride (20d)**

Yellow solid; yield 1.71 g (90%); mp 148-150 °C; IR (KBr) 1875, 1790, 1680  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  2.95-3.20 (AB part of an ABX, 2H,  $J_{\text{AX}}=6.7$ ,  $J_{\text{BX}}=1.6$ ,  $J_{\text{AB}}=12.8$  Hz), 3.15-3.55 (AB, 2H,  $J_{\text{AB}}=19.5$  Hz), 5.70 (X part of an ABX, 1H,  $J_{\text{AX}}=6.7$ ,  $J_{\text{BX}}=1.6$ ), 7.00-8.40 (m, 9H);  $^{13}\text{C}$  NMR ( $\text{CDCl}_3$ )  $\delta/\text{TMS}$  37.2, 41.0, 68.7, 84.7, 115.7-147.5, 166.2, 170.4, 193.6; Anal. Calcd for  $\text{C}_{19}\text{H}_{14}\text{N}_2\text{O}_7$ : C, 59.69; H, 3.69; N, 7.33. Found: C, 59.42; H, 3.81; N, 7.16.

*X-Ray crystal structure analyses of compounds (19d) and (20d)*

For both structures, data were measured from very small prismatic crystals with a Nonius Kappa-CCD area-detector diffractometer, using graphite monochromated Mo-K $\alpha$  radiation, according to the phi and omega scan method, up to  $\theta = 19^\circ$  for (19d) and  $22^\circ$  for (20d).<sup>40</sup> The structures were solved with program *SHELXS86*<sup>41</sup> and refined by full-matrix least-squares, based upon all unique  $F^2$  with program *SHELXL97*.<sup>42</sup> The hydrogen atoms were fitted at theoretical positions and treated as riding, assigned of an isotropic displacement parameter equivalent to 1.10 the one of the bonded atom. CCDC 664441 & 664442 contain the supplementary crystallographic data for (19d) and (20d) respectively. These data can be obtained free of charge from the Cambridge Crystallographic Data Center via [www.ccdc.cam.ac.uk/data\\_request/cif](http://www.ccdc.cam.ac.uk/data_request/cif).

## ACKNOWLEDGEMENTS

We thank Gregorio Crini (CERAC, University of Franche-Comté) for NMR measurements, E. Pousson (University of Bourgogne) for performing the elemental analyses, and Angèle Chiaroni (Laboratory of Crystallochemistry, Institute of Chemistry of Natural Products) for fruitful collaboration on X-ray studies. Sandrine Gerber (Laboratory of Glycochemistry and Asymmetric Synthesis, Institute of Chemical Science and Engineering) is also acknowledged for fruitful discussion. This work was supported by the French Ministry of Research.

## REFERENCES

1. H. M. R. Hoffmann and J. Rabe, *Angew. Chem., Int. Ed. Engl.*, 1985, **24**, 94.
2. L. M. Polo, C. M. Castro, M. C. Cruzado, C. J. G. Collino, F. D. Cuello-Carrion, D. R. Ciocca, O. S. Giordano, M. Ferrari, and L.A. Lopez, *Eur. J. Pharmacology*, 2007, **556**, 19.
3. W. Adam and V. O. Navas-algado, *J. Org. Chem.*, 1995, **60**, 578.
4. S. M. Kupchan, D. C. Fessler, M. A. Eakin, and T. J. Giacobbe, *Science*, 1970, **168**, 376.
5. D. Hwang, N. H. Fischer, B. C. Jang, H. Y. Tak, J. K. Kim, and W. Lee, *Biochem. Biophys. Res. Commun.*, 1996, **226**, 810.
6. *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*, ed. by A. Padwa and W. H. Pearson, John Wiley and Sons, 2002.
7. P. de March, M. el Arrad, M. Figueredo, and J. Font, *Tetrahedron*, 1998, **54**, 11613.
8. R. Grigg, V. Savic, and M. Thornton-Pett, *Tetrahedron*, 1997, **53**, 10633.
9. D. Alonso-Perarnau, P. de March, M. el Arrad, M. Figueredo, J. Font, and T. Parella, *Tetrahedron*, 1997, **53**, 14763.
10. P. Micuch, L. Fisera, V. Ondrus, and P. Ertl, *Molecules*, 1997, **2**, 57.
11. E. M. Peterson, K. Xu, K. D. Holland, A. C. McKeon, S. M. Rothman, J. A. Ferrendelli, and D. F. Covey, *J. Med. Chem.*, 1994, **37**, 275.
12. M. Cacciarini, F. M. Cordero, C. Faggi, and A. Goti, *Molecules*, 2000, **5**, 637.
13. C. Roussel, R. Fihi, K. Ciamala, J. Vebrel, T. Zair, and C. Riche, *Org. Biomol. Chem.*, 2003, **1**, 2689.
14. J. C. Daran, R. Fihi, C. Roussel, N. Laghrib, M. Azrour, K. Ciamala, and J. Vebrel, *Acta Cryst., Sect. E*, 2006, **62**, O329.
15. R. Fihi, K. Ciamala, J. Vebrel, and N. Rodier, *Bull. Soc. Chim. Belg.*, 1995, **104**, 55.
16. C. Roussel, K. Ciamala, P. Audebert, and J. Vebrel, *New J. Chem.*, 1999, **23**, 989.
17. C. Roussel, R. Fihi, K. Ciamala, P. Audebert, and J. Vebrel, *New J. Chem.*, 2000, **24**, 471.
18. C. Roussel, K. Ciamala, J. M. Melot, J. Vebrel, and C. Riche, *J. Chem. Res. (S)*, 2002, 449.
19. C. Roussel, K. Ciamala, J. Vebrel, M. Knorr, and M. M. Kubicki, *Heterocycles*, 2007, **71**, 1517.
20. S. Stverkova, Z. Zak, and J. Jonas, *Liebigs Ann.*, 1995, 477.

21. A. Quilico and P. Grünanger, *Gazz. Chim. Ital.*, 1952, **82**, 140.
22. S. Rigolet, P. Goncalo, J. M. Melot, and J. Vebrel, *J. Chem. Res.(S)*, 1998, 686.
23. S. Rigolet, J. M. Melot, J. Vebrel, A. Chiaroni, and C. Riche, *J. Chem. Soc., Perkin Trans. 1*, 2000, [1095](#).
24. R. Huisgen and W. Mack, *Tetrahedron Lett.*, 1961, [583](#).
25. F. Desarlo, *J. Chem. Soc., Perkin Trans. 1*, 1974, [1951](#).
26. S. W. Baldwin and A. Long, *Org. Lett.*, 2004, **6**, [1653](#).
27. O. Tamura, T. Shiro, M. Ogasawara, A. Toyao, and H. Ishibashi, *J. Org. Chem.*, 2005, **70**, [4569](#).
28. W. B. Jennings, D. R. Boyd, and L. C. Waring, *J. Chem. Soc., Perkin Trans. 2*, 1976, [610](#).
29. M. Joucla, D. Gree, and J. Hamelin, *Tetrahedron*, 1973, **29**, [2315](#).
30. S. Chandrasekhar, M. Ravindranath, B. S. Neela, S. Ramakumar, and M. A. Viswamitra, *J. Chem. Res. (S)*, 1989, 252.
31. R. B. Woodward and R. Hoffmann, *The Conservation of Orbital Symmetry*, Verlag Chemie GmbH, 1971.
32. A. L. Spek, *J. Appl. Cryst.*, 2003, **36**, [7](#).
33. M. Ueda, M. Takahashi, T. Suzuki, Y. Imai, and C. U. Pittman, *J. Polym. Science Part a-Polymer Chemistry*, 1983, **21**, [1139](#).
34. C. R. Hutchinson, *J. Org. Chem.*, 1974, **39**, [1854](#).
35. P. A. Grieco and C. S. Pogonowski, *J. Org. Chem.*, 1974, **39**, [1958](#).
36. K. C. Liu, B. R. Shelton, and R. K. Howe, *J. Org. Chem.*, 1980, **45**, [3916](#).
37. C. Grundmann and R. Richter, *J. Org. Chem.*, 1967, **32**, [2308](#).
38. F. Kröhnke and E. Börner, *Chem. Ber.*, 1936, **69**, [2006](#).
39. A. R. Katritzky and C. W. Rees, *Comprehensive Heterocyclic Chemistry : the structure, reaction, synthesis and uses of heterocyclic compounds*; Oxford a.o. : Pergamon Press, 1984 ed., 1984; Vol. 4.
40. *Collect (Brucker AXS BV) 1997-2000; HKL Denzo and Scalepack (Otinowski & Minor, 1997)*.
41. G. M. Sheldrick, *Acta Cryst., Sect. A*, 1990, **46**, [467](#).
42. G. M. Sheldrick, *SHELXL97. Program for the refinement of Crystal structures, Univ. of Göttingen, Germany*, 1997.