

HETEROCYCLES, Vol. 81, No. 11, 2010, pp. 2465 - 2470. © The Japan Institute of Heterocyclic Chemistry
Received, 17th August, 2010, Accepted, 21st September, 2010, Published online, 22nd September, 2010
DOI: 10.3987/COM-10-12046

DESIGN AND SYNTHESIS OF NOVEL RETINOID SYNERGISTS HAVING A DIBENZODIAZEPINE SKELETON

Kiminori Ohta,^a Emiko Kawachi,^b Koichi Shudo,^c and Hiroyuki Kagechika^{b,*}

^aFaculty of Pharmaceutical Sciences, Tohoku Pharmaceutical University, 4-4-1 Komatsushima, Aoba-ku, Sendai 981-8558, Japan, ^bGraduate School of Biomedical Science, Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, 2-3-10 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-0062, Japan, ^cResearch Foundation Itsuu Laboratory, 2-28-10 Tamagawa, Setagaya-ku, Tokyo 158-0094, Japan; e-mail: kage.omc@tmd.ac.jp

Abstract – Based on the structures of potent RXR agonists **2** and **3**, novel dibenzodiazepine derivatives **4** – **6**, containing two diphenylamine substructures, were designed as RXR modulator candidates and synthesized by utilizing Pd-catalyzed and Cu-promoted diphenylamine-generating reactions as key reactions. These compounds showed retinoid-synergistic activity, enhancing the HL-60 cell differentiation-inducing ability of the RAR agonist Am80.

Retinoids have a broad spectrum of biological activities related to cellular differentiation and proliferation, and are essential for normal embryonic development in vertebrates.¹ Their biological responses are mediated by binding to and activation of retinoic acid receptors (RARs),² which act in the form of heterodimers with another class of retinoid nuclear receptors, retinoid X receptors (RXRs).³ All-*trans* retinoic acid (ATRA) binds to RARs, and its 9-*cis* isomer (9-*cis*RA) binds to both RARs and RXRs (Figure 1).⁴ Various synthetic retinoids, such as Am80 (**1**), bind only to RARs with an affinity that correlates well with most retinoidal activities.⁵ RXRs are transcriptionally silent partners of RARs, and RAR-RXR heterodimers activated by RAR ligands regulate the expression of specific genes.³ We have developed several RXR-specific agonists and reported their retinoid synergistic activities.⁶ For example, RXR agonists, such as HX600 (**2**)⁷ and DA023 (**3**),⁸ themselves exhibited no retinoidal activity, but strongly enhanced the potency of ATRA or Am80 (**1**). Since RXRs form heterodimers with various nuclear receptors, such as vitamin D₃ receptor, thyroid hormone receptors and peroxisome proliferator-activated receptors, RXR ligands may modulate the behaviors of the partner receptors, as well

as retinoidal activities.³ Therefore, we have focused on the development of RXR agonists and designed novel dibenzodiazepine-based RXR modulator candidates **4** – **6** by combining the structures of **2** and **3** (Figure 1). In this paper, we describe the synthesis and biological activities of the novel dibenzodiazepine derivatives **4** – **6**.






Figure 1. (a) Structures of endogenous and synthetic agonists for RARs and RXRs; (b) Design of novel dibenzodiazepine-based RXR agonists

The synthesis of compounds **4** and **5** is summarized in Scheme 1. Compound **8**, which was prepared from 1,2,3,4-tetrahydro-1,1,4,4-tetramethylnaphthalene (**7**) according to the literature procedure,⁹ was reacted with ethyl 4-iodobenzoate in the presence of Pd catalyst to afford diphenylamine **9**.¹⁰ An *o*-bromobenzyl group was introduced at the nitrogen atom of the diphenylamine skeleton of **9** with the corresponding benzyl bromide, followed by reduction of the nitro group with Zn in AcOH to afford compound **10**. Construction of a dibenzodiazepine skeleton was accomplished by means of an intramolecular Pd-catalyzed amination between the aniline and bromobenzene moieties of **10** to afford compound **11** in quantitative yield.¹⁰ Hydrolysis of the ethyl ester group with aqueous 20% KOH solution afforded compound **4**. *N*-Methylation by using NaH as a base and MeI, followed by hydrolysis of the ester group, afforded *N*-methyl dibenzodiazepine derivative **5** in quantitative yield. Dibenzodiazepinone **6** was synthesized as shown in Scheme 2. Compound **8** was reacted with methyl 2-chlorobenzoate in the presence of excess amount of CuI (Ullmann coupling) to afford diphenylamine derivative **12** in 65% yield.¹¹ *N*-Methylation by using NaH as a base and MeI, followed by catalytic hydrogenation of the nitro group with Pd/C afforded aniline derivative **13**. Compound **14** containing two diphenylamine substructures was prepared from **13** by means of Buchwald-Hartwig amination with ethyl 4-iodobenzoate

using Pd-catalyst.¹⁰ Lactamization reaction between the diphenylamine and methyl ester moieties of **14** was performed in the presence of excess NaH, followed by hydrolysis of the ethyl ester group with NaOH, which was prepared in situ from excess NaH and H₂O added to the reaction mixture, afforded compound **6**.



Scheme 1. Synthesis of dibenzodiazepine derivatives **4** and **5**; Reagents and conditions: (a) c.H₂SO₄, HNO₃, AcOH, rt, 44%; (b) Pd/C, H₂, EtOH, rt, 94%; (c) c.H₂SO₄, HNO₃, TFA, rt, 38%; (d) Pd₂(dba)₃, *rac*-BINAP, NaO^tBu, ethyl 4-iodobenzoate, toluene, reflux, 34%; (e) NaH, *o*-bromobenzyl bromide, DMF, rt, 98%; (f) Zn, AcOH, rt, 43%; (g) Pd₂(dba)₃, *rac*-BINAP, NaO^tBu, toluene, reflux, 100%; (h) 20% KOH aq., EtOH, rt, quant; (i) NaH, MeI, DMF, rt, quant.



Scheme 2. Synthesis of compound **6**; Reagents and conditions: (a) K₂CO₃, CuI, methyl 2-chlorobenzoate, xylene, reflux, 65%; (b) NaH, CH₃I, DMF, rt; (c) Pd/C, H₂, rt, 2 steps: 75%; (d) Pd₂(dba)₃, *rac*-BINAP, NaO^tBu, ethyl 4-iodobenzoate, toluene, reflux, 21%; (e) NaH (excess), xylene, reflux, then H₂O, rt, 2steps: 56%.

The biological activities of the synthesized compounds were evaluated in terms of the activity to induce differentiation of HL-60 cells into mature granulocytes in the presence or absence of a synthetic retinoid, Am80 (**1**).¹² The differentiated cells were identified by nitro blue tetrazolium (NBT) reduction assay.¹² None of the test compounds induced cell differentiation of HL-60 alone (data not shown), while all of them showed moderate synergistic retinoid activity with Am80 (Figure 2). Their potency is weaker than that of **2**. Compound **5**, bearing a *N*-methyl group on the diazepine ring, showed more potent synergistic activity than compound **4** without the *N*-methyl group. Introduction of a carbonyl group into the diazepine skeleton led to a decrease of the activity; i.e. the synergistic activity of compound **6** was weaker than that of **5**. These results suggest that polar functional groups on the diazepine skeleton are unfavorable for the expression of retinoid synergistic activity. Further syntheses and structure-activity relationship studies, particularly for compounds with an oxygen and a sulfur atom instead of the *N*-methyl group of compound **5**, are in progress.




Figure 2. Synergistic effects of dibenzodiazepine derivatives **4** – **6** on HL-60 cell differentiation induced by 0.1 nM Am80 (**1**)

In conclusion, newly designed benzodiazepine derivatives **4** – **6** were efficiently synthesized by utilizing Pd-catalyzed and Cu-promoted diphenylamine-generating reactions as key reactions. Compounds **4** – **6** acted as retinoid synergists, and enhanced the cell differentiation induced by 0.1 nM Am80 (**1**). Although the potency of these compounds is not great, this efficient synthetic method for novel dibenzodiazepine

structure bearing two diphenylamine substructures should be applicable elsewhere in the medicinal-chemical study of retinoids and other nuclear receptor ligands.

REFERENCES

1. M. B. Sporn, A. B. Roberts, and D. S. Goodman, 'The Retinoids: Biology, Chemistry, and Medicine', 2nd ed., Raven Press, New York, 1994; M. Mark, N. B. Ghyselinck, and P. Chambon, *Ann. Rev. Pharmacol. Toxicol.*, 2006, **46**, 451; M. Mark, N. B. Ghyselinck, and P. Chambon, *Nuc. Rec. Signal.*, 2009, **7**, 1.
2. D. J. Mangelsdorf, C. Thummel, M. Beato, P. Herrlich, G. Schuetz, K. Umesono, B. Blumberg, P. Kastner, M. Mark, P. Chambon, and R. M. Evans, *Cell*, 1995, **83**, 835; P. Kastner, M. Mark, N. Ghyselinck, W. Krezel, V. Dupe, J. M. Grondona, and P. Chambon, *Development*, 1997, **124**, 313; C. K. Glass and M. G. Rosenfeld, *Gene Dev.*, 2000, **14**, 121.
3. P. Chambon, *FASEB J.*, 1996, **10**, 940; G. R. Benoit, M. Flexor, F. Besançon, L. Altucci, A. Rossin, J. Hillion, Z. Barajthy, L. Legres, E. Ségal-Bendirdjian, H. Gronemeyer, and M. Lanotte, *Mol. Endocrinol.*, 2001, **15**, 1154.
4. P. Germain, P. Chambon, G. Eichele, R. M. Evans, M. A. Lazar, M. Leid, A. R. De Lera, R. Lotan, D. J. Mangelsdorf, and H. Gronemeyer, *Pharmacol. Rev.*, 2006, **58**, 712; P. Germain, P. Chambon, G. Eichele, R. M. Evans, M. A. Lazar, M. Leid, A. R. De Lera, R. Lotan, D. J. Mangelsdorf, and H. Gronemeyer, *Pharmacol. Rev.*, 2006, **58**, 760.
5. H. Kagechika, E. Kawachi, Y. Hashimoto, T. Himi, and K. Shudo, *J. Med. Chem.*, 1988, **31**, 2182; Y. Hashimoto, H. Kagechika, and K. Shudo, *Biochem. Biophys. Res. Commun.*, 1990, **166**, 1300; K. Shudo, H. Kagechika, N. Yamazaki, M. Igarashi, and C. Tateda, *Biol. Pharm. Bull.*, 2004, **27**, 1887; M. Ishido and H. Kagechika, *Drugs Today*, 2007, **43**, 563.
6. H. Kagechika and K. Shudo, *J. Med. Chem.*, 2005, **48**, 5875.
7. H. Umemiya, H. Fukasawa, M. Ebisawa, L. Eyrolles, E. Kawachi, G. Eisenmann, H. Gronemeyer, Y. Hashimoto, K. Shudo, and H. Kagechika, *J. Med. Chem.*, 1997, **40**, 4222; H. Umemiya, H. Kagechika, H. Fukasawa, E. Kawachi, M. Ebisawa, Y. Hashimoto, G. Eisenmann, C. Erb, A. Pornon, P. Chambon, H. Gronemeyer, and K. Shudo, *Biochem. Biophys. Res. Commun.*, 1997, **233**, 121.
8. K. Ohta, M. Tsuji, E. Kawachi, H. Fukasawa, Y. Hashimoto, K. Shudo, and H. Kagechika, *Biol. Pharm. Bull.*, 1998, **21**, 544.
9. L. Eyrolles, H. Kagechika, E. Kawachi, H. Fukasawa, T. Iijima, Y. Matsushima, Y. Hashimoto, and K. Shudo, *J. Med. Chem.*, 1994, **37**, 1508.
10. J. P. Wolfe, S. Wagaw, J.-F. Marcoux, and S. L. Buchwald, *Acc. Chem. Res.*, 1998, **31**, 805; J. F. Hartwig, *Angew. Chem. Int. Ed.*, 1998, **37**, 2046.

11. J. Lindley, [Tetrahedron, 1984, 40, 1433](#); J. Hassan, M. Sevignon, C. Gozzi, E. Schulz, and M. Lemaire, [Chem. Rev., 2002, 102, 1359](#).
12. K. Ohta, E. Kawachi, N. Inoue, H. Fukasawa, Y. Hashimoto, A. Itai, and H. Kagechika, *Chem. Pharm. Bull.*, 2000, **48**, 1504.