

HETEROCYCLES, Vol. 88, No. 2, 2014, pp. 1625 - 1632. © 2014 The Japan Institute of Heterocyclic Chemistry
Received, 30th August, 2013, Accepted, 31st October, 2013, Published online, 8th November, 2013
DOI: 10.3987/COM-13-S(S)115


ACID-PROMOTED AROMATIZATION OF PERYLENE-BASED ENDOXIDES

Alejandro Criado, Manuel Vilas-Varela, Agustín Cobas, Dolores Pérez, Diego Peña, and Enrique Guitián*

Centro de Investigación en Química Biológica e Materiais Moleculares (CIQUS) and Departamento de Química Orgánica, Universidad de Santiago de Compostela, 15782-Santiago de Compostela, Spain; E-mail: enrique.guitian@usc.es

Abstract – The aromatization of perylene-based endoxides by treatment with HCl and EtOH has been studied.

We have recently reported the synthesis of benzo[*a*]perylene (**4**) as shown in Scheme 1.¹ The reaction of **1** with benzyne (**2**) triggers a tandem domino Diels-Alder sequence to afford adduct **3** in excellent yield and high stereoselectivity, which by treatment with HCl/EtOH afforded **4** in 72% yield. Following the same protocol we have prepared several analogues of **3** and the corresponding perylene-derivatives.




Scheme 1. Synthesis of benzo[*a*]perylene (**4**)

In that publication we focused our attention on the synthetic strategy and on the properties of perylene-derivatives, without a detailed mechanistic discussion. Only some details of the computational study of the transformation of **1** into endoxide **3** were included. However the transformation of endoxide **3** into perylene **4** is far from trivial. We will discuss here some mechanistic aspects of this reaction, but


before treating the above transformation we will outline the more reasonable mechanisms for the acid-promoted aromatization of 1,4-endoxides.

Acid-promoted aromatization of cyclohexene-1,4-endoxides such as **I** is usually an easy process.²⁻⁹ It involves protonation to **II** and a formal elimination of a proton and water to afford the aromatic compound **III**. The situation for cyclohexadiene-1,4-endoxides **IV** is more complex. After the opening of the endoxide to the carbenium ion **V**, there is no hydrogen available to accomplish the elimination. A formal reduction of **V** to **VI** must occur to allow dehydration to **III** (Scheme 2). One-pot procedures with combinations of an acid and a reducing agent, such as HOAc/Zn,^{10,11} HOAc/Fe,¹¹ TFA/NaBH₄,¹² TiCl₄/LiAlH₄,¹³⁻¹⁵ and TiCl₄/Zn,¹⁶ or reagents such as Fe₂(CO)₉^{17,18} or TMSI¹⁹ have been used to perform this transformation. Alternatively, cyclohexadiene-1,4-endoxides **IV** have been transformed into arenes **III** in a two-operation process, involving a first reduction step to give **I**, followed by protonation and elimination of water.²⁰⁻²²




Scheme 2. Typical mechanisms for the acid-promoted aromatization of endoxides

According to the above considerations acid-promoted opening of **3** could afford carbenium ions **5** and **6**, which could lose water to afford endoxides **7** and **8**, respectively. From these, a second protonation/opening sequence would afford carbenium ions **9** and **10**, respectively (Scheme 3). DFT calculations at the B3LYP/6-31G(d) level show that the formation of **9** is thermodynamically favoured by 0.011711 Hartrees (7.3 kcal/mol). In any case, transformation of these compounds into benzo[*a*]perylene (**4**) requires a formal reduction and the elimination of water.



Scheme 3. Plausible intermediates in the opening of endoxide **3** (energies in Hartrees)


In our case, having used only HCl and EtOH as reagents for the transformation of **3** into **4**, we thought that EtOH should be the actual reducing agent. In fact, though it is not usually considered as a reducing agent, EtOH acts in this way against strong oxidants and also in biological transformations, such as the reduction of NAD^+ in liver, catalyzed by alcohol dehydrogenase.²³ Closer to our transformation, the reduction of triphenylmethyl cation (**11**) by hydride transfer from ethanol has been reported in the literature (Scheme 4).²⁴⁻²⁶



Scheme 4. Alternative mechanisms for hydride transfer to the triphenylmethyl cation

The study of these reactions using HCl/CH₃CD₂OH had shown that the incorporation of deuterium occurs exclusively at the central atom to give **12**. Two alternative mechanisms have been considered: a direct nucleophilic attack of hydride to the carbocation **11**^{24,25} and the addition of ethanol to an ortho-position to give **13**, followed by an intramolecular hydride transfer to **12** (Scheme 4).²⁶


Extrapolation of this mechanisms to carbenium ions **9** and **10**, led us to predict that the incorporation of deuterium should lead to intermediates **14** and **5**, respectively, which after the elimination of DHO should afford non-deuterated benzo[*a*]pyrene **4** (Scheme 5). In fact, when we treated endoxide **3** with HCl and CH₃CD₂OH we isolated non-deuterated **4**, as proved by NMR analysis.




Scheme 5. Addition and elimination of deuterium

The transformation of endoxide **3** into benzo[*a*]perylene (**4**) has also been achieved using HCl/MeOH or HCl/*n*-BuOH, but in lower yield.

Finally, we tried the aromatization of compound **16** obtained by a similar tandem cycloaddition. However, when we treated **16** with HCl/EtOH as above a new compound was obtained having a CO band at 1653 cm⁻¹ in its IR spectrum and two singlets at 7.26 (1H) and 6.13 (1H) ppm in its ¹H-NMR spectrum. This led us to proposed structure **17** for that compound, which was confirmed by X-ray diffraction studies, as shown in Figure 1.

Figure 1. X-Ray diffraction structure of ketone **17**

We firstly considered that the formation of **17** could involve an initial protonation of **16** followed by opening to carbenium ion **18**,²⁷ deprotonation to **19** and retro-ene fragmentation to **20** (Scheme 6). Aromatization of the endoxide **20** with HCl/EtOH as above would afford **17**.

Scheme 6. Mechanistic proposals for the formation of **17**

However, an alternative mechanism for the transformation of **18** into **20**, involving a hydride migration to give **21** followed by a fragmentation to give the final ketone, could not be discarded.²⁸ To evaluate the feasibility of these mechanisms we carried out the reaction with DCl/CD₃OD, assuming that if the first mechanism were in operation the alcohol **19** would undergo H/D exchange with the medium and the product **20** would incorporate deuterium. On the contrary, if **20** were formed by 1,2-migration of H, deuterium should not be incorporated. After doing the experiment, non-deuterated **17** was isolated, supporting the second mechanistic hypothesis.

In summary, calculations and experimental data for the aromatization of endoxide **3** with HCl/EtOH suggest a mechanism involving the formation of intermediary carbenium ions, its reduction and the elimination of water. A mechanistic proposal involving a hydride donation from the solvent has been discussed.

EXPERIMENTAL

Benzo[*a*]perylene (4).¹ Concentrated HCl solution (50 μ L, 37.5% in water) was added to a solution of adduct **3** (50 mg, 0.15 mmol) in EtOH, MeOH or *n*-BuOH (3 mL), and the mixture was stirred under argon in the absence of light at 78 °C for 5 h. Then, H₂O (1 mL) was added and the mixture was concentrated under reduced pressure. The residue was purified by column chromatography (under argon in the absence of light; SiO₂; 9:1 hexane/CH₂Cl₂), affording benzo[*a*]perylene derivatives (**4**). Yield: 72% in EtOH, 32% in MeOH, 31% in *n*-BuOH.

***p*-Tolyl(2-(9-(*p*-tolyl)-7*H*-benzo[*de*]anthracen-7-yl)phenyl)metanone (17).** Concentrated HCl solution (140 μ L, 0.16 mmol, 37.5% in water) was added to a solution of endoxide **16** (42 mg, 0.08 mmol) in EtOH (5 mL), and the mixture was stirred under argon in the absence of light at 78 °C for 4 h. Then, H₂O (1 mL) was added and the mixture was concentrated under reduced pressure. The residue was purified by column chromatography (under argon in the absence of light; SiO₂; 9:1 hexane/CH₂Cl₂), affording ketone **17** (30 mg, 75% yield) as a green solid, mp 213-216 °C; IR 1653 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 8.13 (d, *J* = 7.9 Hz, 2H), 7.93 (*J* = 8.1 Hz, 2H), 7.77 (*J* = 8.2 Hz, 1H), 7.67 (*J* = 7.7 Hz, 1H), 7.58-7.50 (m, 2H), 7.48 (d, *J* = 7.2 Hz, 1H), 7.45-7.42 (m, 1H), 7.41-7.29 (m, 5H), 7.262 (s, 1H), 7.17 (d, *J* = 8.0 Hz, 2H), 7.15-7.10 (m, 2H), 6.99-6.96 (m, 1H), 6.13 (s, 1H), 2.50 (s, 3H), 2.38 (s, 3H) ppm; ¹³C NMR (100 MHz, CDCl₃), δ 198.2 (C), 147.0 (C), 144.5 (C), 140.1 (C), 139.0 (C), 137.3 (C), 137.1 (C), 137.03 (C), 137.00 (C), 135.4 (C), 133.7 (C), 130.8 (2CH), 130.7 (CH), 130.6 (CH), 130.3 (C), 130.0 (C), 129.4 (2CH), 129.3 (2CH), 128.7 (CH), 127.8 (CH), 127.7 (CH), 127.6 (C), 127.1 (CH), 126.5 (2CH), 126.3 (CH), 126.0 (CH), 125.9 (CH), 125.2 (CH), 124.9 (CH), 123.7 (CH), 118.9 (CH), 44.9 (CH), 21.6 (CH₃), 20.9 (CH₃) ppm. LRMS (EI), *m/z* (%): 500 (63, M⁺), 408 (100), 379 (48); HRMS (EI) calculated for

C₃₈H₂₈O: 500.2140, found: 500.2139; X-Ray Diffraction Studies: Deposition number CCDC-942555 for compound **17**. Free copies of the data can be obtained via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK; Fax: +44 1223 336033; e-mail: deposit@ccdc.cam.ac.uk).

ACKNOWLEDGEMENTS

Financial support from the Spanish Ministry of Economy and Competitiveness (MINECO, CTQ2010-18202, CTQ2010-18208), *Xunta de Galicia* (10PXIB2200222PR) and FEDER is gratefully acknowledged. Generous allocation of computer time was kindly provided by the Centro de Supercomputación de Galicia (CESGA). A. C. thanks the MEC for the award of a FPU fellowship. We thank Benjamin T. King for helpful discussions.

REFERENCES AND NOTES

1. A. Criado, D. Peña, A. Cobas, and E. Guitián, *Chem. Eur. J.*, 2010, **16**, 9736.
2. M. P. Cava and F. M. Scheel, *J. Org. Chem.*, 1967, **32**, 1304.
3. J. Luo and H. Hart, *J. Org. Chem.*, 1987, **52**, 3631.
4. T. Ghosh and H. Hart, *J. Org. Chem.*, 1989, **54**, 5073.
5. S. W. Thomas, III, T. M. Long, B. D. Pate, S. R. Kline, E. L. Thomas, and T. M. Swager, *J. Am. Chem. Soc.*, 2005, **127**, 17976.
6. J. Rybáček, J. Závada, and P. Holy, *Synthesis*, 2008, 3615.
7. H. Qu and C. Chi, *Org. Lett.*, 2010, **12**, 3360.
8. J. E. Rainbolt and G. P. Miller, *J. Org. Chem.*, 2007, **72**, 3020.
9. B.-J. Pei, W.-H. Chan, and A. W. M. Lee, *J. Org. Chem.*, 2010, **75**, 7332.
10. G. Wittig, E. Knauss, and K. Neithammer, *Justus Liebigs Ann. Chem.*, 1960, **630**, 10.
11. D. Chun, Y. Cheng, and F. Wudl, *Angew. Chem. Int. Ed.*, 2008, **47**, 8380.
12. G. W. Gribble, W. J. Kelly, and M. P. Sibi, *Synthesis*, 1982, 143.
13. C. H. Wong, C. W. Hung, and H. N. C. Wong, *J. Organomet. Chem.*, 1988, **342**, 9.
14. B.-J. Pei, W.-H. Chan, and A. W. M. Lee, *Org. Lett.*, 2011, **13**, 1774.
15. Y.-L. Chen, C.-K. Hau, H. Wang, H. He, M.-S. Wong, and A. W. M. Lee, *J. Org. Chem.*, 2006, **71**, 3512.
16. D. K. P. Ng, Y.-O. Yeung, W. K. Chan, and S.-C. Yu, *Tetrahedron Lett.*, 1997, **38**, 6701.
17. W. M. Best, P. A. Collins, R. K. McCulloch, and D. Wege, *Aust. J. Chem.*, 1982, **35**, 843.
18. J. Basset, M. Romero, T. Serra, and M. D. Pujol, *Tetrahedron*, 2012, **68**, 356.
19. J. G. Smith, S. S. Welankiwar, B. S. Shantz, E. H. Lai, and N. G. Chu, *J. Org. Chem.*, 1980, **45**, 1817.

20. G. Wittig and L. Pohmer, [*Chem. Ber.*, 1956, **89**, 1334.](#)
21. M. S. Newman and J. A. Cella, [*J. Org. Chem.*, 1973, **38**, 3482.](#)
22. H. Hart and D. Ok, [*J. Org. Chem.*, 1986, **51**, 976.](#)
23. H. Theorell and J. S. M. McKee, [*Nature*, 1961, **192**, 47.](#)
24. P. D. Barlett and J. D. McCollum, [*J. Am. Chem. Soc.*, 1956, **78**, 1441.](#)
25. G. A. Olah and J. J. Svoboda, [*J. Am. Chem. Soc.*, 1973, **95**, 3794.](#)
26. P. Huszthy, K. Lempert, G. Simig, and J. Tamás, [*J. Chem. Soc., Perkin Trans. 2*, 1982, 1671.](#)
27. DFT calculations at the B3LYP/6-31G(d) level shown that **18** (-1615.71133 au) is thermodynamically more stable than the cation formed by protonation on the left endoxide (-1615.70731 au).
28. This mechanistic proposal has been suggested by a referee.