

CYTOTOXIC XANTHONES FROM *HYPERICUM CHINENSE*

Yinke Li,^{1,3} Yong Xu,² Jia Wu,² Sujuan Wang,² Yanlin Meng,¹ Yanqing Ye,¹
Haiyin Yang,¹ Xuemei Gao,^{1,*} and Qiufen Hu^{1,*}

¹ Key Laboratory of Chemistry in Ethnic Medicinal Resources, State Ethnic Affairs Commission & Ministry of Education, Yunnan University of Nationalities, Kunming 650031, P.R. China; ² Key Laboratory of Tobacco Chemistry of Yunnan Province, Yunnan Academy of Tobacco Science, Kunming 650106, P.R. China; ³ Collge of Resource and Environment, Yuxi Normal University, Yuxi 653100, P. R. China. E-mail: huqiufena@yahoo.com.cn; gao_xuemei@hotmail.com

Abstract – Two new xanthenes, 1-hydroxy-4-(2-hydroxyethyl)-3,8-dimethoxy-9*H*-xanthen-9-one (**1**) and (*S*)-1-hydroxy-4-(1-hydroxy-3-oxobutyl)-3,8-dimethoxy-9*H*-xanthen-9-one (**2**), together with three known xanthenes (**3-5**) were isolated from the leaves and stems of *Hypericum chinense*. Their structures were elucidated by spectroscopic methods, including extensive 1D- and 2D NMR techniques. Compounds **1-5** were tested for their cytotoxicity against five human tumor cell lines (NB4, A549, SHSY5Y, PC3, and MCF7). The results revealed that compound **1** showed high cytotoxicity against A549 and PC3 cell with IC₅₀ values of 2.8 and 3.4 μM, and **2** showed high cytotoxicity against PC3 cell with IC₅₀ valve of 2.8 μM, respectively.

The family Clusiaceae is a rich source of xanthenes.^{1,2} Xanthenes are typically polysubstituted and occur as either fully aromatized, dihydro-, tetrahydro-, or, more rarely, hexahydro- derivatives.² This family of compounds appeals to medicinal chemists because of their pronounced biological activity within a notably broad spectrum of disease states, including anti-hepatitis B virus,³ anti-tobacco mosaic virus,⁴ antibacterial,^{5,6} antioxidant,^{7,8} anti-inflammatory,⁹ tumor-promoting inhibition,¹⁰ cytotoxicity,^{11,12} and the like, as a result of their interaction with a correspondingly diverse range of target biomolecules.

The genus *Hypericum* belonging to Clusiaceae is distributed widely in temperate regions, and has been used for traditional medicines in various parts of the world. In China, *Hypericum. chinese* is used as a folk medicine for treatment of female disorders.¹³ Previous phytochemical investigations on *H. chinese*

resulted in the isolation of xanthenes,¹² acylphloroglucinols,¹⁴ lactones,¹⁵ and norlignans.¹⁶ With the aim of multipurpose utilization of herb plants and identify bioactive natural products from this genus, the phytochemical investigation on *H. chinese* was carried out. As a result, two new xanthenes (**1** and **2**), together with three known xanthenes (**3-5**), were isolated. Compound **2** is the first naturally occurring xanthone possessing a 1-hydroxy-3-oxobutyl moiety. The structures of new compounds were elucidated by comprehensive analysis of their NMR data. In addition, the cytotoxicity of **1-5** were evaluated. The details of the isolation, structure elucidation, and cytotoxicity of the isolates are reported in this article.




Figure 1. The structures of xanthenes from *H. chinese*

A 70% aq. acetone extract prepared from the leaves and stems of *H. chinese* was subjected repeatedly to column chromatography on Silica gel, Sephadex LH-20, RP-18 and Preparative HPLC to afford compounds **1-5**, including two new xanthenes, named 1-hydroxy-4-(2-hydroxyethyl)-3,8-dimethoxy-9H-xanthen-9-one (**1**) and (*S*)-1-hydroxy-4-(1-hydroxy-3-oxobutyl)-3,8-dimethoxy-9H-xanthen-9-one (**2**), together with three known xanthenes, 1,3,7-trihydroxy-2-(2-hydroxy-3-methyl-3-butenyl)xanthone (**3**),¹⁷ 1,7-dihydroxy-2,3-[2''-(1-hydroxy-1-methylethyl)dihydrofurano]xanthone (**4**),¹⁷ cratoxylumxanthone D (**5**).¹⁸ The structures of the compounds **1-5** were shown in Figure 1 and the ¹H and ¹³C NMR data of **1** and **2** were listed in Table 1.

Compound **1** was isolated as a yellow gum. The HRESIMS of **1** gave the pseudomolecular [M+Na]⁺ ion at *m/z* 339.0840, corresponding to a molecular formula of C₁₇H₁₆O₆. Its UV spectrum showed the maximum absorption at 302, 245, and 210 nm. Strong absorption bands accounting for hydroxy (3432 cm⁻¹), carbonyl (1658 cm⁻¹), and aromatic groups (1602, 1546, 1468 cm⁻¹) could also be observed in its IR spectrum. The ¹H- and ¹³C NMR spectrum (Table 1) displayed signals for all 17 carbons and 16 protons, including a xanthenes skeleton¹⁹ (C-1 ~ C-9, C-4a, C-8a ~ C-10a; H-2, H-5 ~ H-7), two

methoxy groups (δ_C 56.0 q, 56.2 q; δ_H 3.85 s, 3.81 s), a hydroxyethyl unit¹¹ [δ_C 34.0 t, 63.8 t; δ_H 2.54 t (7.2), 3.67 t (7.2)], and a phenolic hydroxy group (δ_H 13.43 s). The typical proton signals of ring A [δ_H 6.85 d (8.3), 7.42 t (8.3), 6.70 d (8.3)] and ring B (δ_H 6.59 s) suggested that **1** should be a 1,3,4,8-tetrasubstituted xanthone.¹⁹ The




Figure 2. Key HMBC (\curvearrowright) correlations of **1**

HMBC correlation (Figure 2) of one methoxy proton signal (δ_H 3.85) with C-3 (δ_C 161.0) showed this methoxy group was located at C-3. The correlation between the proton signal (δ_H 3.81) and C-8 (δ_C 165.4) indicated the other methoxy group located at C-8. The long-range correlations of H₂-1' (δ_H 2.54) to C-3 (δ_C 161.0), C-4 (δ_C 108.8) and C-4a (δ_C 155.3), of H₂-2' (δ_H 3.67) to C-4 (δ_C 108.8) were observed in **1**. This led us to conclude that the hydroxyethyl unit was located at C-4. Finally, the HMBC correlations between the phenolic hydroxy proton (δ_H 13.43) and C-1 (δ_C 162.8), C-2 (δ_C 98.1), and C-9a (δ_C 104.1) led to the assignment of the phenolic hydroxy group at C-1. Therefore, compound **1** was assigned as 1-hydroxy-4-(2-hydroxyethyl)-3,8-dimethoxy-9H-xanthen-9-one.

Table 1. ¹H and ¹³C NMR Data of Compounds **1** and **2** (δ in ppm, 500 and 125 MHz)

No.	Compound 1 ^a		Compound 2 ^b	
	δ_C (m)	δ_H (m, J, Hz)	δ_C (m)	δ_H (m, J, Hz)
1	162.8 s		162.2 s	
2	98.1 d	6.59 s	97.1 d	6.40 s
3	161.0 s		161.4 s	
4	108.8 s		110.8 s	
5	110.8 d	6.85 d (8.3)	112.1 d	6.87 d (8.3)
6	135.9 d	7.42 t (8.3)	135.2 d	7.48 t (8.3)
7	106.5 d	6.70 d (8.3)	106.1 d	6.62 d (8.3)
8	165.4 s		165.5 s	
9	182.2 s		183.5 s	
4a	155.3 s		156.8 s	
8a	107.4 s		107.2 s	
9a	104.1 s		105.0 s	
10a	158.2 s		159.2 s	
1'	34.0 t	2.54 t (7.2)	63.1 d	5.13 dd (8.8, 3.2)
2'	63.8 t	3.67 t (7.2)	51.6 t	2.84 dd (15.4, 3.2) 2.43 dd (15.4, 8.8)
3'			205.0 s	
4'			31.5 q	2.14 s
3-OMe	56.0 q	3.85 s	55.8 q	3.83 s
8-OMe	56.2 q	3.81 s	56.1 q	3.80 s
Ar-OH		13.43 s		

^a data obtained in C₅D₅N, ^b data obtained in CD₃OD

Compounds (**2**) was also isolated as a yellow gum, and its molecular formula was determined as C₁₉H₁₈O₇ through HRESI-MS analysis (pseudomolecular ion [M+Na]⁺ at *m/z* 381.0956). The ¹H- and ¹³C spectra data of **2** was very similar to those of **1** (see Table 1), except for the hydroxyethyl unit in **1** was replaced by a 1-hydroxy-3-oxobutyl moiety²⁰ [δ_C 63.1 d, 51.6 t, 205.0 s, 31.5 q; δ_H 5.13 dd (8.8, 3.2), 2.84 dd (15.4, 3.2), 2.43 dd (15.4, 8.8), 2.14 s] in compound **2**. Two methoxy groups located at C-3 and C-8, a phenolic hydroxy group located at C-1, and the 1-hydroxy-3-oxobutyl moiety at C-4 were also be concluded by the analysis of its HMBC spectrum. To determine the absolute configuration of **2**, the circular dichroism (CD) analysis was employed. The experimental CD spectrum of **2** exhibited a positive Cotton effect (CE) at 219 nm and a negative CE near 246 nm. The CEs, optical rotation, and coupling constant values of **2** were in excellent agreement with these of known compound,²⁰ (1'*S*)-7-hydroxy-3-(1'-hydroxy-3'-butanoyl)chromone-5-carboxylic acid. Thus, compound **2** was determined as (*S*)-1-hydroxy-4-(1-hydroxy-3-oxobutyl)-3,8-dimethoxy-9*H*-xanthen-9-one.

Since xanthenes are known to exhibit potent cytotoxicity,^{2,11,12} the cytotoxicity of compounds **1-5** were tested using a previously reported procedure.²¹ All treatments were performed in triplicate. In the MTT assay, the IC₅₀ was defined as the concentration of the test compound resulting in a 50% reduction of absorbance compared with untreated cells. The cytotoxic abilities against NB4, A549, SHSY5Y, PC3, and MCF7 tumor cell lines by MTT-assay (with taxol as the positive control) were shown in Table 2. The results revealed that compound **1** showed high cytotoxicity against A549 and PC3 cell with IC₅₀ values of 2.8 and 3.4 μ M, and **2** showed high cytotoxicity against PC3 cell with IC₅₀ valves of 2.8 μ M, respectively. The other compounds also showed moderate cytotoxicity for some tested cell lines with IC₅₀ values below 10.

Table 2. Cytotoxicity data (IC₅₀, μ M) for compounds **1 - 5**

Compounds	NB4	A549	SHSY5Y	PC3	MCF7
1	5.2	2.8	8.2	3.4	7.5
2	6.1	8.5	7.1	2.8	6.6
3	5.4	>10	7.8	>10	5.5
4	>10	9.6	>10	8.8	7.2
5	6.7	>10	6.5	9.0	>10
taxol	0.03	0.02	0.2	0.2	0.1

NB4, human leukemia cell; A549, carcinomic human alveolar basal epithelial cell; SHSY5Y, human neuroblastoma cell; PC3, Human prostate cancer cell; MCF7, Human breast adenocarcinoma cell.

EXPERIMENTAL

General. Optical rotation was measured in Horiba SEPA-300 high sensitive polarimeter. UV spectra

were obtained using a Shimadzu UV-2401A spectrophotometer. IR spectra were obtained in KBr disc on a Bio-Rad Wininfrared spectrophotometer. ESI-MS were measured on a VG Auto Spec-3000 MS spectrometer. ^1H , ^{13}C , and 2D NMR spectra were recorded on Bruker DRX-500 instrument with TMS as internal standard. Column chromatography was performed on silica gel (200-300 mesh), or on silica gel H (10~40 μm , Qingdao Marine Chemical Inc., China). Second separation was performed by an Agilent 1100 HPLC equipped with ZORBAX- C_{18} (21.2 mm \times 250 mm, 7.0 μm) column and DAD detector.

Plant material. The leaves and stems of *Hypericum chinense* L. were collected in Xishuangbanna Prefecture, Yunnan Province, People's Republic of China, in September 2010. The identification of the plant material was verified by Prof. Ren P. Y (Xishuangbanna Botanical Garden). A voucher specimen (YNNI-2010-9-22) has been deposited in our laboratory.

Extraction and Isolation. The air-dried and powdered leaves and stems of *H. chinense* (4.0 kg) were extracted four times with 70% acetone (4 \times 6 L) at room temperature and filtered. The crude extract (256 g) was applied to silica gel (200–300 mesh) column chromatography, eluting with a CHCl_3 -acetone gradient system (9:1, 8:2, 7:3, 6:4, 5:5), to give five fractions A–E. The further separation of fraction A (9:1, 18.5 g) by silica gel column chromatography, eluted with petroleum ether-EtOAc (9:1, 8:2, 7:3, 6:4, 1:1), yielded the mixtures A1–A5. The subfraction A2 (8:2, 4.2 g) was subjected to preparative HPLC (68% MeOH, flow rate 12 mL/min) to give **5** (14.8 mg). The further separation of subfraction A3 (7:3, 3.8 g) by silica gel column chromatography, and preparative HPLC (55~65% MeOH, flow rate 12 mL/min) to give **1** (8.5 mg), **2** (10.3 mg), **3** (15.7 mg), and **4** (16.0 mg).

Cytotoxicity Assay. The IC_{50} values of compounds were measured using the MTT assay. The MTT assay is a colorimetric assay for measuring the activity of cellular enzymes that reduce the tetrazolium dye, MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide, a yellow tetrazole), to its insoluble formazan, giving a purple color. Firstly, 2500 cells suspended in 100 μL MEM medium were seeded, respectively, in a 96-well plate. After 24 h incubation, fresh medium containing various concentrations of each compound were added into the 96-well plate to replace the old medium. The concentrations were ranged from 100 μM to 1.5625 μM , which was achieved by doing twofold dilutions six times. The OD_{595} values of the control groups at 0 h and 72 h together with the compound treated groups at 72 h from the MTT assay were measured using a plate reader. IC_{50} is the concentration of a compound inhibiting 50% of the cell growth.

1-Hydroxy-4-(2-hydroxyethyl)-3,8-dimethoxy-9H-xanthen-9-one (1): Obtained as a yellow gum; UV (MeOH) λ_{max} (log ϵ) 210 (4.15), 245 (3.38), 302 (3.92) nm; IR (KBr) ν_{max} 3432, 3078, 2930, 2885, 1658, 1602, 1546, 1468, 1375, 1186, 1065, 885, 764 cm^{-1} ; ESIMS m/z (positive ion mode) 339 $[\text{M}+\text{Na}]^+$; HRESIMS (positive ion mode) m/z 339.0840 $[\text{M}+\text{Na}]^+$ (calcd $\text{C}_{17}\text{H}_{16}\text{O}_6\text{Na}$ for 339.0845).

(S)-1-Hydroxy-4-(1-hydroxy-3-oxobutyl)-3,8-dimethoxy-9H-xanthen-9-one (2): Obtained as a yellow gum; $[\alpha]_D^{22.5}$ -45.6 (*c* 0.20, MeOH); UV (MeOH) λ_{\max} (log ϵ) 210 (4.15), 246 (3.78), 305 (3.78) nm; CD (MeOH, *c* 0.25) $\Delta\epsilon_{219}$ +0.94, $\Delta\epsilon_{237}$ -5.56, $\Delta\epsilon_{278}$ +0.28, $\Delta\epsilon_{324}$ -0.92; IR (KBr) ν_{\max} 3425, 3062, 2872, 2806, 1705, 1649, 1600, 1568, 1472, 1349, 1167, 1059, 875, 764 cm^{-1} ; ESIMS *m/z* (positive ion mode) 381 $[\text{M}+\text{Na}]^+$; HRESIMS (positive ion mode) *m/z* 381.0956 $[\text{M}+\text{Na}]^+$ (calcd $\text{C}_{19}\text{H}_{18}\text{NaO}_7$ for 381.0950).

ACKNOWLEDGMENT

This research was supported by the National Natural Science Foundation of China (No. 21002085 and 21362044), the Excellent Scientific and Technological Team of Yunnan High School (2010CI08), the Yunnan University of Nationalities Green Chemistry and Functional Materials Research for Provincial Innovation Team (2011HC008), the National Undergraduates Innovating Experimentation Project (2011HX18), and start-up funds of Yunnan University of Nationalities.

REFERENCES

1. M. M. Pinto, M. E. Sousa, and M. S. Nascimento, [Curr. Med. Chem., 2005, 12, 2517](#).
2. K. S. Masters and S. Bräse, [Chem. Rev., 2012, 112, 3717](#).
3. T. W. Cao, C. A. Geng, Y. B. Ma, K. He, H. L. Wang, N. J. Zhou, X. M. Zhang, Y. D. Tao, and J. J. Chen, [Planta Med., 2013, 79, 697](#).
4. Y. P. Wu, W. Zhao, Z. Y. Xia, G. H. Kong, X. P. Lu, Q. F. Hu, and X. M. Gao, [Molecules, 2013, 18, 9663](#).
5. S. Klaiklay, Y. Sukpondma, V. Rukachaisirikul, and S. Phongpaichit, [Phytochemistry, 2013, 85, 161](#).
6. H. R. Dharmaratne, Y. Sakagami, K. G. Piyasena, and V. Thevanesam, [Nat. Prod. Res., 2013, 27, 938](#).
7. S. Udomchotphruet, P. Phuwapraisirisan, J. Sichaem, and S. Tip-Pyang, [Phytochemistry, 2012 73, 148](#).
8. C. Uvarani, K. Chandraprakash, M. Sankaran, A. Ata, and P. S. Mohan, [Nat. Prod. Res., 2012, 26, 1265](#).
9. M. Ali, M. Arfan, M. Ahmad, K. Singh, I. Anis, H. Ahmad, M. I. Choudhary, and M. R. Shah, [Planta Med., 2011, 77, 2013](#).
10. Q. B. Han, N. Y. Yang, H. L. Tian, C. F. Qiao, J. Z. Song, D. C. Chang, S. L. Chen, K. Q. Luo, and H. X. Xu, [Phytochemistry, 2008, 69, 2187](#).
11. Q. F. Hu, D. Y. Niu, X. L. Li, Y. H. Qin, Z. Y. Yang, G. L. Zhao, Z. X. Yang, X. M. Gao, and Z. Y. Chen, [Heterocycles, 2013, 87, 1127](#).

12. N. Tanaka, Y. Kashiwada, S. Y. Kim, M. Sekiya, Y. Ikeshiro, and Y. Takaishi, [*Phytochemistry*, 2009, **70**, 1456.](#)
13. Z. Y. Xiao and Q. Mu, *Nat. Prod. Res. Dev.*, 2007, **19**, 344.
14. S. Abe, N. Tanaka, and J. Kobayashi, [*J. Nat. Prod.*, 2012, **75**, 484.](#)
15. N. Tanaka, S. Abe, K. Hasegawa, M. Shiro, and J. Kobayashi, [*Org. Lett.*, 2011, **13**, 5488.](#)
16. W. Wang, Y. H. Zeng, K. Osman, K. Shinde, M. Rahman, S. Gibbons, and Q. Mu, [*J. Nat. Prod.*, 2010, **73**, 1815.](#)
17. N. Tanaka and Y. Takaishi, [*Phytochemistry*, 2006, **67**, 2146.](#)
18. S. Udomchotphruet, P. Phuwapraisirisan, J. Sichaem, and S. Tip-pyang, [*Phytochemistry*, 2012, **73**, 148.](#)
19. Y. P. Wu, W. Zhao, Z. Y. Xia, G. H. Kong, X. P. Lu, Q. F. Hu, and X. M. Gao, [*Phytochem. Lett.*, 2013, **6**, 629.](#)
20. M. Gan, Y. Liu, Y. Bai, Y. Guan, L. Li, R. Gao, W. He, X. You, Y. Li, L. Yu, and C. Xiao, [*J. Nat. Prod.*, 2013, **76**, 1535.](#)
21. X. M. Gao, R. R. Wang, D. Y. Niu, C. Y. Meng, L. M. Yang, Y. T. Zheng, G. Y. Yang, Q. F. Hu, H. D. Sun, and W. L. Xiao, [*J. Nat. Prod.*, 2013, **76**, 1052.](#)