

HETEROCYCLES, Vol. 92, No. 6, 2016, pp. 1102 - 1110. © 2016 The Japan Institute of Heterocyclic Chemistry
Received, 4th March, 2016, Accepted, 25th March, 2016, Published online, 5th April, 2016
DOI: 10.3987/COM-16-13452

CONCISE SYNTHESIS AND BIOLOGICAL EVALUATION OF CHALCONE DERIVATIVES BEARING N-HETEROCYCLIC MOIETIES

Zewei Mao,^a Xi Zheng,^b Yuping Lin,^a Yan Qi,^b Chunyan Hu,^a Chunping Wan,^{*b}
and Gaoxiong Rao^{*a}

^a School of Traditional Chinese Medicine, Yunnan University of Traditional Chinese Medicine, Kunming, 650500, China; E-mail: rao13987124569@qq.com.

^b Central laboratory, The NO.1 Affiliated Hospital of Yunnan University of Traditional Chinese Medicine, Kunming, 650021, China; E-mail: wanchunping1012@163.com

Abstract – In this study, we designed and synthesized a series of chalcone derivatives bearing N-heterocyclic moieties, and screened *in vitro* anti-inflammatory in lipopolysaccharide (LPS)-stimulated RAW-264.7 macrophages and anticancer activity against a panel of human tumor cell lines. The results indicated that compound **9** not only had inhibitory effect on the generation of NO (IC₅₀=8.11 μM) and significantly inhibited the production of TNF-α, but also showed better anticancer activity against Hela and SGC7901 (IC₅₀=1.05 μM and 6.47 μM, respectively), which was identified as the most potent anti-inflammatory and anticancer agent.

The secretion of inflammatory mediators by macrophages is involved in both the innate and adaptive immune responses of many human diseases, such as infection and rheumatoid arthritis (RA).¹ LPS, a component of the cell wall of gram-negative bacteria, is known to activate a number of cellular signals in macrophages. The macrophage produce many chemokines (NO, PGE₂) and pro-inflammatory cytokines (TNF-α, IL-6, IL-1β) when the macrophages are stimulated by LPS. So, the pro-inflammatory mediator is considered as potential target in inflammatory-associated diseases. On the other hand, cancer is the second most common cause of death in the world, which still remains a potentially life threatening disease resulting from uncontrolled cell growth.


Chalcones play an important role in living organisms² with a wide range of biological activities, which are widely used as anti-inflammatory, antibacterial, antitumor and so on.³⁻⁶ Moreover, chalcones are

widely present in nature, and a number of anti-inflammatory and anticancer chalcones have been reported in recent years, including sappanchalcone,⁷ butein,⁸ licochalcone A⁹ and echinatin.¹⁰ Due to their flexible character bearing α , β -unsaturated ketone unit, chalcones are playing the vital role in drug research through structural modification.¹¹⁻¹⁴

Nitrogen heterocycles are important class of compounds having versatile biological activities, which are used in drug design and synthesis generally as active units. The combination of the chalcone and imidazole in one frame, has been reported in previous literature and is likely to lead to hybrid compounds with potential activity.¹⁵ Moreover, in previous work, we have reported the synthesis of chalcone-piperazine hybrid compounds and their potential anticancer activities.¹⁶ On the basis of these results, we wanted to design and synthesize molecules towards the recombination of chalcone and active nitrogen heterocycle moieties in the same structure through medicinal chemistry hybridization. Here, we reported the concise synthesis of chalcone derivatives containing nitrogen heterocycle compound, and screened the *in vitro* anti-inflammatory potential of compounds in LPS-stimulated inflammatory response in RAW-264.7 macrophages and anticancer activity against a panel of human tumor cell lines by MTT assay.
















The general concise synthetic route used to obtain title compounds is outlined in Scheme 1. Treatment of commercially available 4-fluoroacetophenone with 4-(dimethylamino)benzaldehyde with 4-fluoroacetophenone gave the 4-(dimethylamino)-4'-fluorochalcone (**1**)¹⁷ in the presence of KOH resulted by the Aldol condensation, and it was in *trans-configuration*. The desired compounds were formed from chalcone intermediate by subsequent substitution reaction with different active nitrogen heterocycle compounds in the presence of Cs₂CO₃ at 110 °C in DMF in good yields (**2-16**) following the literature procedure, and no aza-Michael addition product was observed.^{18,19}

Initially, K₂CO₃ was used in synthesis of title compounds. The results showed that compounds **8-16** were obtained in satisfied yields, but compounds **2-7** were in low yields. Replaced by Cs₂CO₃, chalcone derivatives **2-16** were in excellent yields, since the basicity of imidazole and benzimidazole was less than that of other nitrogen active compounds.



Scheme 1. Concise synthesis of chalcone derivatives

Table 1. Structures and yields of compounds

Compd.	Het	M p (°C)	Yields (%) ^a	Compd.	Het	M p (°C)	Yields (%) ^a
2		164-166	91	10		162-164	85
3		165-167	71	11		179-181	86
4		166-168	79	12		161-163	85
5		179-181	76	13		164-166	77
6		180-181	67	14		174-176	65
7		180-182	69	15		157-158	84
8		170-172	88	16		157-159	83
9		164-165	78				

^a Yields represent isolated yields.

Comparative data for these compounds with respect to structures and yield are provided in Table 1. All of the synthesized compounds were characterized by ¹H NMR and ¹³C NMR, and some representative compounds were characterized by HRMS analysis.

Anti-inflammatory activity: RAW 264.7 cell, a murine macrophage cell line, is widely used to establish inflammatory model in vitro. In this study, we investigated the anti-inflammatory activity of synthetic derivatives in LPS-induced RAW 264.7 on the generation of NO and pro-inflammatory cytokines.

To evaluate the effect of chalcone analogs (**2-16**) on NO generation, the RAW 264.7 cells were stimulated with LPS for 24 h, the level of NO from cell culture supernatant was measured by Griess reagent. The results indicated that compounds **8** and **9** showed significant inhibitory effect on the generation of NO in LPS-induced RAW 264.7, the IC₅₀ value was calculated to be about 10.13 μM and 8.11 μM respectively (Table 2), neither had showed obvious cell cytotoxicity to RAW 264.7 cells at the concentration of 100 μM.

Stimulated macrophages have great impact on inflammatory diseases via excess production of cytokines, including tumor necrosis factor-α (TNF-α), interleukin-1β (IL-1β), IL-6, and other inflammatory mediators such as nitric oxide (NO). In order to detect the effect of compound on pro-inflammatory cytokines (TNF-α, IL-6), we then selected the compound **9** for further study. The production of TNF-α in LPS-induced RAW 264.7 was significantly down-regulated after exposed to indicate concentration of compound **9**. However, interestingly, it had no apparent effects on the generation of IL-6 (Figure. 1). At

present, our results indicated that compound **9** significantly inhibited the production of TNF- α and NO in LPS-induced RAW264.7.

Table 2. Anti-inflammatory activities of compounds

Compd.	NO generation (IC ₅₀ , μ M) ^a	Compd.	NO generation (IC ₅₀ , μ M) ^a
2	17.12	10	16.68
3	21.35	11	26.10
4	30.72	12	>40
5	>40	13	>40
6	>40	14	19.07
7	>40	15	>40
8	10.13	16	>40
9	8.11		

^a Values represent the concentration required to produce 50% inhibition of the response.




Figure 1. The effect of compound **9** on TNF- α and IL-6 production in LPS-induced RAW 264.7. The RAW 264.7 cell was stimulated with LPS in the present or absence of various concentration of compound **9** for 24h, the culture supernatant was harvested to measure the production of TNF- α and IL-6 by ELISA. The results were expressed as mean \pm s.e.m with three independent experiments. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ versus control group.

Anticancer activity: Cytotoxic activity of novel synthesized hybrid derivatives were evaluated against human lung cancer cell line A549, human cervical carcinoma Hela and human gastric carcinoma SGC7901 by MTT assay, cisplatin (DDP) was used as reference drug. The biological results of hybrid compounds are summarized in Tables 3.

As shown in Table 3, the N-heterocycle moieties of the hybrid compounds have an obvious influence on the antitumor activities. To our delight, compounds bearing imidazole, triazole, piperazine or 3-hydroxyazetidine displayed potent or similar cytotoxic activity against human tumor cell lines compared to DDP. Compound **2** displayed potent anticancer activity against A549 (10.21 μ M) and Hela (7.42 μ M), compound **8** appeared good cytotoxic activity against SGC7901 (9.53 μ M), and compound **11** showed potent anticancer activity against Hela (6.58 μ M). Especially, among all derivatives, compound **9** showed the best IC₅₀ value against Hela (1.05 μ M) and SGC7901 (6.47 μ M), which was identified as the most potent anticancer agent. However, substitution at the N-heterocycle with

other substituted groups resulted in lower activity. For example, compounds **3-4** and **10-11** were less active than compounds **2** and **9**, respectively.

Table 3. *In vitro* cytotoxic activities of compounds

Compd.	Cell lines (IC ₅₀ , μM) ^a			Compd.	Cell lines (IC ₅₀ , μM) ^a		
	A549	Hela	SGC7901		A549	Hela	SGC7901
2	10.21	7.42	>40	10	>40	12.53	19.34
3	21.64	17.11	>40	11	17.05	6.58	25.26
4	>40	>40	>40	12	28.78	>40	>40
5	26.50	>40	19.73	13	>40	25.16	>40
6	33.14	>40	24.82	14	7.72	13.01	>40
7	>40	>40	>40	15	>40	>40	27.24
8	>40	16.23	9.53	16	23.21	>40	>40
9	>40	1.05	6.47	DDP	11.54	20.52	12.44

^a Each value was reproduced in triplicate.

Finally, we discussed a clear tendency of structure-activity relationship: Comparison of the activities of the chalcone derivatives **8** and **9** revealed that compounds bearing piperazine or triazole moiety showed significant anti-inflammatory activity, and compounds **2**, **8**, **9**, **11** and **14** bearing imidazole, triazole, piperazine or 3-hydroxyazetidine moiety displayed potent antitumor activity. Interestingly, both anti-inflammatory and anticancer activity would be decreased when substitution at the N-heterocycle moiety with other groups except pyrimidyl. Further derivatization and biological research is under progress in our laboratory and will be reported in due course.

EXPERIMENTAL

Starting materials were commercially available and analytically pure. Melting points were measured on YANACO microscopic melting point meter (Yanaco, Japan) and were uncorrected. ¹H NMR and ¹³C NMR spectras were recorded on a Bruker AV 300 spectrometer (Bruker Company, Germany), using TMS as internal standard and CDCl₃ as solvent, respectively. TLC analysis was carried out on silica gel plates GF254 (Qindao Haiyang Chemical, China). High-resolution mass spectra were performed on an ESI Q-TOF MS spectrometer (Micromass, England).

Synthesis of 4-Dimethylamino-4'-fluorochalcone (1): To a solution of EtOH (15 mL), 4-dimethylaminobenzaldehyde (1.49 g, 10 mmol) and 4-fluoroacetophenone (1.38 g, 10 mmol), was added 20% KOH (10mL) and left to react for 16 h at rt. The reaction was quenched by the addition of water (30 mL) and was extracted with CHCl₃ (3×20 mL). The organic layer was dried using anhydrous sodium sulfate, concentrated in vacuo and purified by column chromatography (10% EtOAc/PE, 2.35g, 87% yield). Yellow solid; mp 139-141°C; ¹H NMR (300 MHz, CDCl₃) δ: 8.01-8.05 (m, 2H), 7.81 (d, *J*=15.3 Hz, 1H), 7.55 (d, *J*=8.7 Hz, 2H), 7.32 (d, *J*=15.2 Hz, 1H), 7.11-7.17 (m, 2H), 6.69 (d, *J*=8.7Hz, 2H), 3.03 (s, 6H);

^{13}C NMR (75 MHz, CDCl_3) δ : 188.9, 166.9, 163.6, 152.1, 146.0, 135.4, 130.9, 130.7, 130.5, 122.5, 116.4, 115.6, 115.3, 111.8, 40.1.

General procedure for the preparation of chalcone derivatives 2-16. To a stirred solution of the 4-dimethylamino-4'-fluorochalcone **1** (1.0 mmol) and Cs_2CO_3 (2.0 mmol) in dried DMF (5 mL), nitrogen heterocycle (1.5 mmol) was added and reaction mixture was stirred for 12-15 h at 110 °C. After completion of the reaction as indicated by TLC, The reaction was quenched by the addition of CHCl_3 (20 mL) and was washed with water (3×20 mL). The organic layer was dried using anhydrous sodium sulfate, concentrated in vacuo and purified by column chromatography to afford **2-16** in 55-91% yields.

4-Dimethylamino-4'-(1-imidazolyl)chalcone (2). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.14 (d, $J=8.7$ Hz, 2H), 7.95 (s, 1H), 7.85 (d, $J=15.3$ Hz, 1H), 7.57 (d, $J=8.7$ Hz, 2H), 7.51 (d, $J=8.4$ Hz, 2H), 7.36 (d, $J=4.5$ Hz, 2H), 7.24 (s, 1H), 6.71 (d, $J=8.7$ Hz, 2H), 3.05 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.8, 152.3, 146.5, 140.0, 137.9, 135.4, 131.0, 130.6, 130.2, 122.4, 120.7, 117.8, 116.1, 111.8, 40.1; HRMS: m/z calcd for $\text{C}_{20}\text{H}_{20}\text{N}_3\text{O}$ ($\text{M}+\text{H}$) $^+$ 318.1601, found 318.1601.

4-Dimethylamino-4'-(2-methyl-1-imidazolyl)chalcone (3). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.11 (d, $J=5.7$ Hz, 2H), 7.90 (d, $J=8.4$ Hz, 1H), 7.55 (d, $J=8.4$ Hz, 2H), 7.34-7.47 (m, 2H), 7.04 (s, 2H), 6.62-6.71 (m, 3H), 3.01 (s, 6H), 2.40 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 186.8, 158.5, 153.8, 152.2, 144.8, 141.3, 138.3, 131.2, 130.9, 129.5, 128.4, 125.4, 125.3, 125.0, 122.5, 120.6, 113.7, 112.1, 111.2, 40.3, 14.3.

4-Dimethylamino-4'-(2-ethyl-1-imidazolyl)chalcone (4). Brown solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.13 (d, $J=5.1$ Hz, 2H), 7.87 (d, $J=15.6$ Hz, 1H), 7.58 (d, $J=9.0$ Hz, 2H), 7.28-7.42 (m, 3H), 7.10 (d, $J=15.6$ Hz, 2H), 6.71 (d, $J=9.0$ Hz, 2H), 3.05 (s, 6H), 2.76 (q, $J=7.5$ Hz, 2H), 1.30 (t, $J=7.5$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 189.2, 152.3, 149.5, 146.7, 140.8, 138.7, 130.7, 129.6, 128.0, 125.5, 122.3, 120.4, 116.1, 111.8, 40.1, 20.7, 12.4.

4-Dimethylamino-4'-(1-benzimidazolyl)chalcone (5). Pale yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.19 (d, $J=8.4$ Hz, 2H), 8.15 (s, 1H), 7.86-7.90 (m, 1H), 7.81 (s, 1H), 7.51 - 7.60 (m, 5H), 7.31-7.36 (m, 3H), 6.66 (d, $J=9.0$ Hz, 2H), 2.99 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.7, 152.2, 146.5, 144.3, 142.0, 139.3, 138.2, 133.2, 130.6, 130.2, 124.0, 123.2, 123.1, 122.3, 120.7, 115.9, 111.8, 110.6, 40.0.

4-Dimethylamino-4'-(2-methyl-1-benzimidazolyl)chalcone (6). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.24 (d, $J=8.4$ Hz, 2H), 7.91 (d, $J=15.6$ Hz, 1H), 7.78 (d, $J=7.8$ Hz, 1H), 7.61 (d, $J=9.0$ Hz, 2H), 7.53 (d, $J=8.7$ Hz, 2H), 7.49 (d, $J=15.6$ Hz, 1H), 7.20-7.30 (m, 3H), 6.73 (d, $J=8.7$ Hz, 2H), 3.07 (s, 6H), 2.57 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 189.3, 152.3, 146.8, 142.7, 139.3, 130.7, 130.0, 126.9, 122.9, 122.7, 119.2, 116.1, 111.8, 109.9, 40.1, 14.6.

4-Dimethylamino-4'-(5, 6-dimethyl-1-benzimidazolyl)chalcone (7). Brown solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.22 (d, $J=8.4$ Hz, 2H), 8.07 (s, 1H), 7.88 (d, $J=15.6$ Hz, 1H), 7.55-7.64 (m, 5H), 7.39 (s, 2H),

6.71 (d, $J=8.7$ Hz, 2H), 3.04 (s, 6H), 2.40 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 189.0, 152.2, 146.6, 142.9, 141.2, 139.7, 138.0, 133.3, 132.1, 131.7, 130.7, 130.2, 123.1, 122.4, 120.7, 116.0, 111.8, 110.7, 40.1, 20.7, 20.3.

4-Dimethylamino-4'-(1-triazolyl)chalcone (8). yellow solid; ^1H NMR (300MHz, CDCl_3) δ : 8.67 (s, 1H), 8.17 (s, 1H), 8.14 (s, 2H), 7.85 (d, $J=15.6$ Hz, 1H), 7.84 (d, $J=8.7$ Hz, 2H), 7.57 (d, $J=8.7$ Hz, 2H), 7.35 (d, $J=15.6$ Hz, 1H), 6.71 (d, $J=8.7$ Hz, 2H), 3.05 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.8, 152.9, 152.3, 146.6, 141.0, 139.4, 138.5, 130.6, 130.1, 122.4, 119.4, 116.1, 111.8, 40.1; HRMS: m/z calcd for $\text{C}_{19}\text{H}_{19}\text{N}_4\text{O}$ ($\text{M}+\text{H}$) $^+$ 319.1553, found 319.1554.

4-Dimethylamino-4'-(1-piperazinyl)chalcone (9). brown solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.01 (d, $J=9.0$ Hz, 2H), 7.80 (d, $J=15.3$ Hz, 1H), 7.56 (d, $J=9.0$ Hz, 2H), 7.41 (d, $J=15.6$ Hz, 1H), 6.92 (d, $J=9.0$ Hz, 2H), 6.70 (d, $J=9.0$ Hz, 2H), 3.33 (t, $J=4.8$ Hz, 4H), 3.01-3.04 (m, 10H), 2.12 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.3, 154.2, 151.8, 144.1, 130.3, 130.1, 129.1, 123.0, 116.8, 113.6, 111.8, 48.5, 45.9, 40.2; HRMS: m/z calcd for $\text{C}_{21}\text{H}_{26}\text{N}_3\text{O}$ ($\text{M}+\text{H}$) $^+$ 336.2070, found 336.2072.

4-Dimethylamino-4'-(N-methyl-1-piperazinyl)chalcone (10). Brown solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.00 (d, $J=9.0$ Hz, 2H), 7.80 (d, $J=15.6$ Hz, 1H), 7.55 (d, $J=8.7$ Hz, 2H), 7.39 (d, $J=15.6$ Hz, 1H), 6.92 (d, $J=8.7$ Hz, 2H), 6.70 (d, $J=9.0$ Hz, 2H), 3.38 (t, $J=5.1$ Hz, 4H), 3.01 (s, 6H), 2.57 (t, $J=5.1$ Hz, 4H), 2.34 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.3, 153.8, 151.8, 144.1, 130.3, 130.1, 129.2, 123.1, 116.9, 113.7, 111.9, 54.8, 47.3, 46.1, 40.1.

4-Dimethylamino-4'-(N-(2-pyrimidyl)-1-Piperazinyl)chalcone (11). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.35 (d, $J=4.8$ Hz, 2H), 8.03 (d, $J=8.7$ Hz, 2H), 7.81 (d, $J=15.6$ Hz, 1H), 7.56 (d, $J=8.7$ Hz, 2H), 7.40 (d, $J=15.3$ Hz, 1H), 6.96 (d, $J=9.0$ Hz, 2H), 6.70 (d, $J=9.0$ Hz, 2H), 6.55 (t, $J=4.8$ Hz, 1H), 4.01 (t, $J=5.4$ Hz, 4H), 3.47 (t, $J=5.4$ Hz, 4H), 3.02 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.4, 161.6, 157.8, 153.7, 151.8, 144.2, 130.4, 130.1, 129.4, 123.1, 116.9, 113.8, 111.9, 110.3, 47.3, 43.3, 40.1.

4-Dimethylamino-4'-(1-pyrazolyl)chalcone (12). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.14 (d, $J=8.7$ Hz, 2H), 8.02 (d, $J=2.1$ Hz, 1H), 7.77-7.85 (m, 4H), 7.58 (d, $J=8.7$ Hz, 2H), 7.38 (d, $J=15.3$ Hz, 1H), 6.70 (d, $J=8.7$ Hz, 2H), 6.52 (t, $J=1.8$ Hz, 1H), 3.04 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 189.0, 152.1, 146.0, 142.7, 141.9, 136.7, 130.5, 129.9, 126.9, 122.5, 118.4, 116.3, 111.8, 108.4, 40.1; HRMS: m/z calcd for $\text{C}_{20}\text{H}_{20}\text{N}_3\text{O}$ ($\text{M}+\text{H}$) $^+$ 318.1601, found 318.1603.

4-Dimethylamino-4'-(3,5-dimethyl-1-pyrazolyl)chalcone (13). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.11 (d, $J=8.7$ Hz, 2H), 7.84 (d, $J=15.6$ Hz, 1H), 7.59 (d, $J=6.6$ Hz, 2H), 7.57 (d, $J=7.2$ Hz, 2H), 7.36 (d, $J=15.6$ Hz, 1H), 6.71 (d, $J=8.7$ Hz, 2H), 6.04 (s, 1H), 3.04 (s, 6H), 2.38 (s, 3H), 2.32 (s, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ : 189.6, 152.1, 149.8, 146.3, 142.9, 139.6, 137.2, 130.5, 129.3, 123.8, 122.5, 116.6, 111.8, 108.1, 40.1, 13.6, 12.8.

4-Dimethylamino-4'-(3-hydroxyazetid-1-yl)chalcone (14). Pale yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 7.94 (d, $J=8.7$ Hz, 2H), 7.78 (d, $J=15.3$ Hz, 1H), 7.54 (d, $J=8.7$ Hz, 2H), 7.38 (d, $J=15.3$ Hz, 1H), 6.70 (d, $J=8.7$ Hz, 2H), 6.40 (d, $J=8.7$ Hz, 2H), 4.77-4.81 (m, 1H), 4.27 (t, $J=8.1$ Hz, 2H), 3.79-3.83 (dd, $J=4.5$ Hz, 4.5 Hz, 2H), 3.03 (s, 6H); HRMS: m/z calcd for $\text{C}_{20}\text{H}_{23}\text{N}_2\text{O}_2$ ($\text{M}+\text{H}$) $^+$ 323.1754, found 323.1750.

4-Dimethylamino-4'-(1-pyrrolyl)chalcone (15). Yellow solid; ^1H NMR (300 MHz, CDCl_3) δ : 8.00 (d, $J=9.0$ Hz, 2H), 7.79 (d, $J=15.3$ Hz, 1H), 7.54 (d, $J=8.7$ Hz, 2H), 7.43 (d, $J=15.3$ Hz, 1H), 6.68 (d, $J=9.0$ Hz, 2H), 6.53 (d, $J=8.7$ Hz, 2H), 3.34 (t, $J=6.9$ Hz, 4H), 2.98 (s, 6H), 1.96-2.01 (m, 4H); ^{13}C NMR (75 MHz, CDCl_3) δ : 187.9, 151.6, 150.7, 143.2, 130.7, 130.0, 126.2, 123.4, 117.2, 111.9, 110.9, 47.6, 40.2, 25.4.

4-Dimethylamino-4'-(1-piperidyl)chalcone (16). Brown solid; ^1H NMR (300 MHz, CDCl_3) δ : 7.99 (d, $J=9.0$ Hz, 2H), 7.80 (d, $J=15.6$ Hz, 1H), 7.55 (d, $J=8.7$ Hz, 2H), 7.41 (d, $J=15.6$ Hz, 1H), 6.90 (d, $J=8.7$ Hz, 2H), 6.69 (d, $J=8.7$ Hz, 2H), 3.35 (s, 4H), 3.01 (s, 6H), 1.65 (s, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ : 188.2, 154.2, 143.8, 130.5, 130.0, 128.2, 123.3, 117.0, 113.5, 111.9, 48.7, 40.2, 25.4, 24.4.

Antimicrobial Activity. Briefly, murine RAW264.7 macrophages (American Type Culture Collection, Manassas) were plated in 96 well plate at a density of 1×10^5 cells/well and stimulated with $1 \mu\text{g}/\text{mL}$ LPS in the present or absence of various concentration of compound for 24 h, the culture supernatant were harvested to measure NO and cytokines production. The production of NO was determined by assaying culture supernatant for NO^{2-} , a stable reaction product of NO, $100 \mu\text{L}$ of supernatant was mixed with an equal volume of Griess reagent [1% sulfanilamide and 0.1% N-(1-naphthyl)ethylenediamine dihydrochloride in 2.5% H_3PO_4] at room temperature for 10 min. Absorbance was measured at 540 nm in a microplate reader. Nitrite concentration was calculated from a NaNO_2 standard curve. For the cytokines detection, culture supernatants were harvested at 36 h to measure IL-6 and TNF- α level by ELISA following the manufacture's instruction.

Antitumor activity: About 1×10^4 cell/well were seeded into 96-well microtiter plates. After 24 h post-seeding, cells were treated with vehicle control or various concentrations of samples for 48 h. $20 \mu\text{L}$ of MTT solution (5 mg/mL) was added to each well and the tumor cells were incubated at 37°C in a humidified atmosphere of 5% CO_2 air for 4 h. Upon removal of MTT/medium, $150 \mu\text{L}$ of DMSO was added to each well and the plate was agitated at oscillator for 5 min to dissolve the MTT-formazan. The assay plate was read at a wavelength of 570 nm using a microplate reader.

ACKNOWLEDGEMENTS

This work was financially supported by the Natural Science Foundation of China (81460624) and the Application Basic Research Program of Yunnan Province (2014FZ087).

REFERENCES

1. W. L. Smith and L. J. Marnett, *Biochim. Biophys. Acta*, 1991, **1083**, 1.
2. F. Berrue, M. W. B. McCulloch, and R. G. Kerr, *Bioorg. Med. Chem.*, 2011, **19**, 6702.
3. L. Jayasinghe, B. A. I. S. Balasooriya, W. C. Padmini, N. Hara, and Y. Fujimoto, *Phytochemistry*, 2004, **65**, 1287.
4. K. Sugamoto, Y. I. Matsusita, K. Matsui, C. Kurogi, and T. Matsui, *Tetrahedron*, 2011, **67**, 5346.
5. N. Mishra, P. Arora, B. Kumar, L. C. Mishra, A. Bhattacharya, S. K. Awasthi, and V. K. Bhasin, *Eur. J. Med. Chem.*, 2008, **43**, 1530.
6. Z. Nowakowska, B. Kedzia, and G. Schroeder, *Eur. J. Med. Chem.*, 2008, **43**, 707.
7. G. S. Jeong, D. S. Lee, B. Li, H. J. Lee, E. C. Kim, and Y. C. Kim, *Eur. J. Pharmacol.*, 2010, **644**, 1.
8. Z. Wang, Y. Y. Lee, J. S. Eun, and E. J. Bae, *Eur. J. Pharmacol.*, 2014, **738**, 40.
9. M. Chen, S. B. Christensen, J. Blom, E. Lemmich, L. Nadelmann, K. Fich, T. G. Theander, and A. Kharazmi, *Antimicrob. Agents Chemother.*, 1993, **37**, 2550.
10. S. I. Ayabe and T. Furuya, *Tetrahedron Lett.*, 1981, **22**, 2097.
11. J. Chen, J. Yan, J. H. Hu, Y. Q. Pang, L. Huang, and X.S. Li, *RSC Adv.*, 2015, **5**, 68128.
12. Y. N. Shen, L. Lin, H.Y. Qiu, W. Y. Zou, Y. Qian, and H. L. Zhu, *RSC Adv.*, 2015, **5**, 23767.
13. K. V. Sashidhara, M. Kumar, R. K. Modukuri, R. Sonkar, G. Bhatia, A. K. Khanna, S. Rai, and R. Shukla, *Bioorg. Med. Chem. Lett.*, 2011, **21**, 4480.
14. C. Tang, L. H. Zhu, Y. Chen, R. Qin, Z. N. Mei, J. Xu, and G. Z. Yang, *RSC Adv.*, 2014, **4**, 10862.
15. V. K. Marrapu, V. Chaturvedi, S. Singh, S. Singh, S. Sinha, and K. Bhandari, *Eur. J. Med. Chem.*, 2011, **46**, 4302.
16. Z. W. Mao, X. Zheng, Y. Qi, M. D. Zhang, Y. Huang, C. P. Wan, and G. X. Rao, *RSC Adv.*, 2016, **6**, 7723.
17. A. Kamal, A. Mallareddy, P. Suresh, V. L. Nayak, R. V. C. R. N. C. Shetti, N. S. Rao, J. R. Tamboli, T. B. Shaik, M. V. P. S. Vishnuvardhan, and S. Ramakrishna, *Eur. J. Med. Chem.*, 2012, **47**, 530.
18. A. J. Watson and A. J. Fairbanks, *Eur. J. Org. Chem.*, 2013, **30**, 6784.
19. M. L. Kantam, J. Yadav, S. Laha, B. Sreedhar, and S. Jha, *Adv. Synth. Catal.*, 2007, **349**, 1938.