

HETEROCYCLES, Vol. 95, No. 1, 2017, pp. 624-635. © 2017 The Japan Institute of Heterocyclic Chemistry  
Received, 29th August, 2016, Accepted, 17th October, 2016, Published online, 15th December, 2016  
DOI: 10.3987/COM-16-S(S)48

## SYNTHESIS OF 2-ARYLAMINO-1-AZAAZULENES

**Satoru Tsukada,\* Makoto Nakazawa, Yuya Okada, Keito Ohtsu, Noritaka Abe, and Takahiro Gunji**


Department of Pure and Applied Chemistry, Faculty of Science and Technology,  
Tokyo University of Science, 2641 Yamazaki, Noda-shi, Chiba 278-8510, Japan  
E-mail: tsukada@rs.tus.ac.jp

**Abstract** – A series of 2-arylamino-1-azaazulenes were synthesized by the Buchwald-Hartwig cross coupling reaction. The resulting compounds were characterized by <sup>1</sup>H NMR, <sup>13</sup>C NMR, HRMS analyses, and elemental analysis. X-Ray crystallographic analysis revealed that 2-(1-naphthylamino)-1-azaazulene exists in dimeric form with two intermolecular N–H···N hydrogen bonds in its crystal structure.


Nitrogen-containing heteroaromatic compounds are one of the most popular and important class of organic compounds studied in different fields of modern day research. Azaazulenes are the structural isomers of quinoline and the heteroatomic analogues of azulene. In particular, 1-azaazulenes exhibit remarkable physical and chemical properties such as a large dipole moment ( $\mu = 3.05$  D),<sup>1</sup> unique reactivity, and potential biological activities.<sup>2-5</sup> In recent years, 2-amino-1-azaazulene derivatives have been studied as a ligand for metal ions,<sup>6,7</sup> and as anticancer agents.<sup>7,8</sup> Especially, 2-arylamino-1-azaazulenes are potentially bioactive and hence deserve more attention from the scientific community. However, there are limited reports on the synthesis, and chemical and physical properties. 2-Arylamino-1-azaazulenes were synthesized by only amination of 2-halo-1-azaazulenes.<sup>6,8-12</sup> The arylation of 2-amino-1-azaazulenes have not been investigated. Here we report the synthesis of a novel series of 2-arylamino-1-azaazulenes via Buchwald-Hartwig cross coupling, along with the study of their UV-Vis absorption properties. The arylations of 2-amino-1-azaazulenes were not succeeded under the same condition of amination of 2-haloazaazulenes. We found that 2-arylamino-1-azaazulenes were successfully synthesized by Buchwald-Hartwig cross coupling using phosphine ligand such as Xantphos and BINAP.

All 2-arylamino-1-azaazulenes were successfully synthesized by the Buchwald-Hartwig cross coupling reaction of aryl halide and arylamine. **1a** was synthesized by using a slightly modified version of the


method in our previous work.<sup>6,8</sup> The reaction of 2-bromo-1-azaazulene **2a** and 2-aminoazulene **3** in the presence of  $\text{Pd}_2(\text{dba})_3$ , Xantphos (9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene),<sup>13</sup>  $\text{Cs}_2\text{CO}_3$ , and 18-crown-6 (Scheme 1) gave **1a** in 24% yield. In this reaction, 1-(1-azaazulene-2-yl)-2-aminoazulene **4** was also obtained in 18% yield. Under a non-catalytic condition, both **1a** and **4** could not be obtained. This result indicated that **4** was generated as by-product of Pd catalyzed cross coupling reaction. However, only 32% of product yield was obtained when the same reagent system was applied for the synthesis of **1b**. In an attempt to enhance the product yield, the palladium catalyst and phosphine ligand were changed to  $\text{Pd}(\text{dppf})\text{Cl}_2$  and BINAP (2,2'-bis(diphenylphosphino)-1,1'-binaphthyl).<sup>14</sup> Under these reaction conditions, **1b** was obtained in 70% yield. **1c** was also obtained in 30% yield by using the same catalyst and ligand (Scheme 2). 2-Arylamino-1-azaazulenes **6**, **8**, **10**, and **12** were synthesized under the conditions similar to those used for **1a**, in 26%, 13%, 28%, and 51% yields, respectively (Scheme 3–5). **1b**, **1c**, and **8** could be also synthesized under a non-catalytic reaction in 9%, 3%, and 75% yields, respectively. On the other hands, **6**, **10**, and **12** were not obtained by a non-catalytic reaction. Therefore, the Buchwald-Hartwig cross coupling reaction of 2-aminoazaazulene was useful for synthesis of 2-arylamino-1-azaazulenes. We note that **8** was already synthesized by another method.<sup>12</sup>






Scheme 1



Scheme 2



Scheme 3



**Figure 1.** (a) ORTEP drawings of **10** with thermal ellipsoids at the 50% probability level. (b) Packing structure of **10**. Hydrogen bonds are displayed as broken lines. Selected bond lengths (Å): C1–N1 1.3629(13), C1–N2 1.3530(14), C2–N2 1.4179(13), N2–H1 0.915(14).


**Table 1.** Summary of crystal data of **10**

Empirical formula	C <sub>19</sub> H <sub>14</sub> N <sub>2</sub>	<i>T</i> / K	103
Formula weight	270.32	<i>D</i> <sub>calcd</sub> / g cm <sup>-3</sup>	1.293
Crystal system	Orthorhombic	$\mu$ (Mo K $\alpha$ ) / mm <sup>-1</sup>	0.077
Space group	Pbca	<i>F</i> (000)	1136
<i>a</i> / Å	21.558(2)	Reflection collected	13851
<i>b</i> / Å	10.2034(11)	Independent reflections	2549
<i>c</i> / Å	19.017(2)	<i>R</i> 1 ( <i>I</i> > 2 $\sigma$ ( <i>I</i> ))	0.0340
<i>V</i> / Å <sup>3</sup>	3793.9(7)	w <i>R</i> 2 (all data)	0.0887
<i>Z</i>	8	Goodness of fit on <i>F</i> <sup>2</sup>	1.040


The molecular structure of **10** was successfully determined by single-crystal X-ray diffraction analysis (Figure 1 and Table 1). To the best of our knowledge, this is the first report of the crystal structure of 2-aryl-amino-1-azaazulenes. Single crystals of **10** were obtained by recrystallization from ethyl acetate at

room temperature. The azaazulene core and amino group including the hydrogen on the N atom are highly planar. The dihedral angle between the plane of the azaazulene core and the naphthalene ring is  $51.62^\circ$ . The C1–N1 (1.3629(13) Å) and C1–N2 (1.3530(14) Å) bonds are slightly longer than the C2–N2 bond (1.4179(13) Å) and comparable to the aromatic C–N bond such as that in pyridine.<sup>15</sup> The aminoazaazulene unit forms two intermolecular N2–H $\cdots$ N1 hydrogen bonds (1.989 Å) between the two monomers (Figure 1b). In other words, **10** exists as a dimer in crystal state. The CH $\cdots$  $\pi$  interactions<sup>16-18</sup> between H on C8 of the azaazulene and naphthalene ring seems to be exist in a dimer form.

UV-Vis absorption spectra of 2-arylamino-1-azaazulenes and **2d** are shown in Figure 2. In comparison to **2d**, all 2-arylamino-1-azaazulenes showed a large molar extinction coefficient  $\epsilon$  and redshift. For example, the longest absorption wavelength of **1a** red-shifted by about 50 nm compared with that of **2d**. This indicates the delocalization of the  $\pi$  electrons on azaazulene and the amine through two tautomeric forms **1a** and **1a'** (Scheme 6). This type of tautomerism is known for bis(1-azaazulene-2-yl)amines.<sup>6</sup> We observed that red shift of absorption spectrum of **1a** by addition of hydrochloric acid to methanol solution of **1a**. This result also indicated that the existence of the two tautomeric forms. The details of tautomerism and p*K*<sub>a</sub> values are currently under investigation. The  $\epsilon$  value for ethyl ester-substituted **1b** at 440 nm, which is the absorption maximum in the visible region, is about 1.5 times larger than that for **1a** and **1c** (Figure 2a), and about 2 times larger than that for other 2-arylamino-1-azaazulenes (Figure 2b).



**Figure 2.** UV-Vis spectra of (a) **1a–1c** and **2d** and (b) **1a**, **2d**, **6**, **8**, **10**, and **12** in THF



**Scheme 6**

In summary, a series of 2-arylamino-1-azaazulenes were successfully synthesized in moderate to yields. X-Ray crystallographic analysis of **10** reveals a highly planar structure that straddles the azaazulene core and amino group including the hydrogen on the N atom. Moreover, the plane of the azaazulene core and the naphthalene ring is twisted. As per the crystal structure, **10** forms a dimer with two intermolecular N–H···N hydrogen bonds. The UV-Vis absorption spectra of the 2-arylamino-1-azaazulenes show a large  $\epsilon$  and redshift compared with the case of amino azulene **2d**, in the visible region. Additionally,  $\epsilon$  of the ethyl ester-substituted **1b** is larger than that for **1a**, **1c**, and other 2-arylamino-1-azaazulenes.

## EXPERIMENTAL

Unless otherwise indicated, all reactions were carried out under argon atmosphere. All solvents were distilled, dried over 3Å or 4Å molecular sieves, and degassed using nitrogen before use. Column chromatography was performed using Wako gel C-200 (spherical, neutral). Aniline, Cs<sub>2</sub>CO<sub>3</sub>, 18-crown-6, and BINAP were purchased from Wako Pure Chemical Industries. 1-Naphthylamine, 2-bromonaphthalene, and Xantphos were purchased from Tokyo Chemical Industry Co. Ltd. Pd<sub>2</sub>(dba)<sub>3</sub> was purchased from Strem Chemicals Inc. Pd(dppf)Cl<sub>2</sub> was purchased from Sigma-Aldrich. 1,4-Dioxane and THF were purchased from KANTO Chemicals.

All NMR spectra were recorded on a JEOL ECP-300 (300 MHz) or ECP-500 (500 MHz) spectrometer. <sup>1</sup>H NMR (300 or 500 MHz) and <sup>13</sup>C-<sup>1</sup>H NMR (75 or 126 MHz) spectra were recorded using tetramethylsilane as an internal standard. IR spectra were recorded with a JASCO FT-IR 6100 spectrometer, and UV-Vis-near infrared (NIR) spectra were recorded with a JASCO V670 spectrometer. Melting points were recorded using a Bibby Stuart Scientific SMP3 instrument; all melting points reported are uncorrected. HR-ESI mass spectra were recorded on a JEOL JMS-T100CS AccuTOF CS spectrometer. Elemental analyses were carried out using a PerkinElmer 2400II CHNS analyzer. Crystal data were collected using a Bruker AXS SMART APEX CCD X-ray diffractometer equipped with monochromatic Mo K $\alpha$  radiation (0.7107 Å). Empirical absorption corrections using equivalent reflections and Lorentzian polarization corrections were performed using the SADABS program.<sup>19</sup> All data were collected with SMART and Bruker SAINTPLUS (Version 6.45) software packages. The structures were solved using the SHELXA-97 program<sup>20</sup> and refined against  $F^2$  by using SHELEXL-97.<sup>21</sup> Deposition number CCDC-1500558 for compound No. **10**. Free copies of the data can be obtained via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> (or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK; Fax: +44 1223 336033; e-mail: [deposit@ccdc.cam.ac.uk](mailto:deposit@ccdc.cam.ac.uk)).

**2d**<sup>22</sup> and **5**<sup>23</sup> were prepared according to literature methods.

### Synthesis of 2a

Compound **2a** was synthesized as per the literature method for the synthesis of 2-bromo-3-methyl-1-azaazulene.<sup>24</sup>

A mixture of 1-azaazulenone<sup>25</sup> (0.22 g, 1.5 mmol) and phosphorus(V) oxybromide (1.5g, 5.2 mmol) was stirred for 30 min at 120 °C. Ice water was added to the reaction mixture, followed by neutralization with aqueous NaHCO<sub>3</sub> solution. The mixture was extracted with CHCl<sub>3</sub>, and the organic layer was washed with water and dried with MgSO<sub>4</sub>. After filtration and evaporation, the crude product was purified by silica gel column chromatography using CHCl<sub>3</sub> as the eluent. The first red fraction to elute from the column was collected. Recrystallization from hexane gave 2,3-dibromo-1-azaazulene **13** as red crystals (27%). And the second orange fraction to elute from the column was collected. Recrystallization from hexane gave **2a** as orange crystals (20%).

**2a**: mp 57.0-57.9 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.33 (s, 1H), 7.67 (t, *J* = 9.5 Hz, 1H), 7.77 (t, *J* = 9.5 Hz, 1H), 7.90 (t, *J* = 9.5 Hz, 1H), 8.45 (d, *J* = 9.5 Hz, 1H), 8.61 (d, *J* = 9.5 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, CDCl<sub>3</sub>): δ 115.8, 130.1, 130.5, 134.4, 135.6, 138.3, 147.1, 147.4, 156.8. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>9</sub>H<sub>7</sub><sup>79</sup>BrN: 207.9762 [M+H]<sup>+</sup>; found: 207.9761.

**13**: mp 143.9-144.7 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.75 (t, *J* = 9.5 Hz, 1H), 7.78 (t, *J* = 9.5 Hz, 1H), 7.95 (t, *J* = 9.5 Hz, 1H), 8.41 (d, *J* = 9.5 Hz, 1H), 8.56 (d, *J* = 9.5 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, CDCl<sub>3</sub>): δ 103.7, 130.5, 130.9, 134.8, 136.6, 139.4, 143.2, 148.4, 155.7. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>9</sub>H<sub>6</sub><sup>79</sup>Br<sup>81</sup>BrN: 287.8847 [M+H]<sup>+</sup>; found: 287.8847.

### Synthesis of 2b

A mixture of ethyl 1,2'-dihydro-2-oxo-1-azaazulene-3-carboxylate<sup>25</sup> (0.33 g, 1.4 mmol) and phosphorus(V) oxychloride (1.8 g, 11.8 mmol) was stirred for 2 h at 110 °C. Ice water was added to the reaction mixture, followed by neutralization with aqueous NaHCO<sub>3</sub> solution. The mixture was extracted with CHCl<sub>3</sub>, and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration and evaporation, the crude product was purified by silica gel column chromatography using CHCl<sub>3</sub> as the eluent. The first yellow fraction to elute from the column was collected. Recrystallization from hexane gave **2b** as yellow crystals (73%).

**2b**: mp 84.5-85.5 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 1.46 (t, *J* = 7.5 Hz, 3H), 4.46 (q, *J* = 7.5 Hz, 2H), 7.93 (t, *J* = 9.5 Hz, 1H), 7.96 (t, *J* = 9.5 Hz, 1H), 8.03 (t, *J* = 9.5 Hz, 1H), 8.66 (d, *J* = 9.5 Hz, 1H), 9.52 (d, *J* = 9.5 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, CDCl<sub>3</sub>): δ 14.3, 60.5, 111.7, 133.0, 133.2, 137.1, 137.5, 139.8, 147.0, 156.7, 158.9, 163.2. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>12</sub>H<sub>10</sub><sup>35</sup>CINNaO<sub>2</sub>: 258.0298 [M+Na]<sup>+</sup>; found: 258.0298. Anal. Calcd for C<sub>12</sub>H<sub>10</sub>NO<sub>2</sub>Cl: C, 61.16; H, 4.28; N, 5.94. Found: C, 61.00; H, 4.10; N, 5.93.

### Synthesis of **2c**

Compound **2c** was prepared by a slight modification of the literature method.<sup>25</sup>

A mixture of 3-cyano-1-azaazulenone<sup>25</sup> (1.20 g, 7.05 mmol) and phosphorus(V) oxychloride (8.20 g, 53.5 mmol) was stirred for 1 h at 115 °C. Ice water was added to the reaction mixture, followed by neutralization using aqueous NaHCO<sub>3</sub> solution. The mixture was extracted with CHCl<sub>3</sub>, and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration and evaporation, the crude product was purified by silica gel column chromatography using CHCl<sub>3</sub> as the eluent. The second yellow fraction to elute from the column was collected. Recrystallization from AcOEt gave **2c** as gold needle-like crystals (57%).

**2c**: mp 231.8-233.0 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 8.03 (t, *J* = 9.5 Hz, 1H), 8.08 (t, *J* = 9.5 Hz, 1H), 8.16 (t, *J* = 9.5 Hz, 1H), 8.70 (d, *J* = 9.5 Hz, 1H), 8.75 (d, *J* = 9.5 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, CDCl<sub>3</sub>): δ 96.0, 113.3, 133.6, 134.2, 135.2, 138.7, 140.9, 149.3, 156.8, 159.5. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>10</sub>H<sub>5</sub><sup>35</sup>ClN<sub>2</sub>Na: 211.0039 [M+Na]<sup>+</sup>; found: 211.0039. Anal. Calcd for C<sub>10</sub>H<sub>5</sub>N<sub>2</sub>Cl: C, 63.68; H, 2.67; N, 14.85. Found: C, 63.45; H, 2.46; N, 14.77.

### Synthesis of **2e**

Compound **2e** was prepared by a slight modification of the literature method.<sup>26</sup>

A mixture of 1-azaazulenone<sup>25</sup> (0.20 g, 1.4 mmol) and phosphorus(V) oxychloride (1.8 g, 11.8 mmol) was stirred for 2 h at 110 °C. Ice water was added to the reaction mixture, followed by neutralization with aqueous NaHCO<sub>3</sub> solution. The mixture was extracted with CHCl<sub>3</sub>, and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>. After filtration and evaporation, the crude product was purified by silica gel column chromatography using CHCl<sub>3</sub> as the eluent. The first yellow fraction to elute from the column gave **2e** as orange crystals (68%).

**2e**: mp 72.2-73.5 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 7.20 (s, 1H), 7.65 (t, *J* = 9.5 Hz, 1H), 7.75 (t, *J* = 9.5 Hz, 1H), 7.84 (t, *J* = 9.5 Hz, 1H), 8.40 (d, *J* = 9.5 Hz, 1H), 8.55 (d, *J* = 9.5 Hz, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, CDCl<sub>3</sub>): δ 112.1, 130.1, 130.5, 134.6, 135.7, 138.0, 147.2, 156.2, 157.7. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>9</sub>H<sub>7</sub><sup>35</sup>ClN: 164.0267 [M+H]<sup>+</sup>; found: 164.0262. Anal. Calcd for C<sub>9</sub>H<sub>6</sub>NCl: C, 66.07; H, 3.70; N, 8.56. Found: C, 65.93; H, 3.53; N, 8.51.

### Synthesis of **3**

Compound **3** was prepared by a slight modification of the reported method.<sup>27</sup>

To a solution of KOH (4.0 g), EtOH (5.0 mL), and water (1.0 mL), diethyl 2-aminoazulene-1,3-dicarboxylate<sup>28,29</sup> (0.75 g, 2.65 mmol) was added. The reaction mixture was refluxed for 30 min with continuous stirring. After cooling to room temperature, the reaction mixture was neutralized using hydrochloric acid to obtain a yellow precipitate. The yellow solid was filtered out,

added to pyridine (5 mL), and refluxed for 30 min. After removal of pyridine under reduced pressure, the crude product was purified by silica gel column chromatography using  $\text{CHCl}_3$  as the eluent. The second red fraction to elute from the column was collected and recrystallized from hexane to give **3** as red needle-like crystals (82%). Compound **3** can also be synthesized by another method.<sup>30</sup>

**3**: mp 92.2-93.4 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  4.56 (s, 2H), 6.66 (s, 2H), 7.15 (t,  $J = 9.5$  Hz, 2H), 7.28 (t,  $J = 9.5$  Hz, 1H), 7.91 (d,  $J = 9.5$  Hz, 2H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  102.8, 124.3, 128.5, 129.8, 141.8, 156.5. HR-MS (HR-ESI-TOF)  $m/z$  calc. for  $\text{C}_{10}\text{H}_{10}\text{N}$ : 144.0813  $[\text{M}+\text{H}]^+$ ; found: 144.0808. Anal. Calcd for  $\text{C}_{10}\text{H}_9\text{N}$ : C, 83.88; H, 6.34; N, 9.78. Found: C, 83.69; H, 6.35; N, 9.88.

### General procedure for the synthesis of 2-arylamino-1-azaazulenes

A mixture of the aryl halide (0.20 mmol), arylamine (0.28 mmol), Pd catalyst (6 mol%), phosphine ligand (4 equiv to Pd),  $\text{Cs}_2\text{CO}_3$  (0.24 mmol), and 18-crown-6 (10 mol%) was refluxed in dry THF or 1,4-dioxane (6 mL) for 24 h in a sealed tube. Then, 20 mL of brine was added and the resulting mixture was extracted with  $\text{CHCl}_3$ . The combined organic layers were dried over anhydrous  $\text{Na}_2\text{SO}_4$ . After filtration and evaporation, the crude product was purified by silica gel column chromatography using  $\text{CHCl}_3$  as the eluent. The product was recrystallized from AcOEt.

### Synthesis of **1a**

The first red purple fraction to elute from the column gave **4** as a red purple powder (18%). The sixth dark red fraction to elute gave **1a** as a dark red powder (24%).

**1a**: mp > 212.8 °C (decomp.).  $^1\text{H}$  NMR (500 MHz,  $\text{DMSO}-d_6$ )  $\delta$  7.11 (s, 1H), 7.21 (t,  $J = 9.5$  Hz, 2H), 7.40 (t,  $J = 9.5$  Hz, 1H), 7.47-7.52 (m, 2H), 7.58-7.63 (m, 1H), 7.65 (s, 2H), 8.13 (d,  $J = 9.5$  Hz, 1H), 8.15 (d,  $J = 9.5$  Hz, 2H), 8.19-8.23 (m, 1H), 11.07 (s, 1H).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{DMSO}-d_6$ ):  $\delta$  101.5, 106.1, 124.2, 127.7, 129.2, 129.4, 130.3, 131.1, 131.5, 131.9, 140.2, 146.9, 149.2, 157.5, 163.6. HRMS (HR-ESI-TOF)  $m/z$  calcd. for  $\text{C}_{19}\text{H}_{15}\text{N}_2$ : 271.1235  $[\text{M}+\text{H}]^+$ , found: 271.1241. Anal. Calcd for  $\text{C}_{19}\text{H}_{14}\text{N}_2$ : C, 84.42; H, 5.22; N, 10.36. Found: C, 84.17; H, 5.20; N, 10.20.

**4**: mp 172.9-174.6 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.69 (s, 1H), 7.16 (t,  $J = 9.0$  Hz, 1H), 7.23 (t,  $J = 9.0$  Hz, 1H), 7.29 (t,  $J = 9.0$  Hz, 1H), 7.51 (t,  $J = 9.5$  Hz, 1H), 7.57 (t,  $J = 9.5$  Hz, 1H), 7.63 (t,  $J = 9.5$  Hz, 1H), 7.82 (d,  $J = 9.5$  Hz, 1H), 7.87 (s, 1H), 8.35 (d,  $J = 9.0$  Hz, 1H), 8.41 (d,  $J = 9.5$  Hz, 1H), 8.57 (d,  $J = 9.0$  Hz, 1H) (NH proton was not appeared).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  105.6, 107.9, 112.4, 126.9, 127.6, 127.8, 129.4, 130.4, 130.7, 131.5, 131.7, 132.4, 134.1, 141.6, 146.0, 147.5, 158.5, 160.4, 165.3. HR-MS (HR-ESI-TOF)  $m/z$  calc. for  $\text{C}_{19}\text{H}_{15}\text{N}_2$ : 271.1235  $[\text{M}+\text{H}]^+$ ; found: 271.1235.

### Synthesis of **1b**

A reaction mixture was refluxed in THF. The first red fraction to elute from the column gave **1b** as a red powder (70%).

**1b**: mp 151.5-153.5 °C. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 1.55 (t, *J* = 6.5 Hz, 3H), 4.55 (q, *J* = 6.5 Hz, 2H), 7.21 (t, *J* = 9.5 Hz, 2H), 7.44 (t, *J* = 9.5 Hz, 1H), 7.64 (t, *J* = 9.5 Hz, 1H), 7.75 (t, *J* = 9.5 Hz, 1H), 7.80-7.84 (m, 3H), 8.22 (d, *J* = 9.5 Hz, 2H), 8.64 (d, *J* = 9.5 Hz, 1H), 8.99 (d, *J* = 9.5 Hz, 1H), 10.54 (s, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, CDCl<sub>3</sub>): δ 14.7, 60.7, 99.3, 107.4, 124.4, 131.4, 132.3, 132.9, 133.2, 133.7, 133.9, 134.1, 140.6, 146.8, 147.2, 160.6, 164.2, 166.6. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>22</sub>H<sub>19</sub>N<sub>2</sub>O<sub>2</sub>: 343.1447 [M+H]<sup>+</sup>; found: 343.1448. IR (KBr, ν/cm<sup>-1</sup>): 3469 (N-H), 3272 (Aryl), 2979 (C-H), 1650 (C=O). Anal. Calcd for C<sub>22</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>: C, 77.17; H, 5.30; N, 8.18. Found: C, 77.01; H, 5.17; N, 8.12.

### Synthesis of **1c**

A reaction mixture was refluxed in 1,4-dioxane. The third yellow brown fraction to elute from the column gave **1c** as a yellow brown powder (30%).

**1c**: mp > 278.3 °C (decomp.). <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 7.24 (t, *J* = 9.5 Hz, 2H), 7.47 (t, *J* = 9.5 Hz, 1H), 7.84 (t, *J* = 9.5 Hz, 1H), 7.87 (s, 2H), 7.92 (t, *J* = 9.5 Hz, 1H), 7.98 (t, *J* = 9.5 Hz, 1H), 8.23 (d, *J* = 9.5 Hz, 2H), 8.30 (d, *J* = 9.5 Hz, 1H), 8.44 (d, *J* = 9.5 Hz, 1H), 11.35 (s, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, DMSO-*d*<sub>6</sub>): δ 81.4, 107.5, 114.9, 124.1, 128.6, 131.6, 132.4, 133.0, 133.7, 134.7, 134.8, 139.6, 148.1, 150.2, 159.0, 162.7. HR-MS (HR-ESI-TOF) *m/z* calc. for C<sub>20</sub>H<sub>13</sub>N<sub>3</sub>Na: 318.1007 [M+Na]<sup>+</sup>; found: 318.1005. IR (KBr, ν/cm<sup>-1</sup>): 3469 (N-H), 3272, 3138 (Aryl), 3023 (C-H), 2214 (C≡N). Anal. Calcd for C<sub>20</sub>H<sub>13</sub>N<sub>2</sub>: C, 81.34; H, 4.44; N, 14.23. Found: C, 81.09; H, 4.40; N, 14.16.

### Synthesis of **6**

The fifth dark red fraction to elute from the column gave **6** as a dark red powder (26%).

**6**: mp > 193.4 °C (decomp.). <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>): δ 7.10 (s, 1H), 7.20 (d, *J* = 4.0 Hz, 2H), 7.50 (t, *J* = 4.0 Hz, 1H), 7.52 (t, *J* = 9.0 Hz, 1H), 7.56 (t, *J* = 9.0 Hz, 1H), 7.64 (t, *J* = 9.0 Hz, 1H), 7.99 (d, *J* = 9.0 Hz, 2H), 8.18 (d, *J* = 9.0 Hz, 1H), 8.26 (d, *J* = 9.0 Hz, 1H), 8.28 (d, *J* = 9.0 Hz, 2H), 10.74 (s, 1H). <sup>13</sup>C{<sup>1</sup>H} NMR (126 MHz, DMSO-*d*<sub>6</sub>): δ 103.0, 112.9, 118.2, 128.7, 129.6, 130.1, 130.4, 131.6, 132.5, 135.4, 136.4, 146.1, 150.4, 158.8, 163.5. HR-MS (HR-APCI-TOF) *m/z* calc. for C<sub>19</sub>H<sub>15</sub>N<sub>2</sub>: 271.1235 [M+H]<sup>+</sup>; found: 271.1238. Anal. Calcd for C<sub>19</sub>H<sub>14</sub>N<sub>2</sub>: C, 84.42; H, 5.22; N, 10.36. Found: C, 84.37; H, 5.09; N, 10.31.

### Synthesis of **8**

The fourth yellow fraction to elute from the column gave **8**<sup>12</sup> as a yellow powder (13%).

**8:** mp 148.8-151.1 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.94 (s, 1H), 7.12 (t,  $J = 7.2$  Hz, 1H), 7.34-7.40 (m, 4H), 7.47-7.52 (m, 3H), 7.99 (t,  $J = 9.5$  Hz, 2H) (NH proton was not appeared).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  98.5, 119.8, 123.3, 126.7, 128.4, 129.3, 129.7, 130.6, 130.7, 140.3, 148.4, 157.6, 165.2. HR-MS (HR-ESI-TOF)  $m/z$  calc. for  $\text{C}_{15}\text{H}_{13}\text{N}_2$ : 221.1079  $[\text{M}+\text{H}]^+$ ; found: 221.1076. Anal. Calcd for  $\text{C}_{15}\text{H}_{12}\text{N}_2$ : C, 81.79; H, 5.49; N, 12.72. Found: C, 81.47; H, 5.43; N, 12.57.

### Synthesis of 10

The fourth yellow fraction to elute from the column gave **10** as a yellow powder (28%).

**10:** mp 195.9-196.9 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  6.70 (s, 1H), 7.36-7.39 (m, 2H), 7.45-7.50 (m, 1H), 7.51-7.56 (m, 3H), 7.76 (d,  $J = 8.0$  Hz, 1H), 7.82 (d,  $J = 8.0$  Hz, 1H), 7.90-7.94 (m, 3H), 8.18 (d,  $J = 8.0$  Hz, 1H) (NH proton was not appeared).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  98.5, 119.0, 122.0, 125.5, 125.7, 126.3, 126.4, 126.5, 128.2, 128.4, 128.5, 129.8, 130.7, 130.8, 134.5, 135.8, 148.3, 157.3, 166.5. HR-MS (HR-ESI-TOF)  $m/z$  calc. for  $\text{C}_{19}\text{H}_{15}\text{N}_2$ : 271.1235  $[\text{M}+\text{H}]^+$ ; found: 271.1221. Anal. Calcd for  $\text{C}_{19}\text{H}_{14}\text{N}_2$ : C, 84.42; H, 5.22; N, 10.36. Found: C, 84.14; H, 5.19; N, 10.29.

### Synthesis of 12

The fourth yellow fraction to elute from the column gave **12** as a yellow powder (51%).

**12:** mp 200.7-202.1 °C.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  7.02 (s, 1H), 7.39-7.46 (m, 3H), 7.47-7.50 (m, 1H), 7.53-7.56 (m, 2H), 7.81 (t,  $J = 7.0$  Hz, 2H), 7.86 (d,  $J = 8.5$  Hz, 1H), 7.95 (d,  $J = 2.0$  Hz, 1H), 8.03 (dd,  $J = 2.0, 8.5$  Hz, 1H), 8.06 (d,  $J = 8.5$  Hz, 1H) (NH proton was not appeared).  $^{13}\text{C}\{^1\text{H}\}$  NMR (126 MHz,  $\text{CDCl}_3$ ):  $\delta$  98.6, 115.4, 120.5, 124.7, 126.7, 126.7, 127.1, 127.7, 129.1, 129.3, 130.2, 130.3, 131.0, 131.4, 134.2, 137.6, 148.4, 156.6, 164.2. HR-MS (HR-ESI-TOF)  $m/z$  calc. for  $\text{C}_{19}\text{H}_{15}\text{N}_2$ : 271.1235  $[\text{M}+\text{H}]^+$ ; found: 271.1234.

### ACKNOWLEDGEMENTS

This work was supported by JSPS KAKENHI Grant Number JP24102008, JP16K17951.

### REFERENCES

1. K. Elblidi, A. Boucetta, B. Benali, A. Kadiri, C. Cazeau-Dubroca, G. Nouchi, and M. Pesquer, *J. Mol. Struct. (Theochem)*, 1995, **343**, 57.
2. T. Nishiwaki and N. Abe, *Heterocycles*, 1981, **15**, 547.
3. N. Abe, *Trends Heterocycl. Chem.*, 2001, **7**, 25.
4. N. Abe, *Recent Res. Devel. Org. Bioorg. Chem.*, 2001, **4**, 17.
5. N. Abe and T. Gunji, *Heterocycles*, 2010, **82**, 201.

6. E. Yoshioka, K. Koizumi, K. Nakashima, H. Fujii, T. Murafuji, T. Gunji, and N. Abe, [\*Heterocycles\*, 2012, \*\*85\*\*, 1683.](#)
7. M. Hyoudou, H. Nakagawa, T. Gunji, Y. Ito, Y. Kawai, R. Ikeda, T. Konakahara, and N. Abe, [\*Heterocycles\*, 2012, \*\*86\*\*, 233.](#)
8. K. Koizumi, K. Shimabara, A. Takemoto, S. Yamazaki, N. Yamauchi, H. Fujii, M. Kurosawa, T. Konakahara, and N. Abe, [\*Heterocycles\*, 2009, \*\*79\*\*, 319.](#)
9. N. Abe, N. Ishikawa, T. Hayashi, and Y. Miura, [\*Bull. Chem. Soc. Jpn.\*, 1990, \*\*63\*\*, 1617.](#)
10. E. Yoshioka, K. Koizumi, S. Yamazaki, H. Fujii, and N. Abe, [\*Heterocycles\*, 2009, \*\*78\*\*, 3065.](#)
11. M. Oda, D. Miyawaki, N. Matsumoto, and S. Kuroda, [\*Heterocycles\*, 2011, \*\*83\*\*, 547.](#)
12. H. Fujii, S. Oka, I. Nakamura, Y. Kawai, R. Ikeda, T. Konakahara, and N. Abe, [\*Heterocycles\*, 2015, \*\*90\*\*, 715.](#)
13. S. Hillebrand, J. Bruckmann, C. Krüger, and M. W. Haenel, [\*Tetrahedron Lett.\*, 1995, \*\*36\*\*, 75.](#)
14. K. Takeda, A. Nakajima, M. Takeda, and E. Yoshii, [\*Org. Synth.\*, 1999, \*\*76\*\*, 6.](#)
15. A. Abbotto, L. Beverina, S. Bradamante, A. Facchetti, C. Klein, G. A. Pagani, M. Redi-Abshiro, and R. Wortmann, [\*Chem. Eur. J.\*, 2003, \*\*9\*\*, 1991.](#)
16. M. Nishio, [\*Phys. Chem. Chem. Phys.\*, 2011, \*\*13\*\*, 13873.](#)
17. M. Nishio, Y. Umezawa, J. Fantini, M. S. Weiss, and P. Chakrabarti, [\*Phys. Chem. Chem. Phys.\*, 2014, \*\*16\*\*, 12648.](#)
18. O. Takahashi, Y. Kohno, and M. Nishio, [\*Chem. Rev.\*, 2010, \*\*110\*\*, 6049.](#)
19. G. M. Sheldrick, SADABS, Program for Siemens Area Detector Absorption Correction, University of Göttingen, Germany, 1996.
20. G. M. Sheldrick, SHELXS-97, Program for Crystal Structure Solution, University of Göttingen, Germany, 1997.
21. G. M. Sheldrick, SHELXL-97, Program for Crystal Structure Refinement, University of Göttingen, Germany, 1997.
22. K. Fujimori, N. Hirako, and K. Yamane, [\*Bull. Chem. Soc. Jpn.\*, 1983, \*\*56\*\*, 1247.](#)
23. R. N. McDonald, J. M. Richmond, J. R. Curtis, H. E. Petty, and T. L. Hoskins, [\*J. Org. Chem.\*, 1976, \*\*41\*\*, 1811.](#)
24. K. Koizumi, C. Miyake, T. Ariyoshi, K. Umeda, N. Yamauchi, S. Tagashira, Y. Murakami, H. Fujii, and N. Abe, [\*Heterocycles\*, 2007, \*\*73\*\*, 325.](#)
25. M. Nagahara, J. Nakano, M. Mimura, T. Nakamura, and K. Uchida, [\*Chem. Pharm. Bull.\*, 1994, \*\*42\*\*, 2491.](#)
26. T. Nozoe, S. Seto, S. Matsumura, and T. Nakazawa, *Chem. Ind. (London)*, 1954, 1357.
27. T. Nozoe, S. Seto, K. Takase, S. Matsumura, and T. Nakazawa, [\*Nippon Kagaku Zasshi\*, 1965, \*\*86\*\*,](#)

[346](#).

28. T. Nozoe, S. Seto, S. Matsumura, and Y. Murase, [Bull. Chem. Soc. Jpn., 1962, 35, 1179](#).
29. T. Nozoe, K. Takase, M. Kato, and T. Nogi, [Tetrahedron, 1971, 27, 6023](#).
30. Y. Sugihara, T. Murafuji, M. Fujinaga, K. Suetake, K. Gyoji, and K. Kurotobi, [Synthesis, 2008, 3745](#).