

HETEROCYCLES, Vol. 97, No. 2, 2018, pp. 1219 - 1225. © 2018 The Japan Institute of Heterocyclic Chemistry
Received, 16th February, 2018, Accepted, 20th March, 2018, Published online, 29th March, 2018
DOI: 10.3987/COM-18-S(T)73

ISOLATION OF AAPTIC ACID FROM THE MARINE SPONGE *AAPTOS LOBATA* AND INHIBITORY EFFECT OF AAPTAMINES ON RANKL-INDUCED FORMATION OF MULTINUCLEAR OSTEOCLASTS

Arina Fukumoto,¹ Yuki Hitora,¹ Aika Kai,¹ Hikaru Kato,¹ Esther D. Angkouw,² Remy E. P. Mangindaan,² Nicole J. de Voogd,³ and Sachiko Tsukamoto^{1*}

¹Graduate School of Pharmaceutical Sciences, Kumamoto University, Kumamoto 862-0973, Japan. ²Faculty of Fisheries and Marine Science, Sam Ratulangi University, Kampus Bahu, Manado 95115, Indonesia. ³Naturalis Biodiversity Center, P.O. Box 9517, 2300 RA Leiden, The Netherlands. E-mail: sachiko@kumamoto-u.ac.jp

Abstract – A new aaptamine congener, aaptic acid (**1**), was isolated from the marine sponge *Aaptos lobata* collected in Indonesia, together with aaptamine (**2**), demethyl(oxy)aaptamine (**3**), and iso-aaptamine (**4**). The chemical structure of **1** was assigned on the basis of NMR data. The isolated compounds were tested for inhibitory activities against osteoclastogenesis in RAW264 cells. At 5 μ M concentration, **2-4** inhibited RANKL-induced multinuclear osteoclast formation by 25%, 54%, and 17%, respectively. In contrast, **1** did not inhibit osteoclastogenesis even at 50 μ M.

Aaptamine (**2**) is a marine alkaloid bearing a benzo[*de*][1,6]-naphthyridine scaffold and was isolated from the sponge *Aaptos aaptos* collected at Manza beach in Okinawa in 1982.¹ Since its isolation, more than thirty congeners have been isolated from sponges of the genera *Aaptos*,² *Xestospongia*,³ *Suberites*,⁴ *Hymeniacidon*,⁵ and *Luffariella*.^{6,7} Interestingly, the congeners have been reported to show broad spectra of biological activities, such as cytotoxic,^{2,3,8} antimicrobial,^{2,3} antiviral,⁹ antifouling,¹⁰ antioxidant,¹¹ and enzymatic inhibitory activities.¹² The cytotoxic effect of **2** was well investigated and found to activate the cyclin dependent kinase inhibitor p21, which leads to inhibition of cell cycle progression and induction of apoptosis.¹³ Further, proteome analysis indicated that **2** exhibited cytotoxic activity by targeting *myc* and p53 and promoted hypusination of eukaryotic initiation factor 5A-1 (eIF5A).¹⁴ In our continuing search

for new bioactive compounds from marine invertebrates, aaptic acid (**1**), a new aaptamine congener, was isolated from the sponge *A. lobata* collected in Indonesia, together with **2**,¹ demethyl(oxy)aaptamine (**3**),² and iso-aaptamine (**4**).⁴ In our in-house screening, **2–4** inhibited RANKL-induced multinuclear osteoclast formation but **1** showed no inhibition.




Figure 1. Chemical structures of **1–4**


The sponge was collected in Indonesia and extracted with EtOH and then MeOH. The extract was partitioned between EtOAc and water, and the aqueous layer was extracted with *n*-BuOH. Purification of *n*-BuOH fraction afforded **1–4**.

Aaptic acid (**1**) was obtained as dark brown amorphous solid. The molecular formula of C₁₆H₁₆N₂O₄ was determined by HRESIMS. ¹H NMR spectrum of **1** showed two sets of mutually coupled protons [δ_{H} 8.09 (d, $J = 5.6$ Hz, H-2)/ δ_{H} 6.23 (d, $J = 5.6$ Hz, H-3) and δ_{H} 7.03 (d, $J = 7.2$ Hz, H-5)/ δ_{H} 6.36 (d, $J = 7.2$ Hz, H-6)], a singlet proton [δ_{H} 6.75 (H-7)], mutually coupled methylene protons [δ_{H} 2.56 (t, $J = 6.2$ Hz, H-2')/ δ_{H} 3.95 (t, $J = 6.2$ Hz, H-3')], and two singlet methyls [δ_{H} 3.85 (8-OMe) and 3.73 (9-OMe)] (Table 1). ¹³C NMR spectrum and HSQC spectrum of **1** revealed the presence of one carbonyl carbon [δ_{C} 172.3 (C-1')], five methine carbons [δ_{C} 149.2 (C-2), 95.5 (C-3), 134.3 (C-5), 109.3 (C-6), and 100.1 (C-7)], six quaternary carbons [δ_{C} 147.0 (C-3a), 130.1 (C-6a), 154.1 (C-8), 135.5 (C-9), 141.8 (C-9a), and 117.8 (C-9b)], two methylene carbons [δ_{C} 31.9 (C-2') and 47.6 (C-3')], and two methoxy carbons [δ_{C} 56.0 (8-OMe) and 59.5 (9-OMe)] (Table 1).

Table 1. ^1H (600 MHz) and ^{13}C (150 MHz) NMR data for **1** ($\text{DMSO-}d_6$)

No.	δ_{C}	δ_{H} , mult (J in Hz)	HMBC
2	149.2, CH	8.09, d (5.6)	3, 3a, 9a
3	95.5, CH	6.23, d (5.6)	9b
3a	147.0, C		
5	134.3, CH	7.03, d (7.2)	6, 3a, 6a
6	109.3, CH	6.36, d (7.2)	5, 7, 9b
6a	130.1, C		
7	100.1, CH	6.75, s	6, 8, 9, 9b
8	154.1, C		
9	135.5, C		
9a	141.8, C		
9b	117.8, C		
8-OMe	56.0, CH_3	3.85, s	8
9-OMe	59.5, CH_3	3.73, s	9
1'	172.3, C		
2'	31.9, CH_2	2.56, t (6.2)	1', 3'
3'	47.6, CH_2	3.95, t (6.2)	3a, 5, 1', 2'

Key HMBC correlations, H-2/C-3a and C-9a, H-3/C-9b, H-5/C-3a and C-6a, H-6/C-7 and C-9b, H-7/C-8, C-9, and C-9b, 8-OMe/C-8, and 9-OMe/C-9, established the aaptamine nucleus. Additional HMBC correlations from H-2' and H-3' to C-1' and from H-3' to C-3a and C-5 assigned the substitution of propionic acid at the N-4 position, and therefore the chemical structure of **1** was determined (Figure 2).

**Figure 2.** COSY and key HMBC correlations of **1**

Compounds **1–4** were evaluated for inhibitory effects on RANKL-induced osteoclastogenesis in RAW264 cells. RANKL induces the osteoclastogenesis of RAW264 cells to form TRAP-positive cells. Subsequently, cell-cell fusion is initiated and then giant multinuclear osteoclasts are formed. In our *in vitro* bioassay, four days after the treatment with RANKL (50 ng/mL) and each compound (5 μ M), cells were fixed and stained with TRAP staining solution (Figure 3a), and the numbers of multinuclear osteoclasts were counted (Figure 3b).^{15,16} Compared with the control containing RANKL, **3** inhibited the formation of multinuclear osteoclasts by 54% without cytotoxicity, and **2** and **4** moderately inhibited it by 25% and 17%, respectively (Figure 3b). In contrast, **1** did not inhibit the osteoclastogenesis even at a concentration of 50 μ M.




Figure 3. Inhibitory effects of **1–4** on RANKL-induced multinuclear osteoclast formation in RAW264 cells. Cells were incubated with RANKL (50 ng/mL) in the presence or absence of **1–4** (5 μ M) for four days. (a) The cells were stained with TRAP-staining solution. (b) Numbers of TRAP-positive multinuclear osteoclasts (nuclei \geq 3) stained in red were counted. Quercetin (25 μ g/mL, 83 μ M) was used as a positive control. The bars represent means \pm S.D. *P < 0.05 and **P < 0.001.

Aaptic acid (**1**), substituted with a propionic acid at N-4, and three known aaptamines (**2–4**) were isolated from the marine sponge *A. lobata* collected in Indonesia. Although more than thirty aaptamine congeners have been isolated from marine sponges so far, **1** is the first compound substituted with a carboxylic acid at the aaptamine nucleus. Aaptamines **2–4** inhibited the formation of RANKL-induced multinuclear

osteoclasts at 5 μM . However, **1** did not exhibit the inhibition even at 50 μM . Introduction of an acyl group and methylation at the N-4 position have been reported to decrease the cytotoxicity, which clearly indicates that the nitrogen atom at N-4 of aaptamines may play a key role in their biological activities.^{8,17,18} The mechanism of the inhibitory effect of aaptamines on osteoclastogenesis is currently under investigation in our laboratory.

EXPERIMENTAL

General Experimental Procedures

UV spectra were measured on a JASCO V-550 spectrophotometer in MeOH. IR spectra were recorded on a Perkin Elmer Frontier FT-IR spectrophotometer. 1D and 2D NMR spectra were recorded on a Bruker Avance III 600 NMR spectrometer. Chemical shifts were referenced to the residual solvent peaks (δ_{H} 2.49 and δ_{C} 39.5 for DMSO- d_6). HRESIMS spectra were measured on a Bruker impact-II mass spectrometer. The preparative HPLC system comprised a Waters 515 HPLC pump, Waters 2489 UV/visible detector. Microscopic images were taken by Keyence BIOREVO BZ-9000.

Animal Material

The marine sponge was collected at a depth of 10 m in North Sulawesi, Indonesia, in December 2010, and immediately soaked in EtOH. The sponge was identified as *Aaptos lobata* by Dr. N. J. de Voogd. A voucher specimen (RMNH POR. 11757) has been deposited in the Naturalis Biodiversity Center, The Netherlands.

Extraction and Isolation

The sponge (562 g wet weight) was extracted with EtOH and then MeOH. After evaporation, the residual aqueous solution was extracted with EtOAc and then *n*-BuOH. The *n*-BuOH fraction (14 g) was subjected to silica gel column chromatography and eluted with a stepwise gradient of $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (4:1 and 3:1), $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{H}_2\text{O}$ (6:4:1) and MeOH to yield 8 fractions (Frs. 1–8). Fr. 2 that eluted with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (4:1) was identified to be **2** (3.2 g). Fr. 1 was subjected to size exclusion HPLC (Asahipak GS-310P, Asahi Chemical Industry Co., Ltd., 21.5 \times 500 mm) with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (4:1) followed by silica gel HPLC (Inertsil Diol, GL Sciences Inc., 20 \times 250 mm) with *n*-hexane/ $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (40:19:1) to afford **3** (1.0 mg). Fr. 4 that eluted with $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (3:1) was purified by size exclusion HPLC (Asahipak GS-310P, 21.5 \times 500 mm) to afford **4** (8.4 mg). Frs. 5 and 6 (1.4 g) were combined and subjected to ODS column chromatography with a stepwise gradient of MeOH/ H_2O (2:3 and 4:1) and $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{H}_2\text{O}$ (6:4:1) to yield 6 fractions. The second fraction (197.8 mg) that eluted with MeOH/ H_2O (2:3) was purified by size exclusion HPLC (Asahipak GS-310P, 21.5 \times 500 mm) with MeOH/ H_2O (1:4) to yield 13 fractions. The third fraction was further purified by silica gel column chromatography with a stepwise gradient of $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (17:3, 5:1, 4:1 and 7:3) and

CH₂Cl₂/MeOH/H₂O (6:4:1) to afford eight fractions. The eighth fraction that eluted with CH₂Cl₂/MeOH/H₂O (6:4:1) yielded **1** (4.7 mg).

Aaptic Acid (1): dark brown amorphous solid; UV (MeOH) λ_{\max} (log ϵ) 218 (4.21), 240 (4.25), 258 (4.19), 268 (4.15), 278 (4.13) nm; IR (film) ν_{\max} 3364, 3083, 2924, 2853, 1651, 1598, 1569, 1459, 1432, 1381, 1350, 1321, 1250, 1196, 1162, 1111, 1093, 1058 cm⁻¹; HRESIMS m/z 301.1182 [M + H]⁺ (calcd for C₁₆H₁₇N₂O₄, 301.1183); ¹H and ¹³C NMR data (DMSO-*d*₆), see Table 1.

Formation of RANKL-Induced Multinuclear Osteoclasts in RAW264 Cells

This experiment was performed as previously described.^{15,16}

ACKNOWLEDGEMENTS

This work was supported by JSPS KAKENHI Grant Numbers JP22406001 (S.T.) and JP17H03994 (S.T.) of Japan.

REFERENCES

1. H. Nakamura, J. Kobayashi, Y. Ohizumi, and Y. Hirata, *Tetrahedron Lett.*, 1982, **23**, 5555.
2. H. Nakamura, J. Kobayashi, Y. Ohizumi, and Y. Hirata, *J. Chem. Soc., Perkin Trans. 1*, 1987, 173.
3. L. Calcul, A. Longeon, A. Al-Mourabit, M. Guyot, and M. L. Bourguet-Kondracki, *Tetrahedron*, 2003, **59**, 6539.
4. S. A. Fedoreev, N. G. Prokofeva, V. A. Denisenko, and N. M. Rebachuk, *Pharm. Chem. J.*, 1988, **22**, 615.
5. G. R. Pettit, H. Hoffmann, J. McNulty, K. C. Higgs, A. Murphy, D. J. Molloy, D. L. Herald, M. D. Williams, R. K. Pettit, D. L. Doubek, J. N. A. Hooper, L. Albright, J. M. Schmidt, J. C. Chapuis, and L. P. Tackett, *J. Nat. Prod.*, 2004, **67**, 506.
6. S. K. Park, S. S. Kim, J. D. Park, J. S. Hong, and I. K. Kim, *J. Korean Chem. Soc.*, 1995, **39**, 559.
7. E. L. Larghi, M. L. Bohn, and T. S. Kaufman, *Tetrahedron*, 2009, **65**, 4257.
8. Y. C. Shen, T. T. Lin, J. H. Sheu, and C. Y. Duh, *J. Nat. Prod.*, 1999, **62**, 1264.
9. A. F. Coutinho, B. Chanas, T. M. L. Souza, I. C. P. P. Frugulhetti, and R. A. Epifanio, *Heterocycles*, 2002, **57**, 1265.
10. J. A. Diers, J. J. Bowling, S. O. Duke, S. Wahyuono, M. Kelly, and M. T. Haman, *Mar. Biotechnol. (NY)*, 2006, **8**, 366.
11. S. Takamatsu, T. W. Hodges, I. Rajbhandari, W. H. Gerwick, M. T. Hamann, and D. G. Nagle, *J. Nat. Prod.*, 2003, **66**, 605.
12. S. Tsukamoto, R. Yamanokuchi, M. Yoshitomi, K. Sato, T. Ikeda, H. Rotinsulu, R. E. P.

- Mangindaan, N. J. de Voogd, R. W. M. van Soest, and H. Yokosawa, [*Bioorg. Med. Chem. Lett.*, 2010, **20**, 3341.](#)
13. S. Aoki, D. X. Kong, H. Suna, Y. Sowa, T. Sakai, A. Setiawan, and M. Kobayashi, [*Biochem. Biophys. Res. Commun.*, 2006, **342**, 101.](#)
 14. S. A. Dyshlovoy, S. Venz, L. K. Shubina, S. N. Fedorov, R. Walther, C. Jacobsen, V. A. Stonik, C. Bokemeyer, S. Balabanov, and F. Honecker, [*J. Proteomics*, 2014, **96**, 223.](#)
 15. S. Tsukamoto, T. Takeuchi, T. Kawabata, H. Kato, M. Yamakuma, K. Matsuo, A. H. El-Desoky, F. Losung, R. E. P. Mangindaan, N. J. de Voogd, Y. Arata, and H. Yokosawa, [*Bioorg. Med. Chem. Lett.*, 2014, **24**, 5315.](#)
 16. A. H. El-Desoky, H. Kato, E. D. Angkouw, R. E. P. Mangindaan, N. J. de Voogd, and S. Tsukamoto, [*J. Nat. Prod.*, 2016, **79**, 1922.](#)
 17. G. R. Pettit, H. Hoffmann, D. L. Herald, J. McNulty, A. Murphy, K. C. Higgs, E. Hamel, N. E. Lewin, L. V. Pearce, P. M. Blumberg, R. K. Pettit, and J. C. Knight, [*J. Org. Chem.*, 2004, **69**, 2251.](#)
 18. G. R. Pettit, H. Hoffmann, D. L. Herald, P. M. Blumberg, E. Hamel, J. M. Schmidt, Y. Chang, R. K. Pettit, N. E. Lewin, and L. V. Pearce, [*J. Med. Chem.*, 2004, **47**, 1775.](#)