

HETEROCYCLES, Vol. 99, No. 1, 2019, pp. 583 - 595. © 2019 The Japan Institute of Heterocyclic Chemistry
Received, 28th March, 2018, Accepted, 18th April, 2018, Published online, 20th April, 2018
DOI: 10.3987/COM-18-S(F)1

A CONVENIENT SYNTHESIS OF 2-(ALKYL(OR ARYL)SULFANYL)-2H-1,4-BENZOTHIAZIN-3(4H)-ONE DERIVATIVES

Kazuhiro Kobayashi,* Kazuki Kawano, and Keita Yamashita


Applied Chemistry Field, Chemistry and Biotechnology Course, Department of Engineering, Graduate School of Sustainability Science, Tottori University, 4-101 Koyama-minami, Tottori 680-8552, Japan. E-mail: kkoba@chem.tottori-u.ac.jp

Abstract – A convenient method for the preparation of 2-(alkyl(or aryl)sulfanyl)-2H-1,4-benzothiazin-3(4H)-one derivatives has been developed. Thus, methyl alkyl(2-{{(alkyl(or aryl)sulfanyl)methyl}sulfanyl}phenyl)carbamates, derived from commercially available 2-aminobenzenethiol by an easily operated three-step sequence, are treated with two equivalents of lithium diisopropylamide (LDA) to generate carbanions stabilized by the adjacent two sulfur atoms, which immediately undergo 2H-1,4-benzothiazin-3(4H)-one ring formation by intramolecular attack onto the carbamate carbonyl accompanying elimination of methoxide. In addition, direct or stepwise introductions of alkyl groups to the 2-position are reported.

Some compounds having the 2H-1,4-benzothiazin-3(4H)-one skeleton have been reported to have interesting biological profile,¹ exhibiting, for example, analgesic^{1a} and antitubercular^{1d} activities. Traditionally, the 2H-1,4-benzothiazin-3(4H)-one structure has been constructed by the reaction of 2-aminobenzenethiols with α -haloalkanoic acid derivatives in the presence of a variety of mediators.² There have been a few reports on the syntheses of 2H-1,4-benzothiazin-3(4H)-ones in which an alkyl(or aryl)sulfanyl group is attached to the 2-position. This type of 2H-1,4-benzothiazin-3(4H)-ones are also of biological interest.³ These derivatives have been conventionally prepared by 2-chlorination of 2H-1,4-benzothiazin-3(4H)-ones with appropriate chlorinating agents, followed by nucleophilic substitution with thiols.³ However, these methods suffer from incomplete generality, cumbersome operations, and/or rather harsh reaction conditions. Accordingly, we were interested in developing a new method for the general preparation of 2-sulfanylated 2H-1,4-benzothiazin-3(4H)-ones. In this paper, we wish to report the results of our investigation, which offer facile route to 2-(alkyl(or

aryl)sulfanyl)-2*H*-1,4-benzothiazin-3(4*H*)-ones (**5**). Our method involves formation of the 2*H*-1,4-benzothiazin-3(4*H*)-one structure by intramolecular cyclization of the carbanions stabilized by the adjacent two sulfur atoms of methyl alkyl(2-[(alkyl(or aryl)sulfanyl)methyl]sulfanyl)phenyl)carbamates (**4**), which are generated on treatment with lithium diisopropylamide (LDA).⁴ It is also reported that this strategy can be extended to the preparation of 2-alkylated 2-(alkyl(or aryl)sulfanyl)-2*H*-1,4-benzothiazin-3(4*H*)-ones (**6**) and (**7**). To the best of our knowledge, this is the first report which demonstrates a practical and general preparation of these classes of 2*H*-1,4-benzothiazin-3(4*H*)-ones.

The preparation of precursors (**4**) was achieved from commercially available 2-aminobenzenethiol (**1**) according to the easily operated three-step sequence illustrated in Scheme 1. Thus, successive treatment of compound (**1**) with sodium hydride and chloromethyl sulfides in DMF at 0 °C afforded the corresponding 2-[(alkyl(or aryl)sulfanyl)methyl]sulfanyl)benzenamines (**2**) in excellent yields. These amines were then *N*-methoxycarbonylated smoothly and cleanly upon treatment with methyl chloroformate in THF in the presence of pyridine at 0 °C to give methyl 2-[(alkyl(or aryl)sulfanyl)methyl]sulfanyl)phenylcarbamates (**3**) in good yields. *N*-Alkylation of these carbamates (**3**) by successive treatment with sodium hydride and alkyl halides in DMF at 0 °C provided **4** in good to excellent yields. These results are summarized in Table 1.



Scheme 1

Table 1. Preparation of 2-sulfanylated 2*H*-1,4-benzothiazin-3(4*H*)-ones (**5**)


Entry	R ¹	2	Yield/% ^a	3	Yield/% ^a	R ² X	4	Yield/% ^a	5	Yield/% ^a
1	Me	2a	93	3a	94	MeI	4a	97	5a	91
2						BnBr	4b	86	5b	77
3	<i>t</i> -Bu	2b	99	3b	77	BnBr	4c	82	5c	57
4	Ph	2c	98	3c	88	MeI	4d	83	5d	94
5	4-ClC ₆ H ₄	2d	94	3d	79	BnBr	4e	95	5e	84

^a Yields of isolated products.

The transformation of **4** into 2-(alkyl(or aryl)sulfanyl)-2*H*-1,4-benzothiazin-3(4*H*)-ones (**5**) is shown in Scheme 1 as well. When compounds (**4**) were treated with two equivalents of LDA in THF at -78 °C,

deprotonation of one of the protons of the methylene between the two sulfur atoms followed by attack of the resulting sulfur-stabilized carbanion onto carbamate carbonyl accompanying elimination of methoxide occurred to afford, after aqueous workup and subsequent purification by column chromatography on silica gel, the desired products (**5**) in generally good yields, as compiled in Table 1 too. However, the cyclization of *tert*-butylsulfanyl substrate (**4c**) resulted in the formation of a somewhat complicated mixture of products, and diminished yield of the product (**5c**) was obtained (Entry 3), probably due to bulkier *tert*-butyl substituent. When an equimolar amount of LDA was used, a considerable amount of the starting material was recovered in each case.

As mentioned in the introduction part, 2-alkyl-2-(alkyl(or aryl)sulfanyl)-2*H*-1,4-benzothiazin-3(4*H*)-ones (**6**) and (**7**) could be obtained conveniently. First, the introduction of alkyl groups to the 2-position of the 2-alkyl(or aryl)sulfanyl-2*H*-1,4-benzothiazin-3(4*H*)-ones was envisaged to be achieved by adding alkyl halides prior to aqueous work up after the treatment of **4** with two equivalents of LDA. In practice, alkyl halides listed in Table 2 were used successfully to give **6** directly, as shown in Scheme 2. The table indicates that yields of the products were generally good, though those of **6e** and **6h** (Entries 5 and 8) are somewhat lower than the others. These results may be ascribed to the steric bulkiness of *tert*-butylsulfanyl group and the lower reactivity of ethyl iodide compared to the other alkyl halides, respectively.



Scheme 2


Table 2. Preparation of 2-sulfanylated 2-alkyl-2*H*-1,4-benzothiazin-3(4*H*)-ones (**6**)

Entry	4	R ¹	R ²	R ³ X	6	Yield/% ^a
1	4a	Me	Me	MeI	6a	85
2	4a			CH ₂ =CHCH ₂ Br	6b	86
3	4a			BnBr	6c	81
4	4b	Me	Bn	MeI	6d	74
5	4c	<i>t</i> -Bu	Bn	MeI	6e	45
6	4d	Ph	Me	MeI	6f	70
7	4d			BnBr	6g	98
8	4e	4-ClC ₆ H ₄	Bn	EtI	6h	40
9	4e			CH ₂ =CHCH ₂ Br	6i	65

^a Yields of isolated products.

Attempts to obtain the corresponding 2-alkylated products in the same manner from **4** using *n*-butyl bromide, *tert*-butyl 2-bromoacetate, and chloromethyl sulfides proved to be unrewarding, probably due to

the reactions of these halides with lithium methoxide formed during the reaction. Fortunately, however, these alkyl substituents could be introduced using a two-step procedure from **4**. Thus, compounds **5** were successively treated with sodium hydride and these halides in DMF at 0 °C to afford the desired 2-alkylated products **7**, as shown in Scheme 3. As listed in Table 3, the yields of the products are moderate to good.



Scheme 3

Table 3. Preparation of 2-sulfanylated 2-alkyl-2*H*-1,4-benzothiazin-3(4*H*)-ones (**7**)

Entry	5	R ¹	R ³ X	7	Yield/% ^a
1	5b	Me	MeSCH ₂ Cl	7a	86
2	5b		<i>t</i> -BuOCOCH ₂ Br	7b	78
3	5c	<i>t</i> -Bu	PhSCH ₂ Cl	7c	54
4	5e	4-ClC ₆ H ₄	<i>n</i> -BuBr	7d	47
5	5e		<i>t</i> -BuSCH ₂ Cl	7e	52
6	5e		<i>t</i> -BuOCOCH ₂ Br	7f	71

^a Yields of isolated products.

In conclusion, we have demonstrated that LDA-mediated cyclization of alkyl(2-[(alkyl(or aryl)sulfanyl)methyl]sulfanyl)phenyl)carbamates opens up facile entry to 2-alkyl(or aryl)sulfanyl-2*H*-1,4-benzothiazin-3(4*H*)-ones and their 2-alkylated derivatives of potential biological interest. In comparison to the previously reported synthesis of 2*H*-1,4-benzothiazin-3(4*H*)-one derivatives, the advantages of the synthesis described here are that the reaction procedure is very simple and the reaction conditions are very mild. Applications of the present process to the synthesis of related heterocyclic derivatives are now in progress in our laboratory.

EXPERIMENTAL

All melting points were obtained on a Laboratory Devices MEL-TEMP II melting apparatus and are uncorrected. IR spectra were recorded with a Perkin–Elmer Spectrum 65 FTIR spectrophotometer. ¹H and ¹³C NMR spectra were recorded in CDCl₃ using TMS as an internal reference with a JEOL ECP500 FT NMR spectrometer operating at 500 and 125 MHz, respectively. High-resolution MS spectra were measured by a Thermo Scientific Exactive spectrometer (ESI and DART, positive) or a JEOL JMS-T100GCV (EI or FI, TOF; 70eV or 2100V, respectively) spectrometer. Elemental analyses were

performed with an Elementar Vario EL II instrument. TLC was carried out on Merck Kieselgel 60 PF₂₅₄. Column chromatography was performed using WAKO GEL C-200E. All of the organic solvents used in this study were dried over appropriate drying agents and distilled prior to use.

Starting Materials. 2-[(Chloromethyl)sulfanyl]-2-methylpropane⁵ and 1-chloro-4-[(chloromethyl)sulfanyl]-benzene⁶ were prepared according to the appropriate reported procedures. *n*-BuLi was supplied by Asia Lithium Corporation. All other chemicals used in this study were commercially available.

Typical Procedure for the Preparation of 2-[(Sulfanylmethyl)sulfanyl]benzenamines (2). 2-[[**(Methylsulfanyl)methyl**]sulfanyl]benzenamine (**2a**).⁷ To a stirred suspension of NaH (60% in mineral oil; 0.40 g, 10 mmol) in DMF (40 mL) at 0 °C was added 2-aminobenzenethiol (**1**) (1.3 g, 10 mmol) dropwise. After evolution of H₂ gas had ceased, ClCH₂SMe (0.97 g, 10 mmol) was added dropwise and stirring was continued at the same temperature for 5 min before addition of saturated aqueous NH₄Cl (80 mL). The resulting mixture was extracted with AcOEt (3 × 30 mL) and the combined extracts were washed with H₂O (3 × 40 mL) and brine (30 mL), and dried (Na₂SO₄). Evaporation of the solvent gave a residue, which was purified by column chromatography on SiO₂ to afford **2a** (1.7 g, 93%): a brown oil; *R*_f 0.46 (Et₂O/hexane 1:5). The spectral (IR and ¹H NMR) data for this compound were identical to those reported previously.⁷

2-[[(1,1-Dimethylethyl)sulfanyl**]methyl]sulfanyl]benzenamine (2b):** a brown oil; *R*_f 0.38 (Et₂O/hexane 1:3); IR (neat) 3457, 3352, 1607 cm⁻¹; ¹H NMR δ 1.35 (s, 9H), 3.88 (s, 2H), 4.38 (br s, 2H), 6.70 (td, *J* = 7.4, 1.1 Hz, 1H), 6.73 (d, *J* = 8.0 Hz, 1H), 7.13 (ddd, *J* = 8.0, 7.4, 1.1 Hz, 1H), 7.43 (dd, *J* = 7.4, 1.1 Hz, 1H); ¹³C NMR δ 31.0, 35.0, 43.8, 115.0, 117.9, 118.5, 130.3, 136.2, 148.3. HR-MS (DART). Calcd for C₁₁H₁₈NS₂ (M+H): 228.0880. Found: *m/z* 228.0870.

2-[[(Phenylsulfanyl)methyl**]sulfanyl]benzenamine (2c):** a brown oil; *R*_f 0.28 (CH₂Cl₂/hexane 1:3); IR (neat) 3458, 3360, 1606 cm⁻¹; ¹H NMR δ 4.19 (s, 2H), 4.29 (br s, 2H), 6.69 (t, *J* = 7.4 Hz, 1H), 6.71 (d, *J* = 8.0 Hz, 1H), 7.14 (ddd, *J* = 8.0, 7.4, 1.1 Hz, 1H), 7.24 (t, *J* = 7.4 Hz, 1H), 7.31 (t, *J* = 7.4 Hz, 2H); 7.39 (dd, *J* = 8.0, 1.1 Hz, 1H), 7.42 (d, *J* = 7.4 Hz, 2H); ¹³C NMR δ 40.7, 115.3, 116.6, 118.5, 126.9, 129.0, 130.4, 130.6, 135.1, 136.8, 148.6. HR-MS (EI). Calcd for C₁₃H₁₃NS₂ (M): 247.0489. Found: *m/z* 247.0485.

2-[[(4-Chlorophenyl)sulfanyl**]methyl]sulfanyl]benzenamine (2d):** a brown oil; *R*_f 0.39 (Et₂O/hexane 1:3); IR (neat) 3459, 3360, 1606 cm⁻¹; ¹H NMR δ 4.16 (s, 2H), 4.29 (br s, 2H), 6.67–6.72 (m, 2H), 7.15 (dd, *J* = 7.0, 6.9 Hz, 1H), 7.27 (d, *J* = 8.6 Hz, 2H), 7.33 (d, *J* = 8.6 Hz, 2H), 7.37 (d, *J* = 8.0 Hz, 1H); ¹³C NMR δ 40.7, 115.1, 116.3, 118.6, 129.1, 130.7, 131.8, 133.0, 133.6, 136.8, 148.6. Anal. Calcd for C₁₃H₁₂ClNS₂: C, 55.41; H, 4.29; N, 4.97. Found: C, 55.38; H, 4.52; N, 4.94.

Typical Procedure for the Preparation of Methyl Carbamates (3). **Methyl (2-{{[(Methylsulfonyl)methyl]sulfonyl}phenyl]carbamate (3a).** To a stirred solution of **2a** (1.72 g, 9.3 mmol) in THF (25 mL) containing pyridine (0.69 g, 9.3 mmol) at 0 °C was added ClCO₂Me (0.88 g, 9.3 mmol) dropwise. After 30 min, saturated aqueous NH₄Cl (30 mL) was added and the resulting mixture was extracted with AcOEt (3 × 25 mL). The combined extracts were washed with H₂O (2 × 20 mL) and brine (20 mL), dried (Na₂SO₄), and concentrated by evaporation. The residue was purified by column chromatography on SiO₂ to afford **3a** (2.1 g, 94%); a pale-yellow oil; *R*_f 0.25 (CH₂Cl₂/hexane 1:1); IR (neat) 3358, 1738 cm⁻¹; ¹H NMR δ 2.23 (s, 3H), 3.80 (s, 3H), 3.81 (s, 2H), 7.02 (td, *J* = 7.4, 1.1 Hz, 1H), 7.37 (td, *J* = 7.4, 1.1 Hz, 1H), 7.55 (dd, *J* = 7.4, 1.1 Hz, 1H), 8.01 (br s, 1H), 8.17 (d, *J* = 7.4 Hz, 1H); ¹³C NMR δ 15.5, 42.7, 52.4, 118.6, 120.8, 123.1, 130.5, 136.3, 140.0, 153.8. HR-MS (EI). Calcd for C₁₀H₁₃NO₂S₂ (M): 243.0388. Found: *m/z* 243.0380.

Methyl [(2-{{[(1,1-Dimethylethyl)sulfonyl]methyl}sulfonyl}phenyl]carbamate (3b): a pale-yellow oil; *R*_f 0.46 (CH₂Cl₂/hexane 1:4); IR (neat) 3359, 1741 cm⁻¹; ¹H NMR δ 1.35 (s, 9H), 3.80 (s, 3H), 3.86 (s, 2H), 7.02 (td, *J* = 7.4, 1.1 Hz, 1H), 7.35 (dd, *J* = 8.0, 7.4 Hz, 1H), 7.56 (d, *J* = 8.0 Hz, 1H), 8.02 (br s, 1H), 8.16 (d, *J* = 7.4 Hz, 1H); ¹³C NMR δ 30.9, 36.8, 44.0, 52.3, 118.7, 122.0, 123.2, 130.4, 135.9, 139.9, 153.9. HR-MS (DART). Calcd for C₁₃H₂₀NO₂S₂ (M+H): 286.0925. Found: *m/z* 286.0922.

Methyl (2-{{[(Phenylsulfonyl)methyl]sulfonyl}phenyl]carbamate (3c): a colorless oil; *R*_f 0.68 (CH₂Cl₂/hexane 1:4); IR (neat) 3359, 1738 cm⁻¹; ¹H NMR δ 3.76 (s, 3H), 4.17 (s, 2H), 7.00 (t, *J* = 7.4 Hz, 1H), 7.26 (dd, *J* = 7.4, 6.9 Hz, 1H), 7.32 (t, *J* = 7.4 Hz, 2H), 7.36 (t, *J* = 7.4 Hz, 1H), 7.39 (d, *J* = 7.4 Hz, 2H), 7.52 (dd, *J* = 6.9, 1.1 Hz, 1H), 7.87 (br s, 1H), 8.16 (d, *J* = 7.4 Hz, 1H); ¹³C NMR δ 42.4, 52.3, 118.6, 120.6, 123.2, 127.2, 129.1, 130.5, 130.7, 134.3, 136.4, 140.2, 153.7. HR-MS (EI). Calcd for C₁₅H₁₅NO₂S₂ (M): 305.0544. Found: *m/z* 305.0553.

Methyl [2-{{[(4-Chlorophenyl)sulfonyl]methyl}sulfonyl}phenyl]carbamate (3d): a white solid; mp 67–69 °C (hexane); IR (KBr) 3358, 1737 cm⁻¹; ¹H NMR δ 3.78 (s, 3H), 4.14 (s, 2H), 7.00 (t, *J* = 7.4 Hz, 1H), 7.28–7.32 (m, 4H), 7.36 (dd, *J* = 8.0, 7.4 Hz, 1H), 7.50 (d, *J* = 7.4 Hz, 1H), 7.78 (br s, 1H), 8.16 (d, *J* = 8.0 Hz, 1H); ¹³C NMR δ 42.5, 52.4, 118.6, 120.3, 123.2, 129.3, 130.8, 131.9, 132.7, 133.4, 136.4, 140.3, 153.6. Anal. Calcd for C₁₅H₁₄ClNO₂S₂: C, 53.01; H, 4.15; N, 4.12. Found: C, 52.88; H, 4.25; N, 4.08.

Typical Procedure for N-Alkylation of Carbamates (3). **Methyl Methyl(2-{{[(methylsulfonyl)methyl]sulfonyl}phenyl]carbamate (4a).** To a stirred suspension of NaH (60% in mineral oil; 0.18 g, 4.5 mmol) in DMF (20 mL) at 0 °C was added a solution of **3a** (1.1 g, 4.5 mmol) in THF (5 mL) dropwise. After evolution of H₂ gas had ceased, MeI (0.64 g, 4.5 mmol) was added dropwise and stirring was continued at the same temperature for 25 min. The resulting mixture was worked up as described for the preparation of **2a**. The crude product was purified by column chromatography on SiO₂

to afford **4a** (1.1 g, 97%); a pale-yellow oil; R_f 0.37 (CH_2Cl_2); IR (neat) 1712 cm^{-1} ; $^1\text{H NMR}$ δ 2.23 (s, 3H), 3.23 (s, 3H), 3.64 and 3.80 (2s, combined 3H), 3.95 (d, $J = 13.2\text{ Hz}$, 1H), 4.01 (d, $J = 13.2\text{ Hz}$, 1H), 7.17 (d, $J = 7.4\text{ Hz}$, 1H), 7.25 (t, $J = 7.4\text{ Hz}$, 1H), 7.30 (t, $J = 7.4\text{ Hz}$, 1H), 7.46 (d, $J = 7.4\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 15.2, 37.3, 38.7, 53.0, 127.3, 128.1, 128.2, 129.8, 134.8, 142.1, 156.2. HR-MS (EI). Calcd for $\text{C}_{11}\text{H}_{15}\text{NO}_2\text{S}_2$ (M): 257.0544. Found: m/z 257.0622.

Methyl (2-{{(Methylsulfanyl)methyl}sulfanyl}phenyl)(phenylmethyl)carbamate (4b): a colorless oil; R_f 0.44 (AcOEt/hexane 1:1); IR (neat) 1707 cm^{-1} ; $^1\text{H NMR}$ δ 2.23 (s, 3H), 3.67 (s, 3H), 3.86 (d, $J = 13.2\text{ Hz}$, 1H), 3.91 (d, $J = 13.2\text{ Hz}$, 1H), 4.32 (d, $J = 14.3\text{ Hz}$, 1H), 5.24 (d, $J = 14.3\text{ Hz}$, 1H), 6.79 (d, $J = 7.4\text{ Hz}$, 1H), 7.09 (t, $J = 7.4\text{ Hz}$, 1H), 7.25–7.29 (m, 6H), 7.46 (d, $J = 7.4\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 15.3, 38.7, 53.2, 55.4, 126.7, 127.5, 128.1, 128.3, 129.0, 129.5, 129.7, 135.1, 137.4, 140.2, 156.4. HR-MS (EI). Calcd for $\text{C}_{17}\text{H}_{19}\text{NO}_2\text{S}_2$ (M): 333.0857. Found: m/z 333.0852.

Methyl [(2-{{(1,1-Dimethylethyl)sulfanyl)methyl}sulfanyl}phenyl)(phenylmethyl)carbamate (4c): a pale-yellow oil; R_f 0.25 (Et_2O /hexane 1:3); IR (neat) 1709 cm^{-1} ; $^1\text{H NMR}$ δ 1.39 (s, 9H), 3.65 and 3.82 (2s, combined 3H), 4.09 (s, 2H), 4.29 (d, $J = 14.3\text{ Hz}$, 1H), 5.23 (d, $J = 14.3\text{ Hz}$, 1H), 6.75 (d, $J = 7.4\text{ Hz}$, 1H), 7.05 (t, $J = 7.4\text{ Hz}$, 1H), 7.23–7.27 (m, 6H), 7.45 (d, $J = 7.4\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 31.0, 32.6, 44.0, 53.1, 53.3, 126.4, 127.4, 128.2, 128.3 (2 overlapped Cs), 128.9, 129.1, 129.6, 136.1, 137.5, 156.3. HR-MS (DART). Calcd for $\text{C}_{20}\text{H}_{26}\text{NO}_2\text{S}_2$ (M+H): 376.1405. Found: m/z 376.1390.

Methyl Methyl(2-{{(phenylsulfanyl)methyl}sulfanyl}phenyl)carbamate (4d): a pale-yellow oil; R_f 0.53 (AcOEt/hexane 1:2); IR (neat) 1712 cm^{-1} ; $^1\text{H NMR}$ δ 3.16 (s, 3H), 3.59 and 3.77 (2s, combined 3H), 4.32 (s, 2H), 7.16 (d, $J = 7.4\text{ Hz}$, 1H), 7.25–7.32 (m, 5H), 7.42 (d, $J = 7.4\text{ Hz}$, 2H), 7.47 (d, $J = 7.4\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 37.3, 38.8, 53.0, 60.4, 127.2, 127.5, 128.16, 128.23, 129.0, 129.8, 130.6, 134.8, 142.0, 156.2. HR-MS (EI). Calcd for $\text{C}_{16}\text{H}_{17}\text{NO}_2\text{S}_2$ (M): 319.0701. Found: m/z 319.0689.

Methyl [(2-{{(4-Chlorophenyl)sulfanyl)methyl}sulfanyl}phenyl)(phenylmethyl)carbamate (4e): a pale-yellow oil; R_f 0.26 (Et_2O /hexane 1:3); IR (neat) 1705 cm^{-1} ; $^1\text{H NMR}$ δ 3.62 and 3.82 (2s, combined 3H), 4.15 (s, 2H), 4.27 (d, $J = 14.9\text{ Hz}$, 1H), 5.14 (d, $J = 14.9\text{ Hz}$, 1H), 6.81 (d, $J = 6.9\text{ Hz}$, 1H), 7.11 (dd, $J = 7.4, 6.9\text{ Hz}$, 1H), 7.22–7.29 (m, 8H), 7.35 (d, $J = 8.0\text{ Hz}$, 2H), 7.45 (d, $J = 7.4\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 39.0, 53.0, 53.4, 127.1, 127.4, 128.1, 128.2 (2 overlapped Cs), 129.0, 129.1, 129.6, 130.0, 131.8, 133.2, 134.6, 137.3, 140.3, 156.2. HR-MS (EI). Calcd for $\text{C}_{22}\text{H}_{20}\text{ClNO}_2\text{S}_2$ (M): 429.0624. Found: m/z 429.0617.

Typical Procedure for the Preparation of Benzothiazinones (5). 4-Methyl-2-(methylsulfanyl)-2H-1,4-benzothiazin-3(4H)-one (5a). To a stirred solution of LDA (1.6 mmol), generated by the standard method from *n*-BuLi and *i*-Pr₂NH, in THF 6 mL at $-78\text{ }^\circ\text{C}$ was added a solution of **4a** (0.21 g, 0.80 mmol) in THF (2 mL) dropwise. After 10 min, the mixture was worked up as described for the preparation of **2a**. The crude product was purified by column chromatography on SiO_2 to afford **5a** (0.17 g, 91%); a

pale-yellow oil; R_f 0.33 (AcOEt/hexane 1:5); IR (neat) 1663 cm^{-1} ; $^1\text{H NMR}$ δ 2.21 (s, 3H), 3.49 (s, 3H), 4.52 (s, 1H), 7.07 (td, $J = 7.4, 1.1\text{ Hz}$, 1H), 7.11 (dd, $J = 8.0, 1.1\text{ Hz}$, 1H), 7.31 (ddd, $J = 8.0, 7.4, 1.1\text{ Hz}$, 1H), 7.36 (dd, $J = 8.0, 1.1\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 15.8, 32.3, 47.7, 117.1, 119.2, 123.6, 127.6, 129.1, 139.4, 163.1. HR-MS (EI). Calcd for $\text{C}_{10}\text{H}_{11}\text{NOS}_2$ (M): 225.0282. Found: m/z 225.0286.

2-(Methylsulfanyl)-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (5b): a white solid; mp 71–73 °C (hexane); IR (KBr) 1656 cm^{-1} ; $^1\text{H NMR}$ δ 2.24 (s, 3H), 4.58 (s, 1H), 4.87 (d, $J = 16.6\text{ Hz}$, 1H), 5.62 (d, $J = 16.6\text{ Hz}$, 1H), 6.99 (dd, $J = 8.0, 1.1\text{ Hz}$, 1H), 7.03 (ddd, $J = 8.0, 7.4, 1.1\text{ Hz}$, 1H), 7.17 (ddd, $J = 8.0, 7.4, 1.1\text{ Hz}$, 1H), 7.23–7.27 (m, 3H), 7.33 (t, $J = 7.4\text{ Hz}$, 2H), 7.36 (dd, $J = 8.0, 1.1\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 15.8, 47.6, 48.5, 117.5, 119.0, 123.8, 125.8, 127.2, 127.6, 128.8, 129.0, 136.5, 138.9, 162.9. HR-MS (EI). Calcd for $\text{C}_{16}\text{H}_{15}\text{NOS}_2$ (M): 301.0595. Found: m/z 301.0595. Anal. Calcd for $\text{C}_{16}\text{H}_{15}\text{NOS}_2$: C, 63.76; H, 5.02; N, 4.65. Found: C, 63.72; H, 5.09; N, 4.54.

2-[(1,1-Dimethylethyl)sulfanyl]-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (5c): a colorless oil; R_f 0.33 (Et₂O/hexane 1:5); IR (neat) 1664 cm^{-1} ; $^1\text{H NMR}$ δ 1.41 (s, 9H), 4.68 (s, 1H), 4.86 (d, $J = 16.6\text{ Hz}$, 1H), 5.58 (d, $J = 16.6\text{ Hz}$, 1H), 7.00 (d, $J = 8.0\text{ Hz}$, 1H), 7.02 (dd, $J = 8.0, 7.4\text{ Hz}$, 1H), 7.15 (ddd, $J = 8.0, 7.4, 1.1\text{ Hz}$, 1H), 7.23 (t, $J = 7.4\text{ Hz}$, 1H), 7.28 (d, $J = 7.4\text{ Hz}$, 2H), 7.32 (t, $J = 7.4\text{ Hz}$, 2H), 7.36 (dd, $J = 8.0, 1.1\text{ Hz}$, 1H); $^{13}\text{C NMR}$ δ 31.0, 44.3, 46.2, 48.7, 117.7, 119.9, 123.7, 126.1, 127.1, 127.7, 128.7, 129.4, 136.7, 139.3, 164.2. HR-MS (ESI). Calcd for $\text{C}_{19}\text{H}_{22}\text{NOS}_2$ (M+H): 344.1145. Found: m/z 344.1128.

4-Methyl-2-(phenylsulfanyl)-2H-1,4-benzothiazin-3(4H)-one (5d): a white solid; mp 65–67 °C (hexane); IR (neat) 1666 cm^{-1} ; $^1\text{H NMR}$ δ 3.44 (s, 3H), 4.87 (s, 1H), 7.07–7.11 (m, 2H), 7.28–7.33 (m, 4H), 7.37 (dd, $J = 6.9, 1.1\text{ Hz}$, 1H), 7.42–7.45 (m, 2H); $^{13}\text{C NMR}$ δ 32.3, 50.4, 117.2, 119.4, 123.7, 127.7, 128.9, 129.0, 129.2, 132.1, 134.0, 139.6, 162.7. HR-MS (EI). Calcd for $\text{C}_{15}\text{H}_{13}\text{NOS}_2$ (M): 287.0439. Found: m/z 287.0429. Anal. Calcd for $\text{C}_{15}\text{H}_{13}\text{NOS}_2$: C, 62.69; H, 4.56; N, 4.87. Found: C, 62.55; H, 4.58; N, 4.87.

2-[(4-Chlorophenyl)sulfanyl]-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (5e): a pale-yellow solid; mp 124–126 °C (hexane/ CH_2Cl_2); IR (KBr) 1642 cm^{-1} ; $^1\text{H NMR}$ δ 4.88 (d, $J = 16.0\text{ Hz}$, 1H), 4.92 (s, 1H), 5.53 (d, $J = 16.0\text{ Hz}$, 1H), 7.03–7.07 (m, 2H), 7.17–7.27 (m, 6H), 7.31 (dd, $J = 8.0, 7.4\text{ Hz}$, 2H), 7.35 (dd, $J = 7.4, 1.1\text{ Hz}$, 1H), 7.40 (d, $J = 8.6\text{ Hz}$, 2H); $^{13}\text{C NMR}$ δ 48.8, 50.3, 117.8, 119.1, 124.0, 126.1, 127.3, 127.9, 128.8, 129.2, 129.3, 130.7, 134.8, 135.1, 136.4, 139.2, 162.6. HR-MS (EI). Calcd for $\text{C}_{21}\text{H}_{16}\text{ClNOS}_2$ (M): 397.0362. Found: m/z 397.0363. Anal. Calcd for $\text{C}_{21}\text{H}_{16}\text{ClNOS}_2$: C, 63.38; H, 4.05; N, 3.52. Found: C, 63.20; H, 4.16; N, 3.48.

Typical Procedure for the Preparation of 2-Alkylated 2H-1,4-Benzothiazin-3(4H)-ones (6). 2,4-Dimethyl-2-(methylsulfanyl)-2H-1,4-benzothiazin-3(4H)-one (6a). Compound **2a** (0.21 g, 0.81 mmol)

was treated with LDA (1.6 mmol) in THF (6 mL) as described for the preparation of **5a**. Then, MeI (0.11 g, 0.81 mmol) was added at the same temperature. After 30 min, the mixture was worked up as described for the preparation of **2a** and the residue was purified by column chromatography on SiO₂ to afford **6a** (1.7 g, 85%); a colorless oil; *R*_f 0.24 (AcOEt/hexane 1:10); IR (neat) 1661 cm⁻¹; ¹H NMR δ 1.89 (s, 3H), 2.03 (s, 3H), 3.50 (s, 3H), 7.06 (td, *J* = 7.4, 1.1 Hz, 1H), 7.09 (dd, *J* = 7.4, 1.1 Hz, 1H), 7.30–7.33 (m, 2H); ¹³C NMR δ 13.3, 23.1, 33.0, 50.8, 116.7, 120.4, 123.3, 127.5, 128.1, 140.0, 165.6. HR-MS (EI). Calcd for C₁₁H₁₃NOS₂ (M): 239.0439. Found: *m/z* 239.0434.

4-Methyl-2-(methylsulfanyl)-2-(prop-2-enyl)-2H-1,4-benzothiazin-3(4H)-one (6b): a pale-yellow oil; *R*_f 0.32 (AcOEt/hexane 1:7); IR (neat) 1659 cm⁻¹; ¹H NMR δ 2.00 (s, 3H), 3.00 (d, *J* = 6.9 Hz, 2H), 3.50 (s, 3H), 5.23 (d, *J* = 10.3 Hz, 1H), 5.26 (d, *J* = 19.5 Hz, 1H), 5.96–6.03 (m, 1H), 7.05 (t, *J* = 7.4 Hz, 1H), 7.06 (d, *J* = 7.4 Hz, 1H), 7.31 (t, *J* = 7.4 Hz, 1H), 7.32 (d, *J* = 7.4 Hz, 1H); ¹³C NMR δ 13.6, 32.8, 39.2, 54.2, 116.4, 119.4, 119.7, 123.3, 127.5, 128.4, 132.3, 139.6, 164.8. HR-MS (EI). Calcd for C₁₃H₁₅NOS₂ (M): 265.0595. Found: *m/z* 265.0583.

4-Methyl-2-(methylsulfanyl)-2-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (6c): a colorless viscous oil; *R*_f 0.29 (AcOEt/hexane 1:10); IR (neat) 1648 cm⁻¹; ¹H NMR δ 2.06 (s, 3H), 3.44 (d, *J* = 14.3 Hz, 1H), 3.50 (s, 3H), 3.68 (d, *J* = 14.3 Hz, 1H), 7.00 (td, *J* = 7.4, 1.1 Hz, 1H), 7.05 (dd, *J* = 8.0, 1.1 Hz, 1H), 7.24 (dd, *J* = 7.4, 1.1 Hz, 1H), 7.26–7.34 (m, 4H), 7.39 (dd, *J* = 8.0, 1.1 Hz, 2H); ¹³C NMR δ 14.0, 33.0, 40.6, 56.0, 116.3, 119.7, 123.9, 127.4, 127.9 (2 overlapped Cs), 128.4, 131.0, 135.2, 139.4, 165.0. HR-MS (EI). Calcd for C₁₇H₁₇NOS₂ (M): 315.0752. Found: *m/z* 315.0749. Anal. Calcd for C₁₇H₁₇NOS₂: C, 64.73; H, 5.43; N, 4.44. Found: C, 64.68; H, 5.53; N, 4.35.

2-Methyl-2-(methylsulfanyl)-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (6d): a white solid; mp 90–92 °C (hexane); IR (KBr) 1661 cm⁻¹; ¹H NMR δ 1.92 (s, 3H), 2.06 (s, 3H), 4.84 (d, *J* = 16.6 Hz, 1H), 5.62 (d, *J* = 16.6 Hz, 1H), 6.98 (d, *J* = 8.0 Hz, 1H), 7.03 (t, *J* = 7.4 Hz, 1H), 7.18 (ddd, *J* = 8.0, 7.4, 1.1 Hz, 1H), 7.24–7.26 (m, 3H), 7.31–7.34 (m, 3H); ¹³C NMR δ 13.1, 22.8, 49.4, 50.6, 117.1, 120.1, 123.5, 125.7, 127.1, 127.6, 128.0, 128.8, 137.0, 139.8, 165.3. HR-MS (FI). Calcd for C₁₇H₁₇NOS₂ (M): 315.0752. Found: *m/z* 315.0760.

2-[2-(1,1-Dimethylethyl)sulfanyl]-2-methyl-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (6e): a pale-yellow oil; *R*_f 0.33 (AcOEt/hexane 1:20); IR (neat) 1660 cm⁻¹; ¹H NMR δ 1.31 (s, 9H), 2.12 (s, 3H), 5.19 (d, *J* = 16.6 Hz, 1H), 5.39 (d, *J* = 16.6 Hz, 1H), 6.99 (ddd, *J* = 8.0, 7.4, 1.1 Hz, 1H), 7.04 (d, *J* = 8.0 Hz, 1H), 7.16 (td, *J* = 7.4, 1.1 Hz, 1H), 7.20 (t, *J* = 7.4 Hz, 1H), 7.23–7.29 (m, 5H); ¹³C NMR δ 26.4, 31.9, 48.6, 48.9, 51.6, 117.3, 121.4, 123.3, 126.6, 127.1, 127.4, 128.1, 128.6, 136.7, 139.4, 166.7. HR-MS (DART). Calcd for C₂₀H₂₄NOS₂ (M+H): 358.1299. Found: *m/z* 358.1285.

2,4-Dimethyl-2-(phenylsulfanyl)-2H-1,4-benzothiazin-3(4H)-one (6f): a white solid; mp 113–115 °C (hexane); IR (KBr) 1659 cm⁻¹; ¹H NMR δ 1.82 (s, 3H), 3.45 (s, 3H), 7.08–7.11 (m, 2H), 7.28–7.38 (m, 7H); ¹³C NMR δ 24.2, 33.0, 54.5, 116.8, 120.9, 123.3, 127.7, 128.2, 128.7, 129.8, 130.2, 136.7, 140.2, 165.2. HR-MS (EI). Calcd for C₁₆H₁₅NOS₂ (M): 301.0595. Found: *m/z* 301.0597. Anal. Calcd for C₁₆H₁₅NOS₂: C, 63.76; H, 5.02; N, 4.65. Found: C, 63.54; H, 4.95; N, 4.67.

4-Methyl-2-(phenylmethyl)-2-(phenylsulfanyl)-2H-1,4-benzothiazin-3(4H)-one (6g): a pale-yellow viscous oil; *R*_f 0.23 (AcOEt/hexane 1:7); IR (neat) 1661 cm⁻¹; ¹H NMR δ 3.37 (s, 3H), 3.41 (d, *J* = 14.3 Hz, 1H), 3.83 (d, *J* = 14.3 Hz, 1H), 7.00–7.04 (m, 2H), 7.26–7.37 (m, 10H), 7.44 (d, *J* = 6.3 Hz, 2H); ¹³C NMR δ 32.8, 42.4, 60.1, 116.3, 120.4, 123.2, 127.4, 127.5, 127.9, 128.5, 128.7, 129.8, 130.7, 131.1, 135.6, 136.5, 139.5, 164.3. HR-MS (ESI). Calcd for C₂₂H₁₉NNaOS₂ (M+Na): 400.0806. Found: *m/z* 400.0802. Anal. Calcd for C₂₂H₁₉NOS₂: C, 69.99; H, 5.07; N, 3.71. Found: C, 70.03; H, 5.11; N, 3.64.

2-[(4-Chlorophenyl)sulfanyl]-2-ethyl-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (6h): a colorless oil; *R*_f 0.36 (AcOEt/hexane 1:30); IR (neat) 1661 cm⁻¹; ¹H NMR δ 1.17 (t, *J* = 7.4 Hz, 3H), 2.03–2.10 (m, 1H), 2.23–2.30 (m, 1H), 5.05 (d, *J* = 16.6 Hz, 1H), 5.39 (d, *J* = 16.6 Hz, 1H), 7.03 (t, *J* = 7.4 Hz, 1H), 7.08 (d, *J* = 8.6 Hz, 1H), 7.18 (d, *J* = 8.6 Hz, 2H), 7.21 (d, *J* = 8.6 Hz, 2H), 7.24–7.35 (m, 7H); ¹³C NMR δ 9.3, 28.8, 49.3, 59.7, 117.3, 119.9, 123.6, 126.6, 127.2, 127.7, 128.5, 128.7, 128.88, 128.92, 135.9, 136.8, 137.3, 139.4, 165.0. HR-MS (DART). Calcd for C₂₃H₂₁ClNOS₂ (M+H): 426.0753. Found: *m/z* 426.0735.

2-[(4-Chlorophenyl)sulfanyl]-4-(phenylmethyl)-2-(prop-2-enyl)-2H-1,4-benzothiazin-3(4H)-one (6i): a colorless oil; *R*_f 0.34 (AcOEt/hexane 1:20); IR (neat) 1662 cm⁻¹; ¹H NMR δ 2.78 (dd, *J* = 14.9, 7.4 Hz, 1H), 3.02 (dd, *J* = 14.9, 6.9 Hz, 1H), 5.04 (d, *J* = 16.6 Hz, 1H), 5.19–5.23 (m, 2H), 5.41 (d, *J* = 16.6 Hz, 1H), 5.98–6.06 (m, 1H), 7.05 (t, *J* = 7.4 Hz, 1H), 7.10 (d, 8.0 Hz, 1H), 7.19 (d, *J* = 8.6 Hz, 2H), 7.22 (d, *J* = 8.6 Hz, 2H), 7.26–7.34 (m, 7H); ¹³C NMR δ 40.2, 49.4, 57.7, 117.3, 119.8, 120.0, 123.7, 126.6, 127.3, 127.8, 128.5, 128.6, 128.7, 129.0, 132.2, 136.1, 136.7, 137.4, 139.4, 164.6. HR-MS (DART). Calcd for C₂₄H₂₁ClNOS₂ (M+H): 438.0753. Found: *m/z* 438.0735.

Typical Procedure for the Preparation of 2-Alkylated 2H-1,4-Benzothiazin-3(4H)-ones (7).

2-(Methylsulfanyl)-2-[(methylsulfanyl)methyl]-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (7a). To a stirred suspension of NaH (60% in mineral oil; 23 mg, 0.57 mmol) in DMF (4 mL) at 0 °C was added dropwise a solution of **5b** (0.17 g, 0.57 mmol) in DMF (1 mL). After evolution of H₂ gas had ceased, ClCH₂SMe (55 mg, 0.57 mmol) was added and stirring was continued for 5 min at the same temperature. The mixture was worked up as described for the preparation of **2a** and the residue was purified by column chromatography on SiO₂ to afford **7a** (0.18 g, 86%); a yellow oil; *R*_f 0.48 (AcOEt/hexane 1:5); IR (neat) 1655 cm⁻¹; ¹H NMR δ 2.04 (s, 3H), 2.31 (s, 3H), 3.30 (d, *J* = 13.7 Hz, 1H),

3.60 (d, $J = 13.7$ Hz, 1H), 4.86 (d, $J = 16.6$ Hz, 1H), 5.62 (d, $J = 16.6$ Hz, 1H), 6.98 (d, $J = 8.0$ Hz, 1H), 7.04 (t, $J = 7.4$ Hz, 1H), 7.18 (dd, $J = 8.0, 7.4$ Hz, 1H), 7.23–7.26 (m, 3H), 7.32 (dd, $J = 8.0, 7.4$ Hz, 2H), 7.37 (d, $J = 7.4$ Hz, 1H); ^{13}C NMR δ 13.6, 18.5, 40.0, 49.4, 56.8, 117.0, 119.3, 123.7, 125.7, 127.2, 127.6, 128.4, 128.8, 136.8, 139.0, 164.1. HR-MS (FI). Calcd for $\text{C}_{18}\text{H}_{19}\text{NOS}_3$ (M): 361.0629. Found: m/z 361.0640.

1,1-Dimethylethyl 2-[2-(Methylsulfanyl)-3-oxo-4-(phenylmethyl)-2H-1,4-benzothiazin-2-yl]acetate (7b): a yellow viscous oil; R_f 0.33 (AcOEt/hexane 1:5); IR (neat) 1733, 1660 cm^{-1} ; ^1H NMR δ 1.50 (s, 9H), 2.06 (s, 3H), 3.18 (d, $J = 15.5$ Hz, 1H), 3.23 (d, $J = 15.5$ Hz, 1H), 4.84 (d, $J = 12.0$ Hz, 1H), 4.87 (d, $J = 12.0$ Hz, 1H), 6.96 (dd, $J = 8.0, 1.1$ Hz, 1H), 7.03 (ddd, $J = 8.0, 7.4, 1.1$ Hz, 1H), 7.16 (ddd, $J = 8.0, 7.4, 1.7$ Hz, 1H), 7.22–7.26 (m, 3H), 7.32 (t, $J = 7.4$ Hz, 2H), 7.36 (dd, $J = 8.0, 1.7$ Hz, 1H); ^{13}C NMR δ 13.9, 28.0, 40.9, 49.4, 51.6, 81.7, 116.9, 119.4, 123.6, 125.7, 125.8, 127.1, 127.5, 128.2, 136.9, 139.0, 163.8, 168.1. HR-MS (ESI). Calcd for $\text{C}_{22}\text{H}_{25}\text{NNaO}_3\text{S}_2$ (M+Na): 438.1174. Found: m/z 438.1170.

2-[2-(1,1-Dimethylethyl)sulfanyl]-4-(phenylmethyl)-2-[(phenylsulfanyl)methyl]-2H-1,4-benzothiazin-3(4H)-one (7c): a yellow oil; R_f 0.28 (AcOEt/hexane 1:20); IR (neat) 1652 cm^{-1} ; ^1H NMR δ 1.30 (s, 9H), 3.99 (d, $J = 13.7$ Hz, 1H), 4.20 (d, $J = 13.7$ Hz, 1H), 5.03 (d, $J = 16.6$ Hz, 1H), 5.62 (d, $J = 16.6$ Hz, 1H), 6.98 (t, $J = 7.4$ Hz, 1H), 7.07 (d, $J = 8.0$ Hz, 1H), 7.16 (dd, $J = 8.0, 7.4$ Hz, 1H), 7.20–7.32 (m, 9H), 7.52 (d, $J = 8.0$ Hz, 2H); ^{13}C NMR δ 31.6, 44.0, 48.2, 49.5, 57.9, 117.4, 121.2, 123.6, 126.6, 126.9, 127.2, 127.3, 128.58, 128.62, 129.0, 130.7, 136.3, 137.2, 138.2, 165.5. HR-MS (ESI). Calcd for $\text{C}_{26}\text{H}_{27}\text{NNaOS}_2$ (M+Na): 488.1153. Found: m/z 488.1138.

2-Butyl-2-[(4-chlorophenyl)sulfanyl]-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (7d): a colorless gum; R_f 0.35 (Et₂O/hexane 1:15); IR (neat) 1659 cm^{-1} ; ^1H NMR δ 0.90 (t, $J = 7.4$ Hz, 3H), 1.25–1.39 (m, 2H), 1.53–1.70 (m, 2H), 1.97–2.03 (m, 1H), 2.17–2.23 (m, 1H), 5.04 (d, $J = 16.0$ Hz, 1H), 5.38 (d, $J = 16.0$ Hz, 1H), 6.99–7.11 (m, 4H), 7.15–7.34 (m, 7H), 7.52 (d, $J = 8.6$ Hz, 2H); ^{13}C NMR δ 13.9, 22.6, 26.9, 35.4, 49.3, 59.2, 117.3, 117.6, 123.9, 126.6, 126.7, 127.4, 127.7, 127.8, 128.7, 128.9, 136.3, 136.5, 137.4, 139.3, 166.4. HR-MS (DART). Calcd for $\text{C}_{25}\text{H}_{25}\text{ClNOS}_2$ (M+H): 454.1066. Found: m/z 454.1050.

2-[(4-Chlorophenyl)sulfanyl]-2-[[2-(1,1-dimethylethyl)sulfanyl]methyl]-4-(phenylmethyl)-2H-1,4-benzothiazin-3(4H)-one (7e): a colorless gum; R_f 0.35 (AcOEt/hexane 1:15); IR (neat) 1659 cm^{-1} ; ^1H NMR δ 1.33 (s, 9H), 3.24 (d, $J = 13.2$ Hz, 1H), 3.28 (d, $J = 13.2$ Hz, 1H), 5.03 (d, $J = 16.0$ Hz, 1H), 5.41 (d, $J = 16.0$ Hz, 1H), 7.05 (t, $J = 7.4$ Hz, 1H), 7.12 (d, $J = 8.0$ Hz, 1H), 7.18 (d, $J = 8.6$ Hz, 2H), 7.22 (dd, $J = 8.0, 7.4$ Hz, 1H), 7.27–7.31 (m, 3H), 7.31–7.34 (m, 4H), 7.36 (dd, $J = 7.4, 1.1$ Hz, 1H); ^{13}C NMR δ 30.9, 34.3, 42.9, 49.5, 58.6, 117.3, 119.9, 123.8, 126.8, 127.4, 127.8, 128.3, 128.70, 128.73, 129.0, 136.3,

136.7, 137.6, 139.1, 164.4. HR-MS (DART). Calcd for C₂₆H₂₇ClNOS₃ (M+H): 500.0943. Found: *m/z* 500.0922.

1,1-Dimethylethyl 2-{2-[(4-Chlorophenyl)sulfanyl]-3-oxo-4-(phenylmethyl)-2*H*-1,4-benzothiazin-2-yl}acetate (7f): a colorless gum; *R*_f 0.29 (Et₂O/hexane 1:5); IR (neat) 1734, 1663 cm⁻¹; ¹H NMR δ 1.49 (s, 9H), 3.00 (d, *J* = 16.0 Hz, 1H), 3.17 (d, *J* = 16.0 Hz, 1H), 5.06 (d, *J* = 16.6 Hz, 1H), 5.36 (d, *J* = 16.6 Hz, 1H), 7.02 (t, *J* = 7.4 Hz, 1H), 7.10 (d, *J* = 8.0 Hz, 1H), 7.21 (d, *J* = 8.6 Hz, 2H), 7.25–7.36 (m, 9H); ¹³C NMR δ 28.1, 42.0, 49.4, 55.9, 81.8, 117.3, 120.3, 123.7, 126.8, 127.3, 127.7, 128.3, 128.4, 128.7, 129.1, 136.5, 136.5, 137.9, 139.0, 163.9, 167.9. HR-MS (ESI). Calcd for C₂₇H₂₆NNaO₃S₂ (M+Na): 534.0941. Found: *m/z* 534.0919.

ACKNOWLEDGEMENTS

Recording mass spectra and performing combustion analyses by Mrs. Miyuki Tanmatsu of this university are gratefully acknowledged.

REFERENCES AND NOTES

- (a) J. Gowda, A. M. A. Khader, B. Kalluraya, P. Shree, and A. R. Shabaraya, *Eur. J. Med. Chem.*, **2011**, *46*, 4100; (b) D. Strøbæk, D. T. Brown, D. P. Jenkins, Y.-J. Chen, N. Coleman, Y. Ando, P. Chiu, S. Jørgensen, J. Demnitz, H. Wulff, and P. Christophersen, *Br. J. Pharmacol.*, **2013**, *168*, 432; (c) J. Goveas, A. M. A. Khader, B. Kalluraya, S. Sheik, and K. R. Chandrashekar, *Der Pharma Chem.*, **2015**, *7*, 248; (d) M. H. Shaikh, D. D. Subhedar, M. Arkile, V. M. Khedkar, N. Jadhav, D. Sarkar, and B. B. Shingate, *Bioorg. Med. Chem. Lett.*, **2016**, *26*, 561; (e) R. Venkatesh, S. Kasaboina, D. Bidayat, N. Kumar, U. N. Jain, S. J. Tangeda, R. Bantu, S. Janardhan, and L. Nagarapu, *Bioorg. Med. Chem. Lett.*, **2017**, *27*, 354.
- (a) H. Shimizu, N. Ueda, T. Kataoka, and M. Hori, *Chem. Pharm. Bull.*, **1984**, *32*, 2571; (b) A. R. Katritzky, H. H. Odens, and S. Zhang, *J. Org. Chem.*, **2001**, *66*, 6792; (c) A. Sharifi, M. S. Abaee, M. Rouzgard, and M. Mirzaei, *Synth. Commun.*, **2013**, *43*, 2079; (d) A. Sharifi, M. Ansari, H. R. Darabi, and M. S. Abaee, *Tetrahedron Lett.*, **2016**, *57*, 529.
- (a) J. Iwao, T. Iso, and M. Oya, *PCT Int. Appl.*, 1986, WO 8605490 (*Chem. Abstr.*, 1987, **106**, 18586); (b) M. Fujita, S. Ito, A. Ota, N. Kato, K. Yamamoto, Y. Kawashima, H. Yamauchi, and J. Iwao, *J. Med. Chem.*, **1990**, *33*, 1898; (c) K. Fujimura, M. Fujita, H. Suhara, and Y. Kawashima, *Bioorg. Med. Chem.*, **1994**, *2*, 235.
- Similar base-mediated cyclization reactions affording 2-alkyl(or aryl)-(1,1-dioxo)-2*H*-1,4-benzothiazin-3(4*H*)-ones have already been reported: (a) Y. Ishikawa, Y. Terao, K. Suzuki, N. Shikano, and M. Sekiya, *Chem. Pharm. Bull.* **1984**, *32*, 438; (b) A. I. Gerasyuto, S. G. Zlotin, and V.

- V. Semenov, [Synthesis, 2001, 300](#).
5. M. Maccoss, F. G. Njoroge, F. Bennett, S. Venkatraman, J. A. Kozlowski, W. Yu, P. E. McNamara, and P. R. Royster, *PCT Int. Appl.*, 2010, WO 2010068714 (*Chem. Abstr.*, 2010, **153**, 88109).
 6. Y. Tamura, H. Annoura, M. Fuji, M. Okura, and H. Ishibashi, [Chem. Pharm. Bull., 1986, 34, 540](#).
 7. H. Shimizu, A. Sugimoto, T. Kataoka, and M. Hori, [Heteroat. Chem., 1995, 6, 167](#).