

HETEROCYCLES, Vol. 96, No. 9, 2018, pp. 1631 - 1637. © 2018 The Japan Institute of Heterocyclic Chemistry
Received, 10th July, 2018, Accepted, 9th August, 2018, Published online, 15th August, 2018
DOI: 10.3987/COM-18-13958

TWO NEW C₁₉-DITERPENOID ALKALOIDS FROM *ACONITUM FRANCHETII* VAR. *VILLOSULUM*

Wenliang Xu,^{a†} Lin Chen,^{b†} Feng Gao,^a and Xianli Zhou^{a*}

^aSchool of Life Science and Engineering, Southwest Jiaotong University, Chengdu 610031, Sichuan, People's Republic of China. ^bSchool of Chemistry and Chemical Engineering, China West Normal University, Nanchong, 637002, Sichuan, People's Republic of China. E-mail: zhouxl@swjtu.edu.cn [†]These authors contributed equally to this work and should be considered co-first authors.

Abstract – Two new C₁₉-diterpenoid alkaloids, named villosudine A (**1**) and villosudine B (**2**), along with seven known diterpenoid alkaloids, were isolated from the root of *Aconitum franchetii* var. *villosulum*. Their structures were elucidated by extensive spectroscopic analyses including 1D, 2D NMR, and HR-ESI-MS. Compounds (**1-7**) were evaluated for their cytotoxicity against the MCF-7 and HepG2 human cancer cell lines.

Aconitum franchetii var. *villosulum* belongs to the family Ranunculaceae and has a wide distribution in Sichuan province of China. Our previous phytochemical studies on this herb have led to the isolation of eleven diterpenoid alkaloids.¹ Diterpenoid alkaloids are typical chemical components of the genus *Aconitum*, and display various bioactivities such as anti-inflammatory, anti-arrhythmia, antifungal, and cytotoxic properties, as well as antiviral, insecticidal and antifeedant activities.² To find more biologically active secondary metabolites, two new C₁₉-diterpenoid alkaloid, named villosudine A (**1**) and villosudine B (**2**), along with seven known diterpenoid alkaloids, were isolated from this plant. Meanwhile, compounds (**1-7**) were tested for their cytotoxicity against the MCF-7 and HepG2 human cancer cell lines. Herein, isolation, structural elucidation, as well as cytotoxicity of these diterpenoid alkaloids are reported. The hydrochloric acid extract of *A. franchetii* was alkalized with 28% NH₄OH soln. and then extracted with CH₂Cl₂ to give the total alkaloid. The alkaloid fraction was chromatographed over silica gel to afford two new C₁₉-diterpenoid alkaloids, villosudine A (**1**) and villosudine B (**2**), together with seven known analogues (**3-9**). The structures of the isolated compounds were as shown in Figure 1, and the ¹H and ¹³C NMR data of **1** and **2** were listed in Table. The known compounds, 6-epichasmanine (**3**),³ ezochasmaconitine (**4**),⁴ ezochasmanine (**5**),⁴ pseudoaconitine (**6**),⁵ ludaconitine (**7**),⁶ leucanthumsine A

(8),⁷ and vilmorisine (9)⁸ were established by comparison of their spectroscopic data with these reported in the literature.




Figure 1. Structures of compounds 1-9

Compound **1** was isolated as a white amorphous powder and gave a positive reaction to Dragendorff's reagent. Its molecular formula, C₃₅H₄₉NO₉ was deduced from the HR-ESI-MS m/z : 628.3488 [M+H]⁺, calcd. for 628.3486 and ¹³C NMR spectroscopic data. The IR (KBr) spectrum showed absorption bands for hydroxy group (3502 cm⁻¹) and ester carbonyl group (1713 cm⁻¹). The NMR data (Table) showed signals characteristic for an *N*-ethyl group [δ_{H} 1.10 (3H, t, J = 7.2 Hz), 2.50 (2H, overlapped); δ_{C} 48.7 t, 13.4 q], five methoxyl groups [δ_{H} 3.12, 3.23, 3.28, 3.29, and 3.52 (each 3H, s); δ_{C} 48.8 q; 56.0 q, 58.7 q, 58.9 q, and 59.2 q]. The quaternary carbon signals (δ_{C} 43.1 and 50.8) characteristic for C-4 and C-11, as well as an O-bearing methylene (δ_{C} 77.4, t) for C-18 suggesting compound **1** might be a typical C₁₉-diterpenoid alkaloid.⁹ In addition, resonances of a *trans*-double bond [δ_{H} 6.42 (1H, d, J = 16.0 Hz) and 7.68 (1H, d, J = 16.0 Hz); δ_{C} 118.5 d and 144.9 d], an aromatic ring [δ_{H} 7.36-7.53 (5H, m); δ_{C} 128.2×2, 129.0×2, 130.3 and 134.7], and a carboxyl group (δ_{C} 166.9) in the NMR spectra, along with the key HMBC correlations (Figure 2), H-9' (δ_{H} 7.68) to C-2' (δ_{C} 128.2) and C-7' (δ_{C} 166.9), suggested the presence of a *trans*-cinnamyl moiety.^{7,10}

The *trans*-cinnamyl moiety was located at C-14 by the long-range correlation in the HMBC experiment from H-14 [δ_{H} 4.76 (1H, d, J = 5.2 Hz)] to C-7' (δ_{C} 166.9). The existence of eight oxygenated carbons deduced from its HR-ESI-MS and ¹³C-NMR spectrum suggested **1** possess two hydroxy groups. A hydroxyl group could be placed at C-13 based on the multiplicity of H-14 and the δ value of 16-OCH₃ (δ_{H} 3.52, s).¹¹ Another hydroxy group was positioned at C-3, due to the HMBC correlations from H-3 (δ_{H}

3.78, m) to C-1 (δ_C 82.6), C-5 (δ_C 48.0), and C-18 (δ_C 77.4). The NMR data of compound **1** were very similar to those of villosutine,¹ a known alkaloid isolated from this plant, except for lacking a signal for acetyl group. The chemical shift of C-8 at δ_C 85.8 in villosutine was shifted upfield to δ_C 78.7 in compound **1** suggesting that 8-OAc was substituted by a methoxyl group,¹² which was confirmed by the difference of 28 mass units between those two compounds. The remaining methoxyl groups were assigned to at C-1, C-6, C-16, and C-18, due to the long-range correlations between 1-OCH₃ (δ_H 3.23)/C-1 (δ_C 82.6), 6-OCH₃ (δ_H 3.28)/C-6 (δ_C 83.1), 16-OCH₃ (δ_H 3.52)/C-16 (δ_C 83.7), and 18-OCH₃ (δ_H 3.29)/C-18 (δ_C 77.4) in HMBC spectrum (Figure 2). Accordingly, the substitution pattern and assigned planar structure of **1** was confirmed by complete ¹H-¹H COSY, HMQC, and HMBC spectroscopic analyses.

The relative configuration of compound **1** was deduced from the vicinal coupling constants (Table) and NOESY experiment (Figure 2). The coupling constant between H-5 and H-6 ($J = 6.4$ Hz) confirmed the β -position of H-6,¹³ which was further supported by the cross-peaks between H-6/H-9 β in the NOESY spectrum. The β -methoxyl group at C-16 was demonstrated by the correlations between the H-16 and H-17 α in the NOESY spectrum. Furthermore, the NOESY cross-peak between H-1/H-3, H-1/H-10 β and H-10 β /H-14 showed that H-1, H-3 and H-14 were β -oriented. Thus, the structure and absolute configuration of villosudine A (**1**) was determined as shown in Figure 1, and the full assignment of its spectroscopic data was achieved based on 1D and 2D NMR analyses.




Figure 2. Key ¹H-¹H COSY, HMBC (a) and NOESY (b) correlations of **1**

Compound **2** exhibited a pseudo-molecular-ion peak at m/z 672.3384 [$M+H$]⁺ in the HR-ESI-MS, corresponding to the molecular formula C₃₆H₄₉NO₁₁. Compound **2** exhibited characteristic NMR (Table) spectral features of an aconitine-type C₁₉-diterpenoid alkaloid¹⁴ bearing an *N*-ethyl group [δ_H 1.08 (3H, t, $J = 7.2$ Hz); δ_C 49.0 t, 12.6 q], four methoxyl groups [δ_H 3.26 and 3.50 (each 3H, s); 3.28 (6H, s); δ_C 55.4 q; 58.8 q, 58.9 q, and 59.1 q], and an acetyl group [δ_H 1.78 (3H, s); δ_C 22.3 q, 169.8 s]. The acetyl signal at higher field (δ_H 1.78) and the chemical shift of C-8 at δ_C 85.9 suggested that the acetyl group was

presented on C-8.⁷ Comparison of the NMR data of **2** with those of the known compound atropurpursine¹⁵ showed that the only difference between **2** and atropurpursine was a substitute group at C-14. The ¹H-NMR spectrum of **2** showed the presence of one CH=CH moiety at δ_{H} 6.58 (1H, d, $J = 16$ Hz) and 7.74 (1H, d, $J = 16$ Hz), together with those of five aromatic protons at δ_{H} 7.43-7.46 (3H, m) and 7.67-7.69 (2H, m), which were assigned to a *trans*-cinnamoyl group.^{7,10} The doublet signal at δ_{H} 4.77 ($J = 5.2$ Hz) could be assigned to H $_{\beta}$ -14, suggesting the presence of an ester function at C-14.¹⁴ The long-rang correlations between H-14 (δ_{H} 4.77) and H-9' (δ_{H} 7.74, d, $J = 16.0$ Hz) with the carbonyl carbon signal at δ_{C} 166.9 (s) in HMBC confirmed the presence of the *trans*-cinnamoyl group at C-14 (Figure 3). In addition, compound **2** possessed a similar configuration to that of **1**, according to the deduction from its NOESY experiment (Figure 3). Accordingly, the structure of villosudine B (**2**) was confirmed by extensive analysis of its NMR spectra.

Table. ¹H (400 MHz) and ¹³C NMR (100 MHz) Data for Compounds **1** and **2** (δ in ppm, J in Hz)

No.	1 in CDCl ₃		2 in (CD ₃) ₂ CO		No.	1 in CDCl ₃		2 in (CD ₃) ₂ CO	
	δ_{H}	δ_{C}	δ_{H}	δ_{C}		δ_{H}	δ_{C}	δ_{H}	δ_{C}
1	3.13 m	82.6 d	3.30 d (5.2)	83.8 d	19	a 2.43 d (12.4)	47.8 t	a 2.21 d (12)	46.5 t
2	a 1.90 m b 2.32 m	33.3 t	3.97 m	66.0 d	21	b 2.91 d (12.4)	48.7 t	b 2.66 d (12)	49.0 t
3	3.78 m	71.8 d	3.43 m	68.5 d	22	2.50 ^a	13.4 q	a 2.50 q (7.2)	12.6 q
4	—	43.1 s	—	44.7 s	1'	1.10 t (7.2)	134.7 s	b 2.70 q (7.2)	135.4 s
5	2.39 d (6.4)	48.0 d	2.35 d (6.4)	46.5 d	2'	—	—	—	129.0 d
6	4.01 d (6.4)	83.1 d	4.15 d (6.4)	83.9 d	3'	7.50-7.53 m	129.0 d	7.67-7.69 m	129.9 d
7	2.07 s	46.1 d	3.12 s	50.5 d	4'	7.36-7.38 m	130.3 d	7.43-7.46 m	131.3 d
8	—	78.7 s	—	85.9 s	5'	7.36-7.38 m	129.0 d	7.43-7.46 m	129.9 d
9	2.58 m	45.9 d	2.88 m	46.8 d	6'	7.36-7.38 m	129.0 d	7.43-7.46 m	129.0 d
10	2.03 ^a	41.4 d	2.28 m	41.3 d	7'	7.50-7.53 m	128.2 d	7.67-7.69 m	129.0 d
11	—	50.8 s	—	53.0 s	8'	—	166.9 s	—	166.9 s
12	a 2.03 ^a b 2.50 ^a	35.9 t	a 2.65 m b 2.09 m	38.8 t	9'	6.42 d (16.0)	118.5 d	6.58 d (16.0)	119.4 d
13	—	75.3 s	—	75.5 s	1-OMe	7.68 d (16.0)	144.9 d	7.74 d (16.0)	145.8 d
14	4.76 d (5.2)	79.1 d	4.77 d (5.2)	79.4 d	6-OMe	3.23 s	56.0 q	3.28 s	55.4 q
15	2.28 m	36.9 t	a 2.42 dd (6, 16) b 2.99 dd (8.8, 16)	40.5 t	8-OMe	3.28 s	58.7 q	3.28 s	58.8 q
16	3.38 t (8)	83.7 d	3.41 dd (6, 8.8)	84.6 d	16-OMe	3.12 s	48.8 q	—	—
17	2.77 br. s	61.2 d	2.82 br. s	60.6 d	18-OMe	3.52 s	58.9 q	3.50 s	59.1 q
18	a 3.48 ABq (8.8) b 3.58 ABq (8.8)	77.4 t	3.47 ABq (8.0) 3.63 ABq (8.0)	72.6 t	<u>-COMe</u> <u>-COMe</u>	3.29 s	59.2 q	3.26 s	58.9 q
						—	—	—	169.8 s
						—	—	1.78 s	22.3 q

^a Overlapped signals.




Figure 3. Key ^1H - ^1H COSY, HMBC (a) and NOESY (b) correlations of **2**

To evaluate the biological activities of these compounds isolated from the roots of *A. franchetii* for future applications, alkaloids **1-7** were tested for their *in vitro* cytotoxicity against the MCF-7 and HepG2 human cancer cell lines. Unfortunately, all of the compounds were inactive ($\text{IC}_{50} > 50 \mu\text{M}$, $n = 3$).

EXPERIMENTAL

General experimental procedure. Optical rotations were measured in CHCl_3 using a PerkinElmer polarimeter with a sodium lamp operating at 598 nm and 20 °C. The IR spectra were obtained using a Thermo Fisher Nicolet 6700 spectrometer. The HR-ESI-MS data were measured using a Q-TOF micro mass spectrometer (Waters). The NMR spectra were recorded on Bruker AV 400 spectrometer. The TLC plates were precoated with silica gel GF₂₅₄ (Qingdao Haiyang Chemical Co., Ltd., China), and it was visualized under a UV lamp at 254 nm or by spraying with Dragendorff's reagent or iodine.

Plant materials. *Aconitum franchetii* var. *villosulum* were collected in Baoxing, Sichuan province of China, in August 2014 and identified by Prof. Liang-ke Song of the Institute of Life Science and Engineering, Southwest Jiaotong University. A voucher specimen (No. SC20140805) was deposited in the School of Life Science and Engineering, Southwest Jiaotong University, Sichuan, China.

Extraction and isolation. Dried and powdered roots of *A. franchetii* var. *villosulum* (9.2 kg) were extracted with 0.1 mol/L hydrochloric acid seven times at room temperature, for two days each soaking. The combined solvent was adjusted to pH 9-10 with 28% NH_4OH soln., and then extracted with CH_2Cl_2 . The CH_2Cl_2 extracts were concentrated to produce the crude alkaloid extract (100 g). Column chromatography of the crude alkaloid extract over silica gel, using a CH_2Cl_2 : MeOH (100:1, v/v) mixture with increasing polarity afforded fractions A-F based on TLC analysis. Fraction C was separated by silica gel CC (petroleum ether: Me_2CO : Et_2NH 20:1:0.1, v/v/v) to obtain **2** (13 mg) and **9** (1 mg). CC (silica gel, petroleum ether: Me_2CO : Et_2NH 15:1:0.1, v/v/v) of fraction D afforded **7** (12 mg) and **8** (1 mg). Fraction E was chromatographed on silica gel column and eluted with petroleum ether: Me_2CO : Et_2N (10:1:0.1-0:1:0.1, v/v/v) to afford **1** (9 mg), **5** (5 mg) and **3** (6 mg). Fraction F was subjected to CC on

silica gel and eluted with petroleum ether: CH₂Cl₂ (1:1-0:1, v/v) to give **6** (6 mg) and **4** (8 mg).

Villosudine A (1)

White amorphous powder; $[\alpha]_{\text{D}}^{20} +10.38$ (*c* 0.55, CHCl₃); IR (KBr) ν_{max} 3502, 3060, 2963, 2931, 2889, 2821, 1713, 1638, 1578, 1496, 1450, 1385, 1335, 1309, 1281, 1202, 1176, 1089, 1043, 983, 921, 766, 709, 684, 491; ¹H NMR (400 MHz, CDCl₃) data and ¹³C NMR (100 MHz, CDCl₃) data, see Table; HR-ESI-MS (*m/z*): 628.3488 [M + H]⁺, calcd. for C₃₅H₅₀NO₉, 628.3486.

Villosudine B (2)

White amorphous powder; $[\alpha]_{\text{D}}^{20} +11.59$ (*c* 0.55, CHCl₃); IR (KBr) ν_{max} 3503, 3060, 2971, 2927, 2890, 2820, 1716, 1637, 1577, 1495, 1450, 1369, 1277, 1228, 1175, 1128, 1098, 985, 964, 921, 880, 841, 769, 709, 685, 620, 498; ¹H NMR (400 MHz, (CD₃)₂CO) data and ¹³C NMR (100 MHz, (CD₃)₂CO) data, see Table; HR-ESI-MS (*m/z*): 672.3384 [M + H]⁺, calcd. for C₃₆H₅₀NO₁₁, 672.3384.

Cell lines and cell culture and cytotoxicity assay

The *in-vitro* growth inhibitory activities of compounds **1-7** were assayed by the MTT method.¹⁶ The HepG2 (human hepatic carcinoma) and MCF-7 (human breast cancer) cell lines were obtained from ATCC. Cells were plated in 96-well plates 24 h before treatment and continuously exposed to different concentrations of compounds for 72 h. DMSO (0.1%, v/v) was used as the negative controls and adriamycin (≥ 98%; Sigma Chemical Co., Ltd., Shanghai, China) was used as the positive control.

ACKNOWLEDGMENTS

This research was supported by grants from NSFC (81773605), the Fundamental Research Funds for the Central Universities (2682017QY04), the Opening Project of Chemical Synthesis and Pollution Control Key Laboratory of Sichuan Province, and the Science and Technology Program of Sichuan, China (2018JY0077).

REFERENCES

- (a) W. Xu, L. Shan, S. Huang, S. Li, and X. Zhou, *Chin. J. Org. Chem.*, 2016, **36**, 2739; (b) J.-F. Zhang, L. Chen, S. Huang, L.-H. Shan, F. Gao, and X.-L. Zhou, *J. Nat. Prod.*, 2017, **80**, 3136.
- (a) F.-P. Wang, Q.-H. Chen, and X.-Y. Liu, *Nat. Prod. Rep.*, 2010, **27**, 529; (b) F. O. A. Frejat, W. Xu, L. Shan, and X. Zhou, *Heterocycles*, 2017, **94**, 1903; (c) C. Li, Y. Hirasawa, H. Arai, H. A. Aisa, and H. Morita, *Heterocycles*, 2010, **80**, 607; (d) J. Zhang, L. Chen, and X. Zhou, *Heterocycles*, 2018, **96**, 304.
- Z.-B. LI and F.-P. Wang, *Chin. Chem. Lett.*, 1996, **7**, 443.
- H. Takayama, M. Ito, M. Koga, S.-i. Sakai, and T. Okamoto, *Heterocycles*, 1981, **15**, 403.
- A. Klásek, V. Simánek, and F. Santavý, *Lloydia*, 1972, **35**, 55.

6. S. L. Shyaula, T. Tamang, N. Ghouri, A. Adhikari, S. Marasini, G. B. Bajracharya, M. D. Manandhar, and M. I. Choudhary, [Nat. Prod. Res.](#), 2016, **30**, 2590.
7. H. Yan, D.-L. Chen, X.-X. Jian, and F.-P. Wang, [Helv. Chim. Acta](#), 2007, **90**, 1133.
8. L. Ding, Y. Chen, and F. Wu, *Acta Chim. Sinica*, 1992, **50**, 405.
9. H. Ji, D.-L. Chen, and F.-P. Wang, [Heterocycles](#), 2004, **63**, 2363.
10. (a) D.-L. Chen, L.-Y. Lin, Q.-H. Chen, X.-X. Jian, and F.-P. Wang, [J. Asian Nat. Prod. Res.](#), 2003, **5**, 209; (b) L.-Y. Li, Q.-H. Chen, X.-L. Zhou, D.-L. Chen, F.-P. Wang, and C.-T. Che, [J. Nat. Prod.](#), 2003, **66**, 269; (c) C.-S. Peng, F.-P. Wang, and X.-X. Jian, [J. Asian Nat. Prod. Res.](#), 2000, **2**, 245; (d) R. Wang and Y.-Z. Chen, [Planta Med.](#), 1987, **53**, 544.
11. (a) S. Zheng, L. Gao, X. Hao, X. Wang, and X. Shen, [Phytochemistry](#), 1997, **46**, 951; (b) S. W. Pelletier, N. V. Mody, R. S. Sawhney, and J. Bhattacharyya, [Heterocycles](#), 1977, **7**, 327.
12. F.-P. Wang, *J. Org. Chem.*, 1982, **3**, 161.
13. D. Csupor, P. Forgo, K. Csedő, and J. Hohmann, [Helv. Chim. Acta](#), 2006, **89**, 2981.
14. S. W. Pelletier, N. V. Mody, B. S. Joshi, and L. C. Schramm, 'Alkaloids: Chemical and Biological Perspectives', Vol. 2, ed. by S. W. Pelletier, John Wiley Press, New York, 1984, pp. 205-462.
15. P. Tang, D. L. Chen, X. X. Jian, and F. P. Wang, [Chin. Chem. Lett.](#), 2007, **18**, 704.
16. S. Huang, J. Zhang, L. Shan, Y. Zhang, and X. Zhou, [Heterocycles](#), 2017, **94**, 485.