

HETEROCYCLES, Vol. 102, No. 1, 2021, pp. 81 - 91. © 2021 The Japan Institute of Heterocyclic Chemistry
Received, 2nd October, 2020, Accepted, 31st October, 2020, Published online, 18th November, 2020
DOI: 10.3987/COM-20-14363

PROBING THE REACTION OF N,N-DIMETHYFORMAMIDE DIMETHYL ACETAL WITH *p*-tert-BUTYLTHIACALIX[4]ARENE TETRAHYDRAZIDES: A FACILE APPROACH FOR THE SYNTHESIS AND APPLICATION OF NOVEL METAL RECEPTORS

Omran A. Omran^a and Ayman Nafady^{b*}

^aChemistry Department, Faculty of Science, Sohag University, 82524, Sohag, Egypt. E-mail: omran3646@gmail.com

^bDepartment of Chemistry, College of Science, King Saud University, Riyadh 11451, Saudi Arabia. E-mail: anafady2004@yahoo.com

Abstract – Recent developments in the field of sensing and recognition revealed that artificial receptors based on calix[4]arenes with π -conjugated fluorophoric or chromophoric moieties are efficient in enhancing selectivity/sensitivity as well as the binding affinity for ionic or molecular recognition. Herein, via studying the reaction of dimethylformamide dimethyl acetal (DMF–DMA) with *p*-tert-butylthiacalix[4]arene tetrahydrazides in three different conformers (i.e., *cone*, *partial-cone*, and *1,3-alternate*), the formation of novel lower rim *p*-tert-butylthiacalix[4]arene derivative (*cone*, *partial-cone*, and *1,3-alternate*) conformers in high yields (80, 95, and 85%, respectively) was achieved. The structure of isolated products was confirmed using different spectroscopic and analytical techniques such as FT-IR, ¹H NMR, ¹³C NMR, and elemental analysis. Importantly, the synthesized receptors showed preferential uptakes of alkali (Na⁺, K⁺ and Cs⁺), heavy (Pb²⁺, Cd²⁺, Hg²⁺, and Ag⁺) and transition (Ni²⁺, Co²⁺, and Cu²⁺) metal ions via liquid–liquid extraction method. The finding of this work adds new compounds to the growing family of thiacalix[4]arenes and validates their high sensitivity and selectivity in detection of heavy metal ions.

INTRODUCTION

Calix[n]arenes¹⁻⁴ are bowl-shaped macrocyclic compounds that have been widely used in catalysis,⁵ molecular recognition,⁶ metal ion receptors,⁷ and sensors.⁸ Among the calixarene family, thiacalix[4]arenes⁹⁻¹¹ are the most popular members as the presence of sulfur atoms in their structure

makes them ideal compounds for facile complexation with various metal ions. In this respect, metal-recognition abilities of thiacalixarenes can be tuned by introducing different functional groups to the active sites in their structures including the upper¹²⁻¹⁴ / lower rims or bridging sulfur atoms.^{15,16} However, the hydroxyl's lower rim groups are considered to be the easiest sites for preparing a large number of thiacalixarenes with various functional groups in different conformers.¹⁷

Thiacalixarenes lower rim functionalization is carried out mainly through alkylation with halo-reagents.¹⁸⁻²¹ In a previous work, alkylation using halo-reagent enabled the synthesis of three different conformers (i.e., *cone*, *partial-cone* and *1,3-alternate*) at the lower rim of *p-tert*-butylthiacalix[4]arene tetraesters,¹⁸ tetraethers,¹⁹ tetraketones,²⁰ and tetraamides.²¹ Among these conformers, *p-tert*-butylthiacalix[4]arene tetraesters were the best precursors for further functionalization of thiacalixarenes to synthesize tetra-acids,²² tetrathioesters,²³ and tetrahydrazides.²⁴ In addition, the *p-tert*-butylthiacalix[4]arene tetrahydrazides have been successfully used for the synthesis of other thiacalixarene derivatives through coupling with carbonyl compounds as well as addition to unsaturated carbon–carbon double bond or carbon–nitrogen triple bonds.²⁵⁻²⁸

As a continuation of our ongoing interest in thiacalixarenes and their potential applications in voltammetric sensors and receptors,²⁵⁻²⁸ we report herein the synthesis of lower rim derivatives, in three different conformers, by reacting N,N-dimethylformamide dimethyl acetal (DMF-DMA) with *p-tert*-butylthiacalix[4]arene tetrahydrazides. Moreover, the affinity of the prepared compounds to recognize and bind to different metal ions such as alkali (Na⁺, K⁺ and Cs⁺), heavy (Pb²⁺, Cd²⁺, Hg²⁺ and Ag⁺) and transition (Ni²⁺, Co²⁺ and Cu²⁺) metal ions using liquid–liquid extraction protocol in picrate solutions has been evaluated.

RESULTS AND DISCUSSION

Novel *p-tert*-butylthiacalix[4]arene derivatives **4a-c**, in three different conformers (i.e., *cone*, *partial-cone* and *1,3-alternate*) as shown in Scheme 2, were prepared through three steps. Firstly, following Miyano et al.¹⁸ who have selectively converted *p-tert*-butylthiacalix[4]arene **1** to *p-tert*-butylthiacalix[4]arene tetraesters **2a-c** in three different conformers (*cone*, *partial-cone*, and *1,3-alternate*) using alkali metal carbonates (M₂CO₃, M = Na, K, and Cs). Secondly, hydrazinolysis of different tetraesters conformers produced the corresponding *p-tert*-butylthiacalix[4]arene tetrahydrazides **3a-c**.²⁴ Finally, *p-tert*-butylthiacalix[4]arene tetrahydrazides **3a-c** were reacted with N,N-dimethylformamide dimethyl acetal (DMF-DMA) in boiling dry toluene to afford *p-tert*-butylthiacalixarene derivatives **4a-c** in good yields (80 to 90%) as illustrated in Scheme 1.


The proposed structures (Scheme 2) of the prepared thiacalixarene derivatives **4a-c** were confirmed by various spectroscopic and analytical methods including FT-IR (Supplementary Materials Figures S1-S3),

^1H NMR (S4-S6), and ^{13}C NMR (S7-S9) as well as elemental analysis. As can be seen in Figures S1-S3, IR spectra obtained for compounds **4a** - **4c** showed the appearance of characteristic IR bands at $\nu = 3378$, 3395, and 3400 cm^{-1} assigned for free NH group, along with strong bands at $\nu = 1701$, 1674 and 1676 cm^{-1} for CO group in the three compounds, respectively. Additionally, the compounds exhibited IR band at 1270 , 1265 , and 1263 cm^{-1} , which are attributed to (COC) group, respectively.


The fact that symmetrical structures of *cone* **4a** and *1,3-alternate* **4c** conformers are originated from their corresponding *cone* **3a** and *1,3-alternate* **3c** conformers, respectively; it is expected to exhibit the same number of peaks in NMR spectra. However, it is possible to differentiate between these conformers by considering the chemical shifts associated with different proton groups. Thus, as illustrated in Figures S4 and S6, the protons of the $-\text{OCH}_2\text{C}(\text{O})-$ group in the *1,3-alternate* conformation, are located in the shielding area of two adjacent benzene rings, and consequently the signals of these protons were detected in a higher field than the signals of the corresponding protons in cone conformer (4.21 ppm and 4.88 ppm, respectively). Whereas, the *tert*-butyl protons of the *1,3-alternate* conformer are noted in the deshielding zone of the attached phenyl unit and the two adjacent benzene rings. Thus, the *tert*-butyl protons signal of *1,3-alternate* should appear in a lower field compared to *tert*-butyl protons signals of the cone conformer (1.20 ppm and 1.11 ppm, respectively). In view of these results, the structures of compounds **4a** and **4c** can be unambiguously assigned to *cone* and *1,3-alternate* conformers, respectively.

The structure of compound **4b** is predicted to be unsymmetrical and therefore, the ^1H NMR spectrum (S5) exhibited marked increase in splitting and multiplicity of protons from different group. The obtained ^1H NMR spectrum of compound **4b** showed the presence of three non-equivalent peaks associated with the *tert*-butyl groups; and the $-\text{N}(\text{Me})_2$ group should exhibit a 2:1:1 ratio of resonance patterns in the proton spectrum. Thus, *tert*-butyl has the following peaks $\delta = 1.15$ (s, 18H), 1.29 (s, 9H), 1.36 (s, 9H) whereas peaks at $\delta = 2.84$ (s, 3H), 2.92(s, 3H), 2.96(s, 6H) are assigned to $-\text{N}(\text{Me})_2$ group. In addition, there are three peaks for $\text{OCH}_2\text{C}(\text{O})-$ protons at $\delta = 4.69$ (s, 2H), 4.8(s, 2H), 4.82(d, 2H); similar peaks are observed for Ar-H and $-\text{NH}$ protons.

In consistence with ^1H NMR results, ^{13}C NMR spectra (Figures S7-S9) also confirmed the symmetrical structure of both **4a** (*cone*) and **4c** (*1,3-alternate*) with the detection of 10 peaks in their spectra. As for the unsymmetrical structure, compound **4b** (*partial-cone*) showed similar number of peaks but with the splitting of each peak in the spectrum of the unsymmetrical structure **4b** (*partial-cone*) (see EXPERIMENTAL).



Scheme 1. Reagents and conditions: i, [7](#); ii, [13](#); and iii, DMF-DMA, toluene, reflux



Scheme 2. Structures of stereoisomers **4a-c**

Although the proposed structures of *p*-*tert*-butylthiacalix[4]arene derivatives **4a-c** contained at their lower rim various functional groups such as C=O, C=N, and NH, which is expected to easily coordinate with the desired metal ion, the conditions used in this study (acidic aqueous/organic) mixture (pH 4.5) are most likely to favor the $-C=N$ group to interact with the metal. Thus, To evaluate the ability of *p*-*tert*-butylthiacalix[4]arene derivatives **4a-c** to bind/host metal ions, a liquid–liquid extraction experiment from aqueous/organic biphasic system (i.e., $H_2O/CHCl_3$ mixture) was carried out using different metal picrate salts with the metal being either monovalent ($M = Na^+$, K^+ , and Cs^+) or divalent heavy and transition metals (Pb^{2+} , Cd^{2+} , Hg^{2+} , Ni^{2+} , Co^{2+} , Cu^{2+} , and Ag^+). The prepared sample solutions were analyzed via UV–Vis spectroscopy. The extraction percent ($E\%$) for alkali and transition metal ions

by using the studied derivatives of *p-tert*-butylthiacalix[4]arene are listed in Table 1 and Figure 1. It is observed that all *p-tert*-butylthiacalixarene derivatives **4a–c**, unlike tertiary amide analogues,²¹⁻²³ are good extractants for all alkali metal ions. However, *1,3-alternate* conformer **4c** is highly selective for K⁺ compared to other alkali metals with a percent of 52% is determined for K⁺ detection. This finding is unusual given that the analogous *p-tert*-butylthiacalixarene tetrahydrazides,²⁴ are only suitable for detection of transition metal ions. Moreover, it can be noted that *p-tert*-butylthiacalixarene derivatives **4a–c** are also highly selective extractants for Cd²⁺, Hg²⁺, and Cu²⁺ by *cone*, **4a** (59-75%); Pb²⁺, Cd²⁺, and Ag⁺ by *partial-cone* **4b** (68-80%). In view of these results, it is clearly established that these conformers have higher selectivity for the environmentally harmful transition metal cations such as Cd²⁺, Hg²⁺, and Ag⁺.

Table 1. Competition between extraction extent (*E*) for different metal ions by *p-tert*-butylthiacalix[4]arene derivatives **4a–c**

Ligand	Na ⁺	K ⁺	Cs ⁺	Pb ²⁺	Cd ²⁺	Hg ²⁺	Ni ²⁺	Co ²⁺	Cu ²⁺	Ag ⁺
4a	33.4	5.2	0.0	47.0	59.6	69.1	52.8	14.9	74.9	52.2
4b	4.3	52.2	0.5	80.0	68.0	5.8	24.0	19.0	37.9	75.1
4c	17.7	24.0	12.9	22.1	30.7	22.0	20.1	22.1	42.4	30.7





Figure 1. Extraction percent (*E*) of different metal ions using *p-tert*-butylthiacalix[4]arene derivatives **4a–c**

The protocol used for extraction of various metal ions involve using aqueous solution phase of metal picrate with concentration of 1.0×10^{-4} M and organic solution phase of the *p-tert*-butylthiacalix[4]arene derivatives **4a–c** (1.0×10^{-3} M) in chloroform. The extraction percentage is estimated using the formula $E\% = (\text{initial aqueous cation concentration}) - (\text{final aqueous cation concentration}) / (\text{initial aqueous cation concentration}) \times 100$.

The stoichiometry and extraction constants of the instantly formed ligand–metal complexes were also determined. Extraction equilibrium is described by Eq. (1), where M^{n+} , L, Pic^- , ML^{n+} , and $ML^{n+}nPic^-$ are the metal ion, ligand (here, **4a**, **4b** and **4c**), picrate anion, metal complex and ion-pair, respectively, while n is the number of ligands that react with one metal ion.



The percentage extraction, $E\%$, and extraction degree, α , were calculated according to Eq. (2).

$$E\% = \alpha \times 100\% = \frac{[ML_n^{n+}nPic^-]_{org}}{[Pic^-]_{aq,init}} \times 100 \quad (2)$$

The thermodynamic extraction constant, K_{ex} , is given by Eq. (3).

$$K_{ex} = [ML_n^{n+}nPic^-]_{org} \cdot f_{M,L,Pic^-} [M^{n+}] \cdot f_{M^{n+}} [nPic^-]_{aq} \cdot f_{pic^-} [L]_{org}^n f [L] \quad (3)$$

Where f is the activity coefficient. The activity coefficients of different species and ion-pair for diluted solution were 1.0×10^{-4} M; thus, they were neglected and assumed to be unity.

To determine the stoichiometric coefficient (n) of the complexes, which is formed in the organic phase using Eq. (4) as the modified form of Eq. 3.

$$\text{Log } K_{ex} = \log\left(\frac{\alpha}{1-\alpha}\right) - \log[M^{n+}]_{aq} - n \log[L]_{org} \quad (4)$$

Virtually, the plot of $\log(\alpha/1 - \alpha)$ versus $\log[L]_{org}$ gave rise to a straight lines, and its slope equals to the (n) value. Furthermore, the extraction constants (K_{ex}) are calculated using the intercept values (b).

$$b = \log K_{ex} + \log[M^{n+}]_{aq} \quad (5)$$

Our obtained results demonstrated that plotting of $\log(\alpha/1 - \alpha)$ vs. $\log[L]_{org}$ for selected metal ions that gave highest extraction percentage such as Ag^+ , Cd^{2+} , and Hg^{2+} (Figure 2) in saturated water/chloroform mixture was linear in all cases, therefore, Eq. 4 is valid, with the values of $\log K_{ex}$, K_{ex} and n are depicted in Table 2.




Figure 2. Plot of $\log(\alpha/1 - \alpha)$ vs. $\log[L]_{\text{org}}$ for metal picrate extraction by *p-tert*-butylthiacalixarenes **4a–c** in a mutually saturated water–chloroform solvent system

Table 2. Extraction constants $\log K_{\text{ex}}$ and stoichiometry complexes for different metal- ligand complexes formed in the organic phase

Ligand	Metal	$\log K_{\text{ex}}$	K_{ex}	n
4a	Cd^{2+}	10.38	2.39×10^{10}	1.5
4b	Ag^+	11.92	8.31×10^{11}	1.8
4c	Hg^{2+}	8.95	8.90×10^8	1.0

The obtained results (Table 2) revealed that the stoichiometric calculations associated with formation of metal-thiacalixarene complexes are highly dependent on the type of *p-tert*-butylthiacalixarenes (**4a–c**) conformers as well as the size and valance of metal ion. Thus, the ligand: metal (L:M) ratio in these type of host-guest associations does not follow a specific trend. In this respect, we found in the case of *cone* conformer (**4a**) with Cd^{2+} $n = 1.5$, thereby implying L:M ratio of 3:2. Whereas for *partial-cone* **4b** with Ag^+ ($n = 1.8$), the L: M ratio is $\sim 1: 2$ and for *1,3-alternate* conformer (**4c**) with Hg^{2+} a 1:1 stoichiometric ratio of L: M is accomplished. The unusual finding of 3:2 L:M ratio in case of cadmium can be explained in terms of the fact that the **4a** conformer of the ligand has four $-\text{C}=\text{N}-$ (azomethine) groups available for host-guest association with the metal ion, and Cd^{2+} favors an octahedral geometry that required the association of six $-\text{C}=\text{N}$ groups. In order to achieve such stoichiometry three molecules of **4a** have to react with two Cd^{2+} ions, thus, giving rise to a 3:2 ratio.

CONCLUSION

A facile synthesis of novel *p-tert*-butylthiacalixarene derivatives in three different conformers at the lower rim is achieved by the reaction of thiacalixarene tetrahydrazides with dimethylformamide dimethyl acetal. The ability of new thiacalixarene derivatives to recognize/bind alkali metal ions (Na^+ , K^+ , and Cs^+), heavy metals (Pb^{2+} , Cd^{2+} , and Hg^{2+}), and transition metal ions (Ni^{2+} , Co^{2+} , Cu^{2+} , and Ag^+) has been studied by the liquid–liquid extraction method. This study revealed that these new compounds are very sensitive and extremely selective towards a wide range of metal ions. Furthermore, given that these new compounds containing good leaving $-\text{N}(\text{Me})_2$ group at their lower rim, it can be used as a starting precursors for various transformations in organic/organometallic reaction processes.

General method for picrate extraction:

Solutions of metal picrates were prepared from an aqueous solution of picric acid and metal hydroxide (i.e., NaOH, KOH or CsOH) or metal nitrate [i.e., $\text{Pb}(\text{NO}_3)_2$, $\text{Cd}(\text{NO}_3)_2$, $\text{Hg}(\text{NO}_3)_2$, $\text{Ni}(\text{NO}_3)_2$, $\text{Co}(\text{NO}_3)_2$, $\text{Cu}(\text{NO}_3)_2$ or AgNO_3] solutions and analyzed by a pH meter. In the latter case, solutions were weakly acidic with the final pH of 4.5. An aqueous picrate solution (5 mL, 1.0×10^{-4} M) and a chloroform solution of the ligand (5 mL, 1.0×10^{-3} M) were stirred together for 0.5 h and then kept for 0.5 h at 25 °C. Blank experiments showed that metal picrate extraction did not occur in the absence of *p-tert*-butylthiacalixarene derivatives **4a-c**. The absorbance of aqueous phase before (A_0) and after extraction (A_i) was measured at 355 nm. The percent extraction ($E\%$) was calculated using the following expression:

$$E\% = \frac{A_0 - A_i}{A_0}$$

EXPERIMENTAL

All chemicals and reagents were of analytical grade and used as received without further purification. Melting points (°C, uncorrected) were determined using an open glass capillary *via* a Melting Point Apparatus (Stuart, model SMP10). The Fourier transform infrared (FT-IR) analysis of the prepared compounds was conducted on a Perkin Elmer Spectrum 1000 FT-IR Spectrometer. ^1H and ^{13}C NMR spectroscopic analyses were performed in deuterated chloroform (CDCl_3) and dimethyl sulfoxide (DMSO) at room temperature using JEOL ECP-400 MHz spectrometers. The chemical shifts were expressed in δ 7.28 downfield from CDCl_3 , which was used as a reference. Absorbance A was measured by a JENWAY, 640S UV - Vis spectrophotometer.

Synthesis of compounds 4a-c (general methods).

A mixture of *p*-*tert*-butylthiacalixarene tetrahydrazides **3a**, **3b** or **3c** (1 g, 0.99 mmol) with DMF-DMA (0.58 g, 4.95 mmol) in 40 mL dry toluene, was heated under reflux for 12 h. The mixture was concentrated to approximately 10 mL then left to cool. The white solid product was separated, filtrated and washed with MeCN. The product was recrystallized from EtOH.

4a, *cone*, yield 80%; white powder; Mp 170 °C, IR (KBr, ν/cm^{-1}): 3378 (NH-free), 1701 (CO), 1270 (COC). ^1H NMR (400 MHz, CDCl_3) δ : 1.11 (s, 36H, Bu^t), 2.96 (s, 24H, $\text{N}(\text{CH}_3)$), 4.88 (s, 8H, OCH_2CO), 7.38 (s, 8H, ArH), 8.02 (s, 4H, $\text{N}=\text{CH}$), 10.60 (s, 4H, CONH). ^{13}C NMR (100 MHz, CDCl_3) δ : 31.0, 34.2, 37.8, 75.9, 128.4, 135.0, 147.7, 158.1, 158.6, 164.2. Anal. Calcd for $\text{C}_{60}\text{H}_{84}\text{N}_{12}\text{O}_8\text{S}_4$: C, 58.61; H, 6.89; N, 13.67. Found: C, 58.50; H, 6.48; N, 13.35.

4b, *partial-cone*, yield 90%; white powder; Mp 242 °C. IR (KBr, ν/cm^{-1}): 3395 (NH), 1674 (CO), 1265 (COC). ^1H NMR (400 MHz, CDCl_3) δ : 1.15 (s, 18H, Bu^t), 1.29 (s, 9H, Bu^t), 1.36 (s, 9H, Bu^t), 2.84 (s, 3H, NCH_3), 2.92 (s, 3H, NCH_3), 2.96 (s, 6H, NCH_3), 4.62 (d, 2H, OCH_2CO), 4.69 (s, 2H, OCH_2CO), 4.8 (s, 2H, OCH_2CO), 4.82 (d, 2H, OCH_2CO), 7.41-8.42 (m, 8H, ArH + 4H, $\text{N}=\text{CH}$). ^{13}C NMR (100 MHz, CDCl_3) δ : 31.0, 31.2, 34.3, 34.5, 34.9, 37.8, 70.7, 73.4, 74.0, 125.2, 126.7, 128.1, 128.8, 129.0, 131.8, 133.5, 135.2, 135.7, 137.8, 147.9, 148.1, 156.8, 157.2, 157.6, 157.9, 158.1, 163.6, 164.0, 164.40. Anal. Calcd for $\text{C}_{60}\text{H}_{84}\text{N}_{12}\text{O}_8\text{S}_4$: C, 58.61; H, 6.89; N, 13.67. Found: C, 58.65; H, 6.54; N, 13.30.

4c, *1,3-alternate*, yield 80%; white powder; Mp 190 °C. IR (KBr, ν/cm^{-1}): 3400 (NH), 1676 (CO), 1263 (COC). ^1H NMR (400 MHz, DMSO) δ : 1.20 (s, 36H, Bu^t), 2.80 (s, 24H, $\text{N}(\text{CH}_3)$), 4.21 (s, 8H, OCH_2CO), 7.42 (s, 8H, ArH), 8.07 (s, 4H, $\text{N}=\text{CH}$), 8.38 (s, 4H, CONH). ^{13}C NMR (100 MHz, DMSO) δ : 31.2, 34.5, 37.5, 69.2, 127.2, 128.9, 147.9, 155.1, 157.9, 162.9. Anal. Calcd for $\text{C}_{60}\text{H}_{84}\text{N}_{12}\text{O}_8\text{S}_4$: C, 58.61; H, 6.89; N, 13.67. Found: C, 58.68; H, 6.51; N, 13.43.

ACKNOWLEDGEMENTS

The authors extend their appreciation for King Saud University, Riyadh, Saudi Arabia for funding this project through Researchers Supporting Project number (RSP-2020/79)

REFERENCES

1. P. Neri, J. L. Sessler, and M.-X. Wang, "[Calixarenes and Beyond](#)," Cham, Switzerland: Springer International Publishing; 2016.
2. C. D. Gutsche, Royal Society of Cambridge, "Calixarenes : an introduction," RSC Pub. Cambridge, 2008.
3. J. Vicens, L. Baklouti, and J. Harrowfield, "[Calixarenes in the Nanoworld](#)," Springer: Dordrecht, 2007.

4. Z. Asfari, V. Bohmer, and J. M. Harrowfield, ["Calixarenes 2001," Kluwer Academic Press: Dordrecht, 2001.](#)
5. R. Salvio, S. Volpi, T. Folcarelli, A. Casnati, and R. Cacciapaglia, [Org. Biomol. Chem., 2019, 17, 7482.](#)
6. G. J. Lumetta, R. D. Rogers, and A. S. Gopalan, In ["Calixarenes for Separations," Washington, DC, 2000, American Chemical Society: Washington, DC.](#)
7. L. Mandolini and R. Ungaro, ["Calixarenes in Action," Imperial College Press, London, 2000.](#)
8. V. M. Mirsky and A. K. Yatsimirsky, [Artificial receptors for chemical sensors, Wiley-VCH Verlag, Weinheim, Germany, 2011.](#)
9. R. Kumar, Y. O. Lee, V. Bhalla, M. Kumar, and J. S. Kim, [Chem. Soc. Rev., 2014, 43, 4824.](#)
10. N. Morohashi, F. Narumi, N. Iki, T. Hattori, and S. Miyano, [Chem. Rev., 2006, 106, 5291.](#)
11. P. Lhoták, [Eur. J. Org. Chem., 2004, 1675.](#)
12. O. Kasyan, E. Healey, A. Drapailo, M. Zaworotko, S. Cecillon, A. Coleman, and V. Kalchenko, [J. Incl. Phenom. Macrocycl. Chem., 2007, 58, 1.](#)
13. P. Lhoták, J. Morávek, and I. Stibor, [Tetrahedron Lett., 2002, 43, 3665.](#)
14. O. Kundrat, I. Cisarova, S. Böhm, M. Pojarova, and P. Lhotak, [J. Org. Chem., 2009, 74, 4592.](#)
15. O. Kundrat, P. Lhotak, V. Eigner, and H. Dvorakova, [Org. Lett., 2011, 13, 4032.](#)
16. T. Landovský, V. Eigner, M. Babor, M. Tichotová, H. Dvoráková, and P. Lhoták, [Chem. Commun., 2020, 56, 78.](#)
17. M. Simánová, H. Dvoráková, I. Stibor, M. Pojarová, and P. Lhoták, [Tetrahedron Lett., 2008, 49, 1026.](#)
18. N. Iki, F. Narumi, T. Fujimoto, N. Morohashi, and S. Miyano, [J. Chem. Soc., Perkin Trans. 2, 1998, 2745.](#)
19. P. Lhotak, M. Himl, S. Pakhomova, and I. Stibor, [Tetrahedron Lett., 1998, 39, 8915.](#)
20. I. I. Stoikov, O. A. Omran, S. E. Solovieva, S. K. Latypov, K. M. Enikeev, A. T. Gubaidullin, I. S. Antipin, and A. I. Konovalov, [Tetrahedron, 2003, 59, 1469.](#)
21. S. E. Solovieva, M. Gro'ner, A. O. Omran, A. T. Gubaidullin, I. A. Litvinov, W. D. Habicher, I. S. Antipin, and A. I. Konovalov, [Russ. Chem. Bull., 2005, 54, 2104.](#)
22. N. Iki, N. Morohashi, F. Narumi, and T. Fujimoto, [Tetrahedron Lett., 1999, 40, 7337.](#)
23. O. A. Omran and I. S. Antipin, [J. Incl. Phenom. Macrocycl. Chem., 2014, 78, 121.](#)
24. I. I. Stoikov, R. Z. Nasibullin, V. A. Smolentsev, L. I. Gafiullina, A. Y. Zhukov, J. B. Pupilampu, I. S. Antipin, and A. I. Konovalov, [Mendeleev Commun., 2006, 16, 248.](#)
25. A. O. Omran, [Chin. J. Chem., 2009, 27, 1937.](#)
26. S. N. Podyachev, N. E. Kashapova, V. V. Syakaev, S. N. Sudakova, R. R. Zainullina, M. Gruner, W.

- D. Habicher, T. A. Barsukova, F. Yang, and A. I. Konovalov, [*J. Incl. Phenom. Macrocycl. Chem.*, 2014, **78**, 371.](#)
27. O. A. Omran, F. A. Elgendy, and A. Nafady, [*Int. J. Electrochem. Sci.*, 2016, **11**, 4729.](#)
28. A. D. M. Mohamad and O. A. Omran, [*J. Iran. Chem. Soc.*, 2019, **16**, 2197.](#)