

HETEROCYCLES, Vol. 106, No. 3, 2023, pp. 479 - 487. © 2023 The Japan Institute of Heterocyclic Chemistry
 Received, 5th December, 2022, Accepted, 16th January, 2023, Published online, 24th January, 2023
 DOI: 10.3987/COM-22-14794

SELECTIVE SYNTHESIS OF 2,2-DIAMINO-4,4,6,6-TETRAKIS(ARYLOXY)CYCLOTRIPHOSPHAZENES N_3P_3 -2,2-(NH₂)₂-4,4,6,6-(ArO)₄

Manabu Kuroboshi,* Fumiya Nakamura, and Hideo Tanaka

Graduate School of Natural Science and Technology, Okayama University,
 Tsushima-naka 3-1-1, Kita-ku, Okayama, JAPAN, mkurohos@cc.okayama-u.ac.jp

Abstract – To synthesize cyclotriphosphazene derivatives having multi-functional groups, aryloxylation of 2,2-diamino-4,4,6,6-tetrachlorocyclotriphosphazene **2** was examined. A mixture of *gem*-disubstituted $N_3P_3(NH_2)_2(ArO)_2Cl_2$ **9_{gem}**, tri-substituted $N_3P_3(NH_2)_2(ArO)_3Cl$ **10**, and tetra-substituted $N_3P_3(NH_2)_2(ArO)_4$ **11** was obtained, especially **11** was obtained selectively when excess amount (6 equiv.) of ArONa was used. On the other hand, mono-substituted $N_3P_3(NH_2)_2(ArO)Cl_3$ **8** and *non-gem*-di-substituted $N_3P_3(NH_2)_2(ArO)_2Cl_2$ **9_{non-gem-cis}** and **9_{non-gem-trans}** were not detected.

Hexachlorocyclotriphosphazene (HCCP, $N_3P_3Cl_6$) has a flat six-membered ring in which three N atoms and three P atoms connect alternately, and two Cl atoms on each P atom.¹ The Cl-P bond of HCCP easily reacts with several kinds of nucleophiles. To prepare multi-functionalized cyclotriphosphazene derivatives for developing multi-functionalized materials,² partial substitution of HCCP would be an important method to introduce more than one kind of substituent (Figure 1).






Figure 1. Preparation of Cyclotriphosphazene Derivatives Having Multi-types of Nucleophiles

We previously reported that gaseous NH₃ was allowed to react with HCCP to afford 2,2-diamino-4,4,6,6-tetrachlorocyclotriphosphazene (**2**) (in Et₂O) and/or 2,2,4,4-tetraamino-6,6-dichlorocyclotriphosphazene


(**3**) (in MeCN), respectively, and **3** was converted to 2,2,4,4-tetraamino-6,6-bis(aryloxy)cyclotriphosphazene (**4**) by treatment with ArOH/K₂CO₃ (Scheme 1).³ HCCP was allowed to react with arylthiol (ArSH) in the presence of NEt₃ in MeCN to give 2,2-bis(arylthio)-4,4,6,6-tetrachlorocyclotriphosphazene (**5**, ArSH, NEt₃ = 2 equiv.) and 2,2,4,4-tetrakis(arylthio)-6,6-dichlorocyclotriphosphazene (**6**, ArSH, NEt₃ = 4 equiv.) (Scheme 2).⁴ In these reactions, nucleophiles were introduced in *gem*-manner, and *non-gem*-derivatives were not detected (Scheme 3). On the other hand, phenols were introduced in HCCP in *non-gem*-manner: *Non-gem-cis*-derivatives and *non-gem-trans*-derivatives were obtained as major isomers.⁵ We also reported the second substitution of **5**.⁶



Scheme 1. Synthesis of Aminochlorocyclotriphosphazenes **2** and **3**




Scheme 2. Synthesis of Arylthiochlorocyclotriphosphazenes **5** and **6**




Scheme 3. Synthesis of Di-substituted Tetrachlorocyclotriphosphazenes

In this paper, we described the second substitution of **2** with phenols **7** to give N₃P₃(NH₂)₂Cl_{4-n}(OAr)_n **9-11**.⁷ In this case, there are several problems in partial aryloxylation: (1) Is number of ArOH introduced controllable? (2) How is the regio/stereochemistry of di-substituted product **9**? *Gem* **9**_{gem}/*non-gem-cis* **9**_{non-gem-cis}/*non-gem-trans* **9**_{non-gem-trans}? (Scheme 4).

Scheme 4. Synthesis of 2,2-Diaminochloroaryloxycyclotriposphazenes **9-11**

Firstly, we examined solvent effect, since solvent played an important role in the reaction of HCCP with ammonia (Table 1).³ Compound **2** was treated with 1 equiv. of 4-chlorophenol (**7c**) and K_2CO_3 , revealing that acetone having highest (relative dielectric constant) value among 7 solvents gave the best result.⁸ Interestingly, only a mixture of di-substituted product **9_{gemC}**⁹ and **2** was obtained even 1 equiv. of **7c** was used, whereas mono-substituted **8c** was not detected. On the other hand, when HCCP was treated with 1 equiv. of sodium 4-chlorophenoxide, a mixture of HCCP, mono-substituted $\text{N}_3\text{P}_3\text{Cl}_5(\text{OC}_6\text{H}_4\text{-Cl-4})$,¹⁰ and *non-gem*-di-substituted products $\text{N}_3\text{P}_3\text{Cl}_4(\text{OC}_6\text{H}_4\text{-Cl-4})_2$ (a 1 : 1 mixture of *cis* and *trans*-isomers) was obtained, and *gem*-di-substituted isomer was not detected.¹¹


Table 1. Solvent Effect in the Reaction of **2** with 4-Chlorophenol in the Presence of K_2CO_3 

Entry	Solv	dielectric constant	Product ratio ^a		Entry	Solv	dielectric constant	Product ratio ^a	
			9_{gemC}	2				9_{gemC}	2
1	Toluene	2.3	7	93	5	acetone	20.5	22	78
2	Et ₂ O	4.3	n.d. ^b	100	6	PrCN	28.9	n.d. ^b	100
3	THF	7.7	4	96	7 ^c	MeCN	35.9	complex	n.d. ^b
4	2-butanone	18.5	n.d. ^b	100					

^aRelative product ratio was determined by ³¹P NMR of the reaction mixture. ^bNot detected. ^cA complex mixture not including **2** was obtained.

Next, we examined equiv. of phenols (Table 2). When **2** was treated with 6 equiv. of 4-hydroxyacetophenone (**7a**) with K_2CO_3 , a mixture of **10a** (tri-substituted) and **11a** (tetra-substituted products) was obtained and **2** and **9_{gem}a** (di-substituted) were not detected (Entry 1). When **7a**/NaH (6 equiv.) was used at 50 °C, only **11a** was obtained (Entry 2), whereas a mixture of **2** and di-substituted **9_{gem}a** was obtained and **10a** and **11a** were not detected when less than 3 equiv. of NaH was used (Entries 3, 4). Li and/or Na phenoxides (**7a'** and **7a''**, 6 equiv.) gave a mixture of **2**, **9_{gem}a**, **10a**, and **11a** (Entries 5-8). As a result, use of phenols (6 equiv.) and NaH (6 equiv.) in acetone at 50 °C gave tetra-substituted **11a** selectively.

Table 2. Reaction Conditions and Products Distribution




Entry	Temp	reagents	Product ratio ^a			
			2	9_{gem}a	10a	11a
1	rt	7a + K_2CO_3 (6.0/6.0 equiv.)	n.d. ^b	n.d. ^b	60	40
2	50	7a + NaH (6.0/6.0 equiv.)	n.d. ^b	n.d. ^b	n.d. ^b	100
3	50	7a + NaH (4.0/2.4 equiv.)	57	43	n.d. ^b	n.d. ^b
4	50	7a + NaH (3.0/1.8 equiv.)	71	29	n.d. ^b	n.d. ^b
5	-20	7a' (Li phenoxide, 6.0 equiv.)	100	n.d. ^b	n.d. ^b	n.d. ^b
6	rt	7a' (Li phenoxide, 6.0 equiv.)	24	68	n.d. ^b	n.d. ^b
7	50	7a' (Li phenoxide, 6.0 equiv.)	n.d. ^b	n.d. ^b	76	24
8	50	7a'' (Na phenoxide, 6.0 equiv.)	n.d. ^b	n.d. ^b	74	26

^aProduct ratio was determined by ^{31}P NMR of the reaction mixture.

^bNot detected.

Finally, we examined the aryloxylation with several phenols (Table 3). Phenols having electron-withdrawing groups (EWG: Entries 1-4) and electron-donating groups (EDG: Entries 5-6) gave tetra-substituted products **11**: a small amount of tri-substituted **10f** was detected with phenol having a strong EDG group (*p*-methoxyphenol (**3f**): Entry 6).

Table 3. Aryloxylation of 2,2-Diamimo-4,4,6,6-tetrachlorocyclotriphosphazene **2** with Phenols



Entry	ArOH (7)	Hammett parameter as <i>p</i> -substituent	ratio of 11 / 10 ^a	Isolated yield of 11 / 10 ^c
1	HO-C ₆ H ₄ -C(=O)Me- <i>p</i> (7a)	0.50	11a 100	62 ^c
2	HO-C ₆ H ₄ -C(=O)OMe- <i>p</i> (7b)	0.45	11b 100	58 ^c
3	HO-C ₆ H ₄ -Cl- <i>p</i> (7c)	0.23	11c 100	86 ^d
4	HO-C ₆ H ₄ -F- <i>p</i> (7d)	0.06	11d 100	84 ^d
5	HO-C ₆ H ₄ -Me- <i>p</i> (7e)	-0.17	11e 100	88 ^d
6	HO-C ₆ H ₄ -OMe- <i>p</i> (7f)	-0.27	11f 85 ^b	68 ^d


^aProduct ratio was determined by ³¹P NMR. ^b11% **10f** and 4% unidentified product.

^cIsolated by recrystallization. ^dIsolated by silica gel column chromatography.

In conclusion, aryloxylation of 2,2-diamino-4,4,6,6-tetrachlorocyclotriphosphazene **2** was examined. When 1 equiv. of ArOH/K₂CO₃ was used, a mixture of **2** and *gem*-di-substituted **9_{gem}** was obtained, whereas mono-substituted N₃P₃(NH₂)₂(OAr)Cl₃ **8** and *non-gem*-di-substituted **9_{non-gem-cis}** and **9_{non-gem-trans}** were not detected. On the other hand, tetra-substituted N₃P₃(NH₂)₂(OAr)₄ **11** and a small amount of tri-substituted N₃P₃(NH₂)₂(OAr)₃Cl **10** were obtained under conditions of phenols (6 equiv.), NaH (6 equiv.), acetone, 50 °C.

EXPERIMENTAL


Synthesis of 2,2-Diamino-4,4,6,6-tetrachlorocyclotriphosphazene (**2**):



Et₂O (200 mL) solution of HCCP (6.96 g, 20 mmol) was stirred for 20 h under NH₃ atmosphere at room temperature to give colorless precipitates. The reaction mixture was filtered, and the precipitates were washed with Et₂O (20 mL x 2). Then the precipitates were washed with hot MeCN (50 mL x 4), and the MeCN washings were concentrated under reduced pressure and dried in vacuo to give **2** (colorless solid, 5.97 g, 19.3 mmol, 96% yield).¹²

2,2-Diamino-4,4,6,6-tetrachlorocyclotriphosphazene (**2**); ³¹P NMR (162 MHz, CDCl₃) δ 21.63 (d, *J* = 50.8 Hz, 2P), 9.17 (t, *J* = 50.8 Hz, 1P).

Reaction of **2** with Aryloxides (Table 2)



2 was treated with ArOH and base under conditions summarized in Table 2. The reaction mixture was poured into sat. aq. K₂CO₃ and extracted with EtOAc. The combined organic layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was analyzed by ³¹P NMR.


2,2-Diamino-4,4-bis(4-chlorophenoxy)-6,6-dichlorocyclotriphosphazene (**9_{gemc}**); ³¹P NMR (162 MHz, CDCl₃) δ 24.16 (dd, *J* = 53.3, 68.1 Hz 1P), 16.31 (dd, *J* = 59.5, 68.1 Hz, 1P), 11.86 (t, *J* = 58.2 Hz, 1P).

2,2-Diamino-4,4-bis(4-acetylphenoxy)-6,6-dichlorocyclotriphosphazene (**9_{gema}**); ³¹P NMR (162 MHz, CDCl₃) δ 24.16 (dd, *J* = 53.3, 70.6 Hz, 1P), 15.93 (dd, *J* = 57.0, 66.9 Hz, 1P), 11.79 (t, *J* = 57.0 Hz, 1P).

2,2-Diamino-4,4,6-tris(4-acetylphenoxy)-6-chlorocyclotriphosphazene (**10a**); ³¹P NMR (162 MHz, CDCl₃) δ 21.02 (dd, *J* = 65.6, 83.0 Hz, 1P), 15.69 (t, *J* = 68.1 Hz, 1P), 6.36 (dd, *J* = 68.1, 83.0 Hz, 1P).

2,2-Diamino-4,4,6-tris(4-methoxyphenoxy)-6-chlorocyclotriphosphazene (**10f**); ³¹P NMR (162 MHz, CDCl₃) δ 22.74 (dd, *J* = 63.2, 76.8 Hz, 1P), 17.05 (t, *J* = 63.2 Hz, 1P), 9.20 (dd, *J* = 63.2, 76.8 Hz, 1P).

Synthesis of 2,2-Diamino-4,4,6,6-tetrakis(aryloxy)cyclotriphosphazene:



To an acetone (5 mL) solution of ArOH (**7**, 3 mmol) was added NaH (3 mmol) at room temperature, and the reaction mixture was stirred at room temperature for 0.5 h. To the resultant mixture was added **2** (0.5 mmol), and the resultant was stirred at 50 °C for 20 h. The reaction mixture was poured into sat. aq. K₂CO₃ and extracted with EtOAc (20 mL x 4). The combined organic layer was dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography or recrystallization to give the desired product.

2,2-Diamino-4,4,6,6-tetrakis(4-acetylphenoxy)cyclotriphosphazene (**11a**); ³¹P NMR (162 MHz, CDCl₃) δ 16.94 (t, *J* = 70.6 Hz, 1P), 8.29 (d, *J* = 70.6 Hz, 2P); ¹H NMR (400 MHz, CDCl₃) δ 7.84 (d, *J* = 8.7 Hz, 8H), 7.18 (d, *J* = 8.7 Hz, 8H), 2.55 (s, 12H); ¹³C NMR (100 MHz, CDCl₃) δ 196.83, 154.36, 134.20, 121.05, 26.64; ESI-MS (neg. mode) 742.1180 (M + Cl⁻), Calcd for C₃₂H₃₂Cl₅N₅O₈P₃ 742.1152.

2,2-Diamino-4,4,6,6-tetrakis(4-methoxycarbonylphenoxy)cyclotriphosphazene (**11b**); ³¹P NMR (162 MHz, CDCl₃) δ 16.95 (t, *J* = 70.6 Hz, 1P), 8.27 (d, *J* = 68.1 Hz, 2P); ¹H NMR (400 MHz, CDCl₃) δ 7.92 (d, *J* = 8.7 Hz, 8H), 7.16 (d, *J* = 8.7 Hz, 8H), 3.89 (s, 12H); ¹³C NMR (100 MHz, CDCl₃) δ 166.27, 154.35, 131.38, 131.38, 127.09, 120.96, 52.32; ESI-MS (neg. mode) 770.1188 (M-H⁻), Calcd for C₃₂H₃₁N₅O₁₂P₃ 770.1182.


2,2-Diamino-4,4,6,6-tetrakis(4-chlorophenoxy)cyclotriphosphazene (**11c**); ³¹P NMR (162 MHz, CDCl₃) δ 17.72 (t, *J* = 69.4 Hz, 1P), 10.10 (d, *J* = 69.4 Hz, 2P); ¹H NMR (400 MHz, CDCl₃) δ 7.21 (d, *J* = 9.2 Hz, 8H), 7.02 (d, *J* = 9.2 Hz, 8H); ¹³C NMR (100 MHz, CDCl₃) δ 149.27, 130.58, 129.58, 122.54; ESI-MS (neg. mode) 709.9159 (M + Cl⁻), Calcd for C₂₄H₂₀Cl₅N₅O₄P₃ 709.9171.

2,2-Diamino-4,4,6,6-tetrakis(4-fluorophenoxy)cyclotriphosphazene (**11d**); ³¹P NMR (162 MHz, CDCl₃) δ 17.62 (t, *J* = 66.9 Hz, 1P), 10.13 (d, *J* = 66.9 Hz, 2P); ¹H NMR (400 MHz, CDCl₃) δ 7.05 (dd, *J* = 9.2, 4.4 Hz, 8H), 6.93 (dd, *J* = 9.2, 8.2 Hz, 8H), ¹³C NMR (100 MHz, CDCl₃) δ 243.45 (d, *J* = 243.5 Hz), 146.68, 122.58 (d, *J* = 7.67 Hz), 116.08 (d, *J* = 24.0 Hz); ¹⁹F NMR δ from -117.48 to -117.7 (m); ESI-MS (neg. mode) 646.0395 (M + Cl⁻), Calcd for C₂₄H₂₀ClF₄N₅O₄P₃ 646.0353.

2,2-Diamino-4,4,6,6-tetrakis(4-methylphenoxy)cyclotriphosphazene (**11e**); ^{31}P NMR (162 MHz, CDCl_3) δ 17.94 (t, $J = 66.7$ Hz, 1P), 10.72 (d, $J = 66.7$ Hz, 2P); ^1H NMR (400 MHz, CDCl_3) δ 7.50-6.80 (m, 16H), 2.29 (s, 12H), ^{13}C NMR (100 MHz, CDCl_3) δ 148.84, 134.28, 129.88, 121.16, 20.90; ESI-MS (neg. mode) 630.1329 ($\text{M} + \text{Cl}^-$), Calcd for $\text{C}_{28}\text{H}_{32}\text{ClN}_5\text{O}_4\text{P}_3$ 630.1356.


2,2-Diamino-4,4,6,6-tetrakis(4-methoxyphenoxy)cyclotriphosphazene (**11f**); ^{31}P NMR (162 MHz, CDCl_3) δ 19.04 (t, $J = 65.6$ Hz, 1P), 11.52 (d, $J = 65.6$ Hz, 2P); ^1H NMR (400 MHz, CDCl_3) δ 7.02 (s, 8H), 6.74 (s, 8H), ^{13}C NMR (100 MHz, CDCl_3) δ 156.58, 144.54, 122.23, 114.37, 55.62; ESI-MS (neg. mode) 694.1140 ($\text{M} + \text{Cl}^-$), Calcd for $\text{C}_{28}\text{H}_{32}\text{ClN}_5\text{O}_8\text{P}_3$ 694.1152.

Synthesis of Lithium 4-Acetylphenoxide (**7a'**)



A mixture of 4-hydroxyacetophenone (**3a**, 1.36 g, 10 mmol), $\text{LiOH} \cdot \text{H}_2\text{O}$ (0.422 g, 10.1 mmol), and H_2O (25 mL) was stirred at room temperature for 30 min, and poured into toluene (100 mL). The mixture was evaporated at 1 atm to remove H_2O by azeotrope, and the residue was dried under reduce pressure to obtain lithium 4-acetylphenoxide (**7a'**).

Synthesis of Sodium 4-Acetylphenoxide (**7a''**)



A suspension of 4-hydroxyacetophenone (**3a**, 1.36 g, 10 mmol) and aq. NaOH (0.5 M, 20 mL, 10.0 mmol) was stirred at room temperature for 30 min, and poured into toluene (100 mL). The mixture was evaporated at 1 atm to remove H_2O by azeotrope, and the residue was dried under reduce pressure to obtain sodium 4-acetylphenoxide (**7a''**).

ACKNOWLEDGEMENTS

This study was financially supported by JSPS KAKENHI, Grant Number JP20K05665. HCCP was a gift from Otsuka Chemical Co Ltd. Mass spectra were measured by Mrs. Tsugumi SHIOKAWA, Department of Instrumental Analysis & Cryogenics, Advanced Science Research Center, Okayama University.

REFERENCES AND NOTES

1. L. W. Daasch and D. C. Smith, *Anal. Chem.*, 1951, **23**, 853; A. Wilson and D. F. Carroll, *Chem. Ind.* 1958, 1558; A. Wilson and D. F. Carroll, *J. Chem. Soc.*, 1960, 2548.
2. M. Kuroboshi and H. Tanaka, *Heterocycles*, 2020, **100**, 1513, and references cited herein.
3. M. Kuroboshi, M. Morita, Y. Masumoto, M. Mikasa, R. Toza, and H. Tanaka, *Heterocycles*, 2019, **98**, 931.
4. M. Kuroboshi, M. Misaki, and H. Tanaka, *Heterocycles*, 2019, **98**, 940.
5. Y. W. Chen-Yang and B. D. Tsai, *Polyhedron*, 1993, **12**, 59.
6. M. Kuroboshi, A. Ueno, A. Kawano, and H. Tanaka, *Heterocycles*, 2022, **104**, 69.
7. G. E. Peters, R. J. Radel, and R. Medina, *J. Agric. Food Chem.*, 1988, **36**, 384.
8. Nucleophilicity of OAr^- would high in polar solvent because solvation of counter cationic metal would occur efficiently. Dielectric constant was used as a measure of polarity of the solvent. Solubility of **2** (high in a polar solvent) would also have an effect on the reactivity.
9. PCl_2 of HCCP and **2** appeared at 20-24 ppm, whereas $\text{P}(\text{NH}_2)_2$ and $\text{PCl}(\text{OAr})$ appeared at 9.2 and 12-14 ppm (unpublished result) in ^{31}P NMR. Substituted product showed at 24.2 (PCl_2), 11.9 ($\text{P}(\text{NH}_2)_2$), and 16.3 ppm. Therefore, substituted product was deduced to be a *gem*-di-substituted species.
10. D. Dell, E. W. Fitzsimmons, and R. A. Shaw, *J. Chem. Soc.*, 1965, 4070.
11. Though the selectivity (HCCP gave *non-gem*-products, whereas **2** only gave **9_{gem}**) is a very interesting problem, the authors have no evidence for this problem in this stage. When mono-substituted compound **7** will be isolated and analyzed by X-ray crystallography and IR, some evidences will be obtained (bond length, vibration wave number, etc). Now, the authors are trying to make the 2,2-diamino-4-mono-phenoxyphosphazene derivatives.
12. Ether solution, hot MeCN washings, and remained precipitates included unreacted **1** (HCCP), **2**, and NH_4Cl , respectively.